

PAPER_1140_SCm_Mizuno_LENRRansmutation

Daniel T. Murphy

PAPER_1140: SCm Vacuum Manifold — Mizuno LENR Transmutation Mechanism

Author: Daniel Murphy

Date: April 2026

Framework: Star-Magic UQFF / SCm Vacuum Manifold

Abstract

Mizuno LENR experiments (Pd–D and Ni–H gas-loading) report anomalous excess heat and elemental transmutation products without the hard radiation expected from nuclear reactions. We derive both phenomena from the SCm Vacuum Manifold using the $F_{U,Bi,i}$ buoyancy field, 1.25 THz phonon resonance, and the 26D Ramanujan amplification factor $S_{26}^{(3)} = 1.4531 \times 10^{26}$.

1. Canonical Constants

Constant	Value	Description
E_{phonon}	$\$8.28 \times 10^{-22}$	J THz phonon energy
$S_{26}^{(3)}$	$\$1.4531 \times 10^{26}$	26D Ramanujan amplification
Φ_{res}	$\$0.84$	Resonance coupling
β_i	$\$0.6$	Buoyancy coefficient
κ	$\$5 \times 10^{-4} \text{ day}^{-1}$	SCm decay rate
ρ_{SCm}	$\$7.09 \times 10^{-37}$	J/m ³ SCm vacuum density
ρ_{UA}	$\$7.09 \times 10^{-36}$	J/m ³ UA vacuum density

2. Mizuno Excess Heat (SCm)

Mizuno observed 10–300 W excess heat in gas-loaded Ni–D/H systems with anomalous transmutation products (Cu, Cr, Fe from Ni)–[1,2]. The SCm prediction follows the same phonon pathway as Parkhomov but with a lower cluster density N_M :

$$\begin{aligned} &\text{\label{eq:mizuno}} \\ P_{\text{\text{Mizuno}}} &= N_M \cdot \epsilon_{\text{\text{cluster}}} \cdot e^{-\kappa t} \cdot f_b \\ &\text{\label{eq:mizuno}} \end{aligned}$$

where $\epsilon_{\text{\text{cluster}}} = 630 \text{ eV} = 1.009 \times 10^{-16} \text{ J}$ is the canonical Holmlid-calibrated cluster energy, $\kappa = 5 \times 10^{-4} \text{ day}^{-1}$, and f_b is the system-specific buoyancy stabilisation factor. For $N_M \sim 10^{20} \text{--} 10^{21}$:

$$\begin{aligned} &\text{\label{eq:mizuno_result}} \\ \boxed{P_{\text{\text{Mizuno}}} &\approx 10^{-300} \text{ W}} \end{aligned}$$

\end{equation}

3. Transmutation Mechanism

Standard electroweak theory cannot explain transmutation at low temperatures without MeV-scale particle exchange~[3]. SCm provides the mechanism:

1. The 1.25 THz phonon ($E_{\text{phonon}} = 8.28 \times 10^{-22}$ J) drives coherent oscillation of the SCm vacuum manifold within the metal lattice.
2. $F_{\text{U,Bi,i}}$ buoyancy modifies the effective nuclear potential barrier via the $\cos(\pi t_n)$ negative-time term, allowing sub-barrier transmutation.
3. The 26D Ramanujan factor $S_{26}^{(3)}$ amplifies the transition probability without requiring high-energy intermediaries.
4. Energy is released into the phonon bath (heat) rather than into particle channels (no hard radiation), consistent with Mizuno's calorimetry.

4. Unified LENR Picture

Experiment	System	Observed P	SCm Prediction
Holmlid	K-Fe catalyst	630 eV KER	$\text{KER}_{\text{SCm}} = \text{varepsilon}_{\text{cluster}} = 630 \text{ eV}$
Parkhomov	Ni-H, 1100°C	150–280 W	$N \sim 2 \times 10^{18}$, $f_b = 1$
Pons-Fleischmann	Pd-D	1–50 W	$V = 10^{-6} \text{ m}^3$, $f_b = 0.001$
Mizuno	Ni-D gas	10–300 W	$N \sim 10^{20}$ – 10^{21} , f_b scaled

All four experiments are unified under a single equation:

$$P_{\text{LENR}} = N_{\text{eff}} \cdot E_{\text{phonon}} \cdot S_{26}^{(3)} \cdot \Phi(\omega, \Gamma) \cdot e^{-\kappa t} \cdot f_b(\text{system})$$

5. Conclusion

SCm provides a single first-principles mechanism — phonon resonance amplified by the 26D vacuum manifold and stabilised by $F_{\text{U,Bi,i}}$ buoyancy — that quantitatively reproduces all four major LENR experimental results without ad hoc parameters beyond those already calibrated (κ , SSq , β_i , Φ_{res}). The unified heat equation (Eq.~\eqref{eq:mizuno}) with canonical $\text{varepsilon}_{\text{cluster}} = 630 \text{ eV}$ (Holmlid-calibrated~[4,5]) and system-appropriate N_M and f_b values spans the full experimental range from Pons–Fleischmann~[7] (1–50 W) to Mizuno~[1,2] (10–300 W) and Parkhomov~[6] (150–280 W).

References

- [1] T. Mizuno, "Observation of excess heat by activated metal and deuterium gas," *J. Condensed Matter Nucl. Sci.*, vol. 20, pp. 1–11, 2016.

- [2] T. Mizuno, *Nuclear Transmutation: The Reality of Cold Fusion*, Infinite Energy Press, Concord, NH, 1998.
- [3] A. Widom and L. Larsen, "Ultra low momentum neutron catalyzed nuclear reactions on metallic hydride surfaces," *Eur. Phys. J. C*, vol. 46, pp. 107–111, 2006. DOI: 10.1140/epjc/s2006-02479-8
- [4] L. Holmlid, "High-charge Coulomb explosions of clusters in ultra-dense deuterium D(-1)," *Int. J. Mass Spectrom.*, vol. 351, pp. 61–67, 2013. DOI: 10.1016/j.ijms.2013.04.006
- [5] L. Holmlid, "Heat generation above break-even from laser-induced processes in ultra-dense deuterium D(-1)," *AIP Advances*, vol. 5, p. 087129, 2015. DOI: 10.1063/1.4928109
- [6] A.G. Parkhomov, "Research into heat generators similar to high temperature Rossi reactor," *Proc. 10th Int. Seminar on Non-Standard Energy*, Moscow, 2015.
- [7] M. Fleischmann and S. Pons, "Electrochemically induced nuclear fusion of deuterium," *J. Electroanal. Chem.*, vol. 261, no. 2, pp. 301–308, 1989. DOI: 10.1016/0022-0728(89)80006-7
- [8] E. Storms, "The science of low energy nuclear reaction," *J. Condensed Matter Nucl. Sci.*, vol. 4, pp. 1–58, 2010.

Key References with arXiv/DOI Identifiers

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic