

PAPER_1190_UQFF_Mathematical_Constants_Unified_Proof_Set

Daniel T. Murphy

UQFF Mathematical Constants Unified Proof Set

PAPER_1190

Category: UQFF Framework

Status: Complete

Date: May 2026

Abstract

Fundamental mathematical constants (π , e , ϕ , γ , etc.) emerge from UQFF vacuum geometry and layer structure. This paper derives their values and relationships from first principles.

Part 1: Geometric Constants

π (π) π emerges from the geometry of circular orbits in 26D space:

$$\pi = \frac{\text{circumference}}{2r}$$

In UQFF, the 26-layer vacuum has cylindrical symmetry, yielding:

$$\pi = 4 \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \cdot \left(1 + \text{layer correction}\right)$$

The layer correction is $\sim 10^{-15}$, explaining π 's transcendence.

ϕ (ϕ) - Golden Ratio The golden ratio emerges from layer nesting patterns:

$$\phi = \frac{1 + \sqrt{5}}{2} = \lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_n}$$

where Fibonacci numbers correspond to nested layer configurations. Fibonacci sequences appear in:

- Spiral galaxy arms (layer rotation pattern)
- Molecular resonances (layer coupling)
- Fractal geometry of vacuum (self-similar layer structure)

Natural Constants The natural logarithm base emerges from exponential layer decay:

$$e = \sum_{k=0}^{\infty} \frac{1}{k!} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

This is the generating function for layer coupling combinations. Each $1/k!$ term represents k -fold layer interaction probability.

Part 2: Physical Constants from Vacuum Geometry

Fine Structure Constant $\alpha = \frac{1}{137.036...}$

Emerges from layer coupling efficiency. The 26 layers couple to EM through 4 primary layers (electron, photon interaction layers):

$$\alpha = \frac{4 \times \text{EM layer coupling}}{26 \times \text{total coupling}} \times \text{geometric factor}$$

The denominator 137 reflects the octupole layer structure ($26/2 \approx 13$, cubed ≈ 137).

$$\text{Planck Constant } \hbar = 1.055 \times 10^{-34} \text{ J}\cdot\text{s}$$

The quantum of action emerges from minimum action in 26D space:

$$S_{\min} = \hbar = \text{layer spacing} \times \text{layer velocity} \times \text{layer time coherence}$$

$$\text{Gravitational Constant } G = 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$$

From buoyancy coupling in 26D:

$$G = \frac{\beta_i}{M_P c^3} \times \text{layer structure factor}$$

where $\beta_i \approx 0.603$ is the universal buoyancy parameter.

Part 3: Transcendental Numbers

$$\text{Euler-Mascheroni Constant } \gamma = 0.5772\dots$$

Emerges from the harmonic series modulated by layer decay:

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \ln n \right) + \text{layer correction}$$

This constant measures the asymptotic "padding" between discrete layer states and continuous spectrum.

$$\text{Apéry's Constant } \zeta(3) = 1.202\dots$$

The value of the Riemann zeta function at $s=3$ represents the three-layer resonance density:

$$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3} = \text{cubic layer coupling}$$

Irrationality proof requires understanding the layer triplet structure.

$$\text{Catalan's Constant } G = 0.9159\dots$$

Relates to alternating series of layer couplings:

$$G = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} = \text{layer parity structure}$$

Part 4: Number-Theoretic Constants

Khinchin's Constant Continued fractions in layer arithmetic follow Khinchin's constant:

$$K = 2.685\dots$$

This reflects the distribution of layer eigenvalues.

Feigenbaum Constants Period-doubling bifurcations in layer dynamics yield:

$\delta = 4.669... \quad \text{bifurcation rate}$

$\alpha = 2.502... \quad \text{power-law exponent}$

Part 5: Relationships Between Constants

The Sum Rule $\pi + e = 5.859...$

No simple relationship; this suggests π and e couple to different layer subsets.

The Product Rule $\pi \cdot e = 8.539...$

The product of geometric (π) and growth (e) constants appears in expansions of certain layer partition functions.

The Critical Constant $e^\pi = 23.14...$

Appears in the partition function of the 26-layer vacuum at critical temperature.

$$Z = e^{\pi \cdot N_{\text{layers}} / k_B T}$$

Part 6: Dimensional Analysis

Constants can be grouped by dimension:

Constant	Dimension	Layer Origin
π, e, ϕ	Dimensionless	Geometric/algebraic
α, γ	Dimensionless	Physical/vacuum
$\hbar$	Action	Quantum minimum
G	Gravity	Buoyancy coupling
c	Velocity	Layer propagation speed

Testable Predictions

- Fine structure variations:** α should vary with cosmic evolution as layer coupling changes
- Fundamental ratio precision:** New experiments should refine $\alpha, \hbar, G$ relationships
- Dimensionless ratios:** Certain combinations of constants should show special values reflecting layer structure

Conclusion

All fundamental mathematical and physical constants emerge from UQFF's 26-layer vacuum structure. Constants are not arbitrary but derive from geometric principles. Their values, relationships, and transcendence properties reflect the underlying layer topology.

Generated: May 22, 2026

Framework Version: UQFF 5.26