

PAPER_1191_GW190425_Mass_Gap_Bayesian_UQFF

Daniel T. Murphy

■---

paper_id: PAPER_1191
title: "GW190425 Mass-Gap Bayesian Posterior: MC Error Bars on f_{gap} Under UQFF Aether Modulation"
session: 299
date: 2026-05-17
author: "Daniel T. Murphy"
status: production
cvw: "v2.0.0"
tags: ["GW190425", "mass-gap", "Bayesian", "Monte-Carlo", "UQFF", "LIGO"]
crosslinks: [PAPER_1185, PAPER_365]
sm_anchor: "CVW v2.0.0 -- G6 SM Anchor Gate compliant"
arxiv_class: "gr-qc"

PAPER_1191: GW190425 Mass-Gap Bayesian Posterior --- MC Error Bars on f_{gap} Under UQFF Aether Modulation

Abstract

We close audit gap \#9 of the UQFF calibration program by adding Bayesian error bars to the GW190425 mass-gap fraction. A Monte-Carlo posterior over chirp mass $\mathcal{M}_c$ and mass ratio q propagates LIGO observational uncertainty into the per-component masses (m_1, m_2) and returns $f_{\text{gap}} \equiv P(2 \leq m_1 \leq 5, \mathcal{M}_c \odot)$ with 16/84-percentile error bars. Validation on four anchors (GW190425, GW190814, GW170817, mock BBH) recovers published gap fractions; calculator registered as `cp4_id = 443`. 20/20 smoke tests pass.

1. Motivation

The previous UQFF treatment of GW190425 returned a point-estimate f_{gap} without uncertainty propagation, which is incompatible with the CVW v2.0.0 G6 SM Anchor Gate requirement of falsifiable error bars. Audit entry \#9 flagged this gap as MEDIUM priority.

2. Forward Model

For chirp mass $\mathcal{M}_c$ and mass ratio $q \equiv m_2/m_1 \in (0, 1]$, component masses are $m_1 \equiv \mathcal{M}_c \frac{(1+q)^{1/5}}{q^{3/5}}, m_2 \equiv q m_1$.

3. Monte-Carlo Posterior

We draw $N_{\text{MC}}=4000$ samples,

$$\mathcal{M}_c^{(k)} \sim \mathcal{N}(\mathcal{M}_c^{\text{obs}}, \sigma), \text{quad } q^{(k)} \sim \mathcal{U}(q_{\text{lo}}, q_{\text{hi}}).$$

The mass-gap fraction is

$$f_{\text{gap}} = \frac{1}{N_{\text{MC}}} \sum_{k=1}^{N_{\text{MC}}} \mathbb{1}_{\left[2, M_{\text{dot}} \leq m_1^{(k)} \leq 5, M_{\text{dot}}\right]}.$$

Per-component median and 16/84 percentiles are reported as $m_{1, \text{med}}$ with asymmetric error bars $\Delta_+ = m_{1, \text{hi}} - m_{1, \text{med}}$ and $\Delta_- = m_{1, \text{med}} - m_{1, \text{lo}}$.

4. UQFF Aether Correction

GW strain residuals carry a small $|\delta| \leq 10^{-3}$ Aether modulation $f_A = 1 + \beta_i (\rho_{\text{SCm}} / \rho_{\text{amb}}) \cos(\pi t_n)$, which does not affect f_{gap} at the precision of present LIGO observations.

5. Anchor Validation

Anchor	$\mathcal{M}_c$ (M_{dot})	σ	range	$f_{\text{gap}}^{\text{pred}}$	$f_{\text{gap}}^{\text{obs}}$	Match
A1 GW190425	1.44	0.02	[0.8, 1.0]	~ 0.05	[0, 0.10]	yes
A2 GW190814	6.09	0.06	[0.10, 0.13]	~ 0.65	[0.30, 1.00]	yes
A3 GW170817	1.188	0.001	[0.7, 1.0]	~ 0	≤ 0.05	yes
A4 mock BBH	28	1	[0.5, 1.0]	~ 0	≤ 0.05	yes

6. Code Registration

Module `_session299_gw190425_mass_gap_errors.py`; class `GW190425MassGapBayesianCalculator`; `cp4_id = 443`. Registered in `CondensedPhysics3.py` under `SESSION_299_CALCULATORS`. Smoke-test result: 20 / 20 (4/4 anchors match).

7. Falsifiable Predictions

- (i) For GW190425 the posterior median m_1 must lie in $[1.4, 2.5] M_{\text{dot}}$.
- (ii) For any future BBH with $\mathcal{M}_c > 20 M_{\text{dot}}$ the mass-gap fraction must be statistically zero (≤ 0.05).

8. Status

`[[CLOSED]]` quad audit gap `#9` quad `cp4_id = 443` quad session = 299.

9. References

Abbott et al. 2020, ApJ 892, L3 (GW190425). Abbott et al. 2020, ApJ 896, L44 (GW190814). UQFF_CALIBRATION_AUDIT.md.