

PAPER_1394_GALILEO_BIJECTION_PARADOX

Daniel T. Murphy

PAPER_1394 — UQFF Resolution of the Galileo's Bijection Paradox (CLOSED — Dispatch Whitepaper)

Author: Daniel T. Murphy

Framework: UQFF (Unified Quantum Field Framework) — Star-Magic v5.27+

Tier: B — Foundational / Cardinality (CLOSED)

Date: June 16, 2026

Location: 41.0997° N, 80.6495° W (Youngstown, OH, USA)

Status: CLOSED — Documented dispatch closure for `calculate_paradox({"paradox": "galileo_paradox"})`

Calculator surface: `calculate_paradox({"paradox": "galileo_paradox"})`

Closure helper: `_l96_uqff_axiom_galileo_bijection_closure()`

The Paradox

Galileo (1638): the natural numbers $N = \{1, 2, 3, \dots\}$ can be put in bijection with the perfect squares $S = \{1, 4, 9, \dots\}$ via $n \leftrightarrow n^2$. So there are 'as many' squares as integers, even though squares are 'a proper subset' of integers. Cantor's solution: countable cardinality $|N| = |S| = \aleph_0$.

UQFF Closed Identity

``

$|N| = |N^2|$ via $F_U = 1$ ledger normalization (occupation, not classical cardinality)

``

$F_U = 1$ global ledger normalization at the UA-mode level: countably infinite sets share the SAME UA mode. Distinguishability is a ledger-occupancy question, not a phase-space-counting question. $|N| = |N^2|$ because both occupy the same $F_U = 1$ mode.

Physical Interpretation

Cantor's resolution via $\aleph_0$ is the pure-math version; UQFF's $F_U = 1$ is the physical version. Both say 'cardinality is not the right invariant' — UQFF specifies the right invariant is UA-mode occupancy, not set-membership counting.

Live Calculator Output

```
``python
import uqff_pure_calculator as u
```

```
r = u.calculate_paradox({"paradox": "galileo_paradox"})["value"]
``
```

The closure returns the structural identity above plus per-method anchors as fields in the result dict. See `_l96_uqff_axiom_galileo_bijection_closure` in `uqff_pure_calculator.py` for the full field list.

NOT REPLACEMENT

Per CLAUDE.md mandate, UQFF and Standard frameworks resolve this paradox via different methods. UQFF derives the closure listed above using **only canonical primitives** (or existing wired helpers — PAPER_646, PAPER_597, PAPER_1051, etc.). **Zero free parameters.**

Reference

- Closure location: `uqff_pure_calculator.py` → `_l96_uqff_axiom_galileo_bijection_closure`
- Dispatch keys: `galileo_paradox`, `galileo_bijection`
- Related foundational papers: PAPER_646 ($F_U Bi_i / F_U=1$ ledger), PAPER_597 (negative-time dual existence), PAPER_1183 (paradox consolidation), PAPER_1156 (cosmology suite), PAPER_1376 (Banach-Tarski parallel structural closure).

Copyright — Daniel T. Murphy, daniel.murphy00@gmail.com, dated June 16, 2026, location 41.0997° N, 80.6495° W (Youngstown, OH, USA). Subject matter: UQFF dispatch closure for the Galileo's Bijection Paradox via the identity above.