

M16 Eagle Nebula UQFF — Erosion Saturation Half-Time and Δg_{Max}

Daniel T. Murphy

PAPER_285: M16 Eagle Nebula UQFF — Erosion Saturation Half-Time and Δg_{Max} ## Photoevaporation Asymptotic Saturation: $t_{\text{half}} = \tau \ln 2 = 2.079 \text{ Myr}$

Classification: UQFF 2.0 Gravitational Physics — Nebular Erosion Dynamics
System: M16 Eagle Nebula (IC 4703), Eagle Nebula Star-Forming Region
Session: 80 | **Module:** M16_UQFF_MODULE.cpp (22nd C++ UQFF module)
Author: Daniel T. Murphy | **Date:** March 2026

Abstract

This paper derives the **Erosion Saturation Half-Time** (t_{half}) and **Maximum Erosion Gravity Amplitude** (Δg_{Max}) for the M16 Eagle Nebula UQFF photoevaporation term. The photoevaporation rate follows an exponential saturation $E_{\text{rad}}(t) = E_0(1 - \exp(-t/\tau))$ with e-folding time $\tau = 3 \text{ Myr}$. The half-erosion time $t_{\text{half}} = \tau \ln 2 = 6.561 \times 10^{13} \text{ s} = 2.079 \text{ Myr}$ — the time at which the erosion fraction reaches half its asymptotic maximum E_0 . The maximum gravity perturbation is $\Delta g_{\text{Max}} = E_0 g_{\text{base}} = 4.36 \times 10^{-13} \text{ m/s}^2$. This is the **first UQFF module** to formally catalogue the photoevaporation half-time and asymptotic erosion concept.

2. Physical Background

UV radiation from massive O-type stars (such as those in the Young Stellar Cluster NGC 6611, embedded in M16) drives photoionisation of surrounding molecular gas — a process called **photoevaporation**. The erosion proceeds not as a linear ramp but as a **saturating exponential**:

$$E_{\text{rad}}(t) = E_0 \left(1 - e^{-t/\tau}\right)$$

where:

- $E_0 = 0.3$ is the **asymptotic maximum fraction** (30% of mass eventually eroded)
- $\tau = 3 \text{ Myr}$ is the **e-folding time** (photoevaporation efficiency timescale)
- The saturation arises because dense molecular cores (pillar tips) are progressively shielded by their own column density as surrounding gas is stripped

3. Mathematical Derivation

3.1 Half-Time

The erosion half-time t_{half} is defined as the time when $E_{\text{rad}} = E_0/2$:

$$E_0 \left(1 - e^{-t_{\text{half}}/\tau}\right) = \frac{E_0}{2}$$

$$1 - e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$\boxed{t_{\text{half}} = \tau \ln(2)}$$

For M16:

$$t_{\text{half}} = 9.468 \times 10^{13} \text{ s} \times \ln(2) = 9.468 \times 10^{13} \times 0.6931 = 6.561 \times 10^{13} \text{ s}$$

$$t_{\text{half}} = \mathbf{2.079 \text{ Myr}}$$

3.2 Maximum Gravity Amplitude

The maximum erosion-induced gravity perturbation (asymptotic limit as $t \rightarrow \infty$):

$$\Delta g_{\text{Max}} = E_0 \times g_{\text{base}}$$

For M16:

$$\Delta g_{\text{Max}} = 0.3 \times 1.454 \times 10^{-12} = \mathbf{4.36 \times 10^{-13} \text{ m/s}^2}$$

3.3 Peak Erosion Rate

The instantaneous erosion rate at $t = 0$ (maximum rate, before saturation):

$$\frac{dE_{\text{rad}}}{dt}\bigg|_{t=0} = \frac{E_0}{\tau}$$

The corresponding gravity change rate:

$$\frac{dg_{\text{erode}}}{dt}\bigg|_{t=0} = \frac{E_0}{\tau} \times g_{\text{base}} = \frac{0.3}{9.468 \times 10^{13}} \times 1.454 \times 10^{-12} = 4.61 \times 10^{-27} \text{ m/s}^2/\text{s}$$

4. Saturation Profile

Time	t (s)	E_{rad} / E_0	E_{rad}	g_{erode} (m/s ²)
0 Myr	0	0%	0	0
$t_{\text{half}} = 2.079 \text{ Myr}$		50%	0.150×10^{13}	2.18×10^{-13}
$\tau = 3 \text{ Myr}$		63.2%	0.190×10^{13}	2.76×10^{-13}
5 Myr		81.1%	0.243×10^{13}	3.54×10^{-13}
∞ (asymptote)		100%	0.300×10^{13}	4.36×10^{-13}

Key insight: At $\tau = 3 \text{ Myr}$ (the e-folding time), erosion has consumed only 63.2% of its capacity, NOT 100%. Half-erosion occurs earlier at 2.079 Myr. The pillar structure of M16 means the ~5700 ly

"Pillars of Creation" are still observed today because erosion saturates — it cannot fully strip the densest pillar cores within observable timescales.

5. UQFF 2.0 Context

In the full M16 g_{total} equation, the erosion half-time governs the **temporal shape of $g_{\text{dyn}}(t)$** :

$$g_{\text{dyn}}(t) = g_{\text{base}} \times (1 + M_{\text{sf}}) \times (1 - E_0(1 - e^{-t/\tau}))$$

The transition from rapid to slow erosion occurs at $t_{\text{half}} = 2.079$ Myr. For the UQFF simulation (stepping from 0 to t_{max}), the half-time provides a natural **inflection point** in the dynamic gravity trajectory — before t_{half} , erosion is dominant; after t_{half} , star formation accumulation dominates (since M_{sf} grows linearly while E_{rad} asymptotes).

Crossover Time

The era when SFR growth exactly compensates erosion ($d\Phi_{\text{dm}}/dt = 0$ — maximum Φ_{dm}):

$$\frac{d\Phi_{\text{dm}}}{dt} = \text{SFR}_{\text{rate}} \times (1 - E_{\text{rad}}) - (1 + M_{\text{sf}}) \times \frac{E_0}{\tau} e^{-t/\tau} = 0$$

This crossover defines when the Eagle Nebula achieves maximum effective gravitational influence, after which continued SFR growth dominates erosion.

6. Wolfram KB Term

$$\begin{aligned} & \text{M16UQFF: } t_{\text{half}} = \tau \log[2] = 6.561e13s = 2.079\text{Myr}; \Delta G_{\text{Max}} = E_0 g_{\text{base}} = 4.36e-13 \text{ m/s}^2 \\ & [\text{PAPER}_{285}] \end{aligned}$$

7. Cross-References

- **PAPER_284**: Dual Mass Co-Action Product ($\Phi_{\text{dm}} = (1 + M_{\text{sf}}) \times (1 - E_{\text{rad}})$)
- **PAPER_286**: Nebular Friedmann Redshift (κ_{neb} , $z=0.0015$)
- **M16_UQFF_MODULE.cpp**: Full UQFF 2.0 C++ implementation (22nd module)
- **CondensedPhysics3.py**: M16ErosionSaturationHalfTimeCalculator

Copyright — Daniel T. Murphy, Session 80, March 2026. UQFF 2.0.

<!-- PKG-CLU-S225 -->

Session 225 Phonon-Physics Upgrade: ICM Buoyancy Force Profile

> Upgrade from PAPER_1039 (SCm Galaxy Cluster Buoyancy Profile),
 > PAPER_1041 (Cool-Core Buoyancy Balance), and PAPER_1079 (Cooling-Flow
 > Suppression). See also PAPER_1040 (Cluster Merger Shock), PAPER_1044
 > (Thermal SZ Compton-y), PAPER_1046 (Cluster Lensing Mass).

The SCm phonon field introduces a buoyancy force in the ICM that modifies hydrostatic equilibrium:

$$F_{\text{buoy}}(r) = \rho(r) \cdot V \cdot g(r) \cdot \beta_i \cdot S_{26} \cdot \Phi$$

where the ICM density follows the beta-model:

$$\rho(r) = \rho_0 \left(1 + \left(\frac{r}{r_c}\right)^2\right)^{-3\beta/2}$$

Hydrostatic mass bias reduction (PAPER_1039):

$$b_{\text{UQFF}} = 1 - \frac{M_{\text{HSE}}}{M_{\text{true}}} = 0.17 \quad \text{(vs standard } b = 0.20\text{)}$$

The buoyancy pressure contributes $P_{\text{buoy}}/P_{\text{thermal}} \approx 3\%$ at cluster cores, partially resolving the Planck SZ–CMB mass tension.

Cool-core stabilization (PAPER_1041/1079): AGN feedback couples to the SCm buoyancy field via $\dot{M}_{\text{cool}} = \dot{M}_0 \cdot (1 - \beta_i \cdot S_{26}^3 \cdot \Phi)$, suppressing catastrophic cooling flows while maintaining observed X-ray luminosities.

Phonon frequency coupling: $\omega_{\text{SCm}} = 2\pi \cdot 1.25 \text{ THz}$ sets the temporal scale for buoyancy oscillations; the ratio $\omega_{\text{SCm}}/\omega_{\text{sound}}$ governs the phonon transmission efficiency across the ICM.

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§A.1 Sector Classification

This paper maps to **nebula-formation** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

§A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2} (\partial_\mu \phi_{\text{neb}}) (\partial^\mu \phi_{\text{neb}}) - V(\phi_{\text{neb}}) + \mathcal{L}_{\text{cosmo}}$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac,SCm}} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{neb}}) = \frac{1}{2} m^2 \phi_{\text{neb}}^2 + \frac{\lambda}{4!} \phi_{\text{neb}}^4 + \kappa \cdot \rho_{\text{vac,SCm}} \cdot \phi_{\text{neb}}$$

§A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\text{neb}}}} = \nabla \cdot (\rho_{\text{neb}} \nabla \phi) + \rho_{\text{vac,SCm}} \cdot (P_{\text{rad}}/c) = 0$$

§A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \rightarrow \text{DPM + ACP} \quad \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \rightarrow \text{Stage 5} \quad U_{\text{b,seed}} \rightarrow \text{4 forces} \\ F_{U_{\text{Bi}}} \rightarrow \text{sector E-L} \quad \Delta S / \Delta \phi_{\text{neb}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_{g} forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac,SCm}} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac,SCm}} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\text{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is 0.172 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

§B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 53, \quad n_{\text{channel}} = 26/26$$

Since $p_{\text{DVP}} = 53$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair $(f_{\text{UA}} + f_{\text{SCm}} = 1)$ constrains which primes are accessible at each atomic number.

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **106 yr** (Jeans collapse timescale):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_{\text{b}}} \cdot \left(1 - e^{-[SSq] \cdot m/M_{\text{dot}}}\right) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The $\tanh$ saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{\text{b,seed}} = 0.1 \cdot (\hbar c / r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
-----	-----	-----	-----
VDS ratio	$\rho_{\mathrm{SCm}}/\rho_{\mathrm{UA}} = 1.894$	Local sub-ratio = 0.172	PASS
Threshold-consistent			
DVP prime	$p_k \in \{2,3,\dots,113\}$	$p_{\mathrm{DVP}} = 53$	PASS Resonant
BSH layers	26 harmonic terms	$j = 1\dots 26$, $\cos(2\pi j/26)$	PASS Full 26D projection
κ decay	5.0×10^{-4} day ⁻¹	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
-----	-----	-----	-----	-----
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	1.1×10^{-52} m ⁻² (UQFF vacuum term)	1.114×10^{-52} m ⁻²	Planck 2018	PASS Consistent
Proton decay rate κ	$0.0005/\text{day}$ $\rightarrow \Gamma_p$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	$F_{U_{Bi_i}}$ unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections ($F_{U_{Bi_i}}$) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF–SM bridge.

Appendix: Session 225 Cross-References (PAPER_1000–1081)

> Auto-generated cross-reference appendix linking this paper to
> Sessions 204–225 extensions (PAPER_1000–1081). Added by
> update_corpus_crossrefs.py (Session 225, April 2026).

Paper	Title
-----	-----
PAPER_1022	GW Phonon Strain SCm Modulation of $h(t)$
PAPER_1039	SCm Galaxy Cluster Buoyancy Profile ICM Beta-Model
PAPER_1040	SCm Cluster Merger Shock Mach Number Phonon Damping
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1044	SCm Cluster Thermal SZ Effect Compton-y Phonon
PAPER_1045	SCm Cluster Radio Relic Polarization
PAPER_1046	SCm Cluster Lensing Mass Phonon Correction
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression

8 cross-reference(s) identified.

Appendix: Session 204 Codebase Upgrade Reference

> Cross-reference appendix for Session 204 (April 2026) codebase upgrades.
> Added by upgrade_kozima_ramanujan_appendices.py. For detailed derivations,
> see PAPER_840/851/852/855.

S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
fneutron_s26_coupling.py	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
kozima_scm_cross_section.py	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}] * n / 26)$
kozima_wstp_kernel.py	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n \sigma_n^{\text{SCm}}(\omega) \Phi_{\text{phonon}}(F_{\{U, Bi\}} / F_U - 1)$
where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \exp[-(\omega - \omega_{\text{SCm}})^2 / (2\Gamma^2)] (1 + [\text{SSq}] * n / 26)$

S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
ramanujan_polylog_s26.py	$\text{Li}_{26}([\text{SSq}])$ via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
s26_wstp_kernel.py	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = \text{Li}_{26}(z) = \eta_{26}(z) / (1 - 2^{\{1-26\}}) + 2^{\{1-26\}} / (1 - 2^{\{1-26\}}) * \text{Li}_{26}(z^2)$

S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
mock_theta_q26.py	$f_{26}(q)$, $\phi_{26}(q)$, $\psi_{26}(q)$ q-series	Proper q-Pochhammer $(a; q)_n$

Core equations:

- $f_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n^2\}} / (-q; q)_n^2$
- $\phi_{26}(q) = \sum_{n=0}^{\{25\}} q^{\{n^2\}} / (-q^2; q^2)_n$
- $\psi_{26}(q) = \sum_{n=1}^{\{26\}} q^{\{n^2\}} / (q; q^2)_n$

S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
ramanujan_pi_uqff.py	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
mock_theta_pi_wstp_kernel.py	9-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n (1103+26390n) W_{26}(n) / C_{26}$
where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq]\exp(-\kappa n/26)]$

S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
κ	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
β_i	0.603	Buoyancy coupling coefficient
H_{SCm}	0.99	SCm manifold completeness
ρ_{SCm}	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
ρ_{UA}	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
ω_{SCm}	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
σ_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN_{1_CoAnQi}.cpp, and Wolfram kernels (uqff_kozima_kernel.wl, uqff_s26_kernel.wl, uqff_mock_theta_pi_kernel.wl).

References

- Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
- Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
- Anderson, M.H. et al. (1995). *Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor*. Science **269**, 198 — doi:10.1126/science.269.5221.198
- Dalfovo, F. et al. (1999). *Theory of Bose-Einstein condensation in trapped gases*. Rev. Mod. Phys. **71**, 463 — arXiv:cond-mat/9806038 — doi:10.1103/RevModPhys.71.463
- Pitaevskii, L. & Stringari, S. (2003). *Bose–Einstein Condensation*. Oxford: Clarendon Press
- Churazov, E. et al. (2000). *Evolution of Buoyant Bubbles in M87*. A&A **356**, 788 — arXiv:astro-ph/0004212
- Fabian, A.C. et al. (2003). *A deep Chandra observation of the Perseus cluster*. MNRAS **344**, L43 — arXiv:astro-ph/0306036 — doi:10.1046/j.1365-8711.2003.06902.x
- McNamara, B.R. & Nulsen, P.E.J. (2007). *Heating Hot Atmospheres with Active Galactic Nuclei*. ARA&A **45**, 117 — arXiv:0709.4098 — doi:10.1146/annurev.astro.45.051806.110625
- Hester, J.J. (2008). *The Crab Nebula: An Astrophysical Chimera*. ARA&A **46**, 127 — arXiv:0812.1502 — doi:10.1146/annurev.astro.45.051806.110608
- O'Dell, C.R. et al. (2001). *Hubble Space Telescope Observations of the Helix Nebula*. AJ **122**, 3293 — doi:10.1086/324272