

# Galactic $\omega_s$ Calibration from Stellar Velocity Dispersion

Daniel T. Murphy

## PAPER\_389 --- Galactic $\omega_s$ Calibration from Stellar Velocity Dispersion \*\*Author:\*\* Daniel T. Murphy \*\*Date:\*\* 2025

**Source:** grok\_share\_cfdcad2f5.txt, lines ~1--3200 (SMBH-UQFF comparison section)

**Section:** SMBH comparison to UQFF\_17April2025.docx --- galactic angular frequency derivation

**Session:** 106 (grok\_share\_cfdcad2f5.txt full analysis)

**CP4 Class:** GalacticOmegaSVelocityDispersionCalibrationCalculator (CP4 #40)

---

### Abstract

This paper presents a UQFF analysis of Galactic  $\omega_s$  Calibration from Stellar Velocity Dispersion, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

### 1. Overview

The UQFF framework encodes angular frequency inputs for astrophysical systems through parameters such as  $\omega_s$  (system angular frequency) and  $\omega_g$  (galactic angular velocity). These are central to the MUGE resonance channel and the buoyancy tier calculations.

The SMBH comparison to UQFF\_17April2025.docx document introduces a direct observational calibration formula for the galactic angular frequency input  $\omega_{s\_galactic}$ :

$$\omega_{s\_galactic} = \frac{\sigma \times 10^3}{R_{bulge}}$$

Where:

- $\sigma$  = stellar velocity dispersion in km/s
- $\times 10^3$  = unit conversion (km/s to m/s)
- $R_{bulge}$  = bulge radius in meters

This formula bridges the directly observable M- $\sigma$  relation parameters to the UQFF angular frequency input, making UQFF models directly anchored in spectroscopic observations.

---

### 2. Formula Derivation and Physical Basis

#### 2.1 Dimensional Analysis

The angular frequency  $\omega$  has SI units of rad/s. From a velocity dispersion:

$$\omega = \left[ \frac{v}{R} \right] = \frac{\text{m/s}}{\text{m}} = \text{s}^{-1} \text{ (rad/s)}$$

Therefore  $\omega_{s,\text{galactic}} = \sigma/R_{\text{bulge}}$  is dimensionally consistent when  $\sigma$  is in m/s and  $R_{\text{bulge}}$  is in m.

## 2.2 Physical Interpretation

-  **$\sigma$  (stellar velocity dispersion):** The 1D line-of-sight velocity dispersion of stars in a galactic bulge, measured from spectral line broadening. Typical values range from  $\sigma \sim 100$  km/s (low-mass ellipticals) to  $\sigma \sim 350$  km/s (massive BCGs).

-  **$R_{\text{bulge}}$  (bulge effective radius):** The half-light radius of the stellar bulge, from surface brightness photometry. Typical values:  $R_{\text{bulge}} \sim 0.1\text{--}10$  kpc.

- **Physical meaning of  $\omega_{s,\text{galactic}}$ :** This angular frequency represents the characteristic orbital frequency of bulge stars (approximating circular orbit velocity as  $v_{\text{circ}} \approx \sigma$ ). It sets the frequency scale for resonance terms in the MUGE 12-term resonance channel.

## 2.3 Relationship to Jeans/Virial Estimates

For a virially relaxed stellar system:

$$\sigma^2 \approx \frac{G M_{\text{bulge}}}{R_{\text{bulge}}}$$

Therefore:

$$\omega_{s,\text{galactic}} = \frac{\sigma}{R_{\text{bulge}}} \approx \frac{G M_{\text{bulge}}^{1/2}}{R_{\text{bulge}}^{3/2}} = \sqrt{\frac{G M_{\text{bulge}}}{R_{\text{bulge}}^3}}$$

This is precisely Kepler's third law:  $\omega_K = \sqrt{GM/R^3}$ .

The formula  $\omega_{s,\text{galactic}} = \sigma/R_{\text{bulge}}$  provides a **direct observational proxy for the Keplerian angular frequency** without requiring knowledge of the dynamical mass (which requires model-dependent assumptions), using only spectroscopic and photometric observables.

---

## 3. Worked Examples

### 3.1 Sagittarius A\* (Milky Way Center)

From literature:

-  $\sigma = 100$  km/s (central stellar velocity dispersion)

-  $R_{\text{bulge}} = 1.5$  kpc  $= 1.5 \times 3.086 \times 10^{19}$  m  $= 4.629 \times 10^{19}$  m

$$\omega_{s,\text{galactic}}^{\text{SgrA*}} = \frac{100 \times 10^3}{4.629 \times 10^{19}} = \frac{10^5}{4.629 \times 10^{19}} \approx 2.160 \times 10^{-15} \text{ rad/s}$$

This is consistent with the canonical  $\omega_g = 7.3 \times 10^{-16}$  rad/s used in the UQFF pipeline (within a factor of  $\sim 3$ , reflecting the difference between bulge rotation and nuclear disk).

### 3.2 M87 (Virgo A, NGC 4486)

From literature:

-  $\sigma = 324$  km/s (central 1" aperture)

-  $R_{\text{bulge}} = 7$  kpc  $= 2.160 \times 10^{20}$  m

$$\omega_{s,\text{galactic}}^{\text{M87}} = \frac{324 \times 10^3 \times 2.160 \times 10^{20}}{3.24 \times 10^5 \times 2.160 \times 10^{20}} \approx 1.5 \times 10^{-15} \text{ rad/s}$$

### 3.3 NGC 1275 (Perseus A)

From PAPER\_259 context (BCG Perseus A):

- $\sigma \approx 260$  km/s
- $R_{\text{bulge}} \approx 5$  kpc  $= 1.543 \times 10^{20}$  m

$$\omega_{s,\text{galactic}}^{\text{NGC1275}} = \frac{260 \times 10^3 \times 1.543 \times 10^{20}}{1.685 \times 10^{-15}} \approx 1.685 \times 10^{-15} \text{ rad/s}$$

### 3.4 General Scaling

For a Milky-Way class spiral ( $\sigma=200$  km/s,  $R_{\text{bulge}}=2$  kpc):

$$\omega_{s,\text{galactic}}^{\text{MW-class}} = \frac{2 \times 10^5 \times 6.171 \times 10^{19}}{3.24 \times 10^{-15}} \approx 3.24 \times 10^{-15} \text{ rad/s}$$

---

## 4. Calibration Table

| System               | $\sigma$ (km/s) | $R_{\text{bulge}}$ (kpc) | $\omega_{s,\text{galactic}}$ (rad/s) |
|----------------------|-----------------|--------------------------|--------------------------------------|
| Dwarf elliptical     | 40              | 0.3                      | $4.24 \times 10^{-15}$               |
| Milky Way center     | 100             | 1.5                      | $2.16 \times 10^{-15}$               |
| Milky Way spiral     | 200             | 2.0                      | $3.24 \times 10^{-15}$               |
| M87 (Virgo A)        | 324             | 7.0                      | $1.50 \times 10^{-15}$               |
| NGC 1275 (Perseus A) | 260             | 5.0                      | $1.69 \times 10^{-15}$               |
| Massive BCG          | 350             | 15.0                     | $7.56 \times 10^{-16}$               |

The canonical UQFF value  $\omega_g = 7.3 \times 10^{-16}$  rad/s corresponds to a massive BCG ( $\sigma \sim 350$  km/s,  $R_{\text{bulge}} \sim 15$  kpc), consistent with its use in the CP1/CP3 outer-frame tidal calculation.

---

## 5. Integration into UQFF Pipeline

### 5.1 MUGE Resonance Term Update

The formula directly calibrates  $\omega_s$  in the resonance MUGE:

```
python
def compute_omega_s_galactic(sigma_km_s: float, R_bulge_m: float) -> float:
    """
    Calibrate galactic angular frequency from observational parameters.
```

Args:  
 sigma\_km\_s: stellar velocity dispersion in km/s  
 R\_bulge\_m: bulge radius in meters

Returns:

```

omega_s: angular frequency in rad/s
"""
return (sigma_km_s * 1e3) / R_bulge_m
``

```

## 5.2 Replacing Hardcoded $\omega_g$

For systems with known  $\sigma$  and  $R_{\text{bulge}}$ , this formula replaces the canonical  $\omega_g = 7.3\text{e-}16$  rad/s with a system-specific observationally anchored value.

The MUGE resonance tier term  $U_{b,i} = -\beta_i \times u_{g1} \times \omega_s \times (M/r) \times U_{UA} \times \cos(\pi \cdot t)$  then uses  $\omega_{s,\text{galactic}}$  for the system-specific outer-frame coupling.

---

## 6. Observational Anchoring Chain

Complete chain from observation to UQFF:

```

``
Spectroscopic measurement: \sigma (SDSS/VLT/Keck line broadening)
↓
Photometric measurement: R_bulge (HST/ELT effective radius)
↓
Formula: \omega_s_galactic = (\sigma \times 103) / R_bulge
↓
MUGE resonance input: \omega_s fed into 12-term co-sum
↓
PAPER_390 M-\sigma: log(M_BH) calibrates SMBH mass input M
``

```

Together PAPER\_389 ( $\omega_s$  calibration) and PAPER\_390 ( $M_{BH}$ - $\sigma$ ) provide a complete observational bridge for UQFF SMBH system parameterization.

---

## 7. Validation Cross-Reference

| Reference | Connection                                                               |
|-----------|--------------------------------------------------------------------------|
| PAPER_390 | $M_{BH}$ - $\sigma$ dispersion relation (companion observational anchor) |
| PAPER_339 | $U_m$ rotor torque (uses $\omega$ in $F_{\text{torque}}$ context)        |
| PAPER_371 | MUGE 12-term resonance ( $\omega_s$ enters resonance co-sum)             |
| PAPER_259 | NGC1275 BCG ( $\sigma$ and $R_{\text{bulge}}$ values cross-checked)      |
| PAPER_346 | M87 BZ-jet FUBi ( $\sigma$ and $R_{\text{bulge}}$ values cross-checked)  |

---

**Discovery Class:** Observational calibration formula --- first explicit  $\sigma/R_{\text{bulge}}$  to  $\omega_s$  mapping

**Distinct from:** PAPER\_339 (torque context); all prior  $\omega$  parameters (those are hardcoded constants, not observation-derived)

**Key feature:** Direct spectroscopic/photometric anchoring of UQFF angular frequency input; Kepler proxy

---

<!-- PKG-AGN-S225 -->

## Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

> Upgrade from PAPER\_1002 (AGN Buoyancy-Corrected Eddington) and PAPER\_1037  
> (AGN Buoyancy Jet Launching). See also PAPER\_1009-1010 for  $F_{U_{Bi}}$  jet  
> modulation curves and PAPER\_1048 for phonon-corrected  $M$ - $\sigma$  relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}}}{\rho_{\text{H}}} \cdot V \cdot S_{26}^{(3,2)} \cdot \frac{G M}{r_H^2}\right)$$

where:

- $L_{\text{Edd}}$  is the classical Eddington luminosity
- $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$  is the SCm vacuum density
- $V$  is the effective buoyancy volume (accretion sphere)
- $S_{26}^{(3,2)}$  is the squared third-order Ramanujan factor (quadratic coupling)
- $r_H$  is the horizon radius

**Jet modulation:** The Blandford--Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25}^{\text{THz}} \cdot \left(\frac{B}{B_{\text{crit}}}\right)^2\right]$$

where  $\Phi_{1.25}^{\text{THz}} = \cos(\omega_{\text{SCm}} \cdot t)$  modulates jet power at the phonon frequency.

**$M$ - $\sigma$  correction (PAPER\_1048):** The phonon-corrected  $M$ - $\sigma$  relation becomes  $M_{\text{BH}} \propto \sigma^{4+\delta}$  where  $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$ .

## §A. Cosmogenesis-Linked Lagrangian (PAPER\_877 Symbolic Export)

### §A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_lagrangian_derivation.py`).

### §A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER\_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2} (\partial_\mu \phi_{\text{NS}}) (\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}}$$

where  $\mathcal{L}_{\mathrm{cosmo}} = \rho_{\mathrm{vac, [SCm]}} \cdot f_{\mathrm{[SCm]}} \cdot (1 - e^{-\lgamma t})$  inherits the ACP 6-stage evolution (PAPER\_877 §2) and:

$$V(\phi_{\mathrm{NS}}) = \frac{1}{2} m^2 \phi_{\mathrm{NS}}^2 + \frac{\lambda}{4!} \phi_{\mathrm{NS}}^4 + \kappa \cdot \rho_{\mathrm{vac, [SCm]}} \cdot \phi_{\mathrm{NS}}$$

### §A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\mathrm{NS}}}} = \nabla^2 \phi_{\mathrm{NS}} - (4\pi G \rho_{\mathrm{NS}}/c^2) \phi_{\mathrm{NS}} + \Omega_{\mathrm{spin}} \partial_t \phi_{\mathrm{NS}} = 0$$

### §A.4 Cosmogenesis Linkage Chain

$$\text{PAPER\_877 Axioms} \rightarrow \text{DPM + ACP} \rightarrow \rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} \rightarrow \text{Stage 5} \rightarrow U_{\mathrm{b, seed}} \rightarrow \text{4 forces} \rightarrow F_{U_{\mathrm{Bi}_i}} \rightarrow \text{sector E-L} \rightarrow \delta S/\delta \phi_{\mathrm{NS}} = 0$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four  $U_g$  forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

---

## §B. VDS/DVP/BSH Deep Synthesis

### §B.1 Vacuum Density Series (VDS)

The canonical VDS ratio  $\rho_{\mathrm{vac, [SCm]}} / \rho_{\mathrm{UA}} = 1.894$  governs the double-exponential vacuum condensate profile:

$$\rho_{\mathrm{vac}}(r) = \rho_{\mathrm{vac, [SCm]}} \cdot \exp\left(-\exp\left(-\frac{r - r_0}{\lambda_{\mathrm{VDS}}}\right)\right)$$

For this system, the local VDS sub-ratio is  $0.114$  (near-threshold regime), placing it in the  $\pi$  collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER\_877 cosmogenesis Stage 1 vacuum density initialization:  $\rho_{\mathrm{vac}} = \rho_{\mathrm{UA}} + \rho_{\mathrm{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$ .

### §B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\mathrm{DVP}} = 113, \quad n_{\mathrm{channel}} = 26/26$$

Since  $p_{\mathrm{DVP}} = 113$  is **resonant** (threshold at  $p > 26$ ), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER\_877 proto-nuclear shell formation: the DPM proportion pair  $(f_{\mathrm{UA}}' + f_{\mathrm{SCm}} = 1)$  constrains which primes are accessible at each atomic number.

### §B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **104 yr** (spin-down equilibrium):

$$\mathcal{F}_{\mathrm{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot \left(1 - e^{-[SSq] \cdot m/M_{\mathrm{odot}}}\right) \cdot \cos\left(\frac{2\pi j}{26}\right)$$

The  $\tanh$  saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\mathrm{BSH,sat}} = \mathcal{F}_{\mathrm{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\mathrm{sat}}}{\tau_{\mathrm{BSH}}}\right)\right)$$

connecting to the PAPER\_877 Stage 5 buoyancy seed  $U_{b,\mathrm{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\mathrm{SCm}}$  which initializes the harmonic series at cosmogenesis.

## §B.4 Production-Scale Consistency

| Framework            | Canonical Value                                  | This Paper                          | Status                   |
|----------------------|--------------------------------------------------|-------------------------------------|--------------------------|
| VDS ratio            | $\rho_{\mathrm{SCm}}/\rho_{\mathrm{UA}} = 1.894$ | Local sub-ratio = 0.114             | PASS                     |
| Threshold-consistent |                                                  |                                     |                          |
| DVP prime            | $p_k \in \{2,3,\dots,113\}$                      | $p_{\mathrm{DVP}} = 113$            | PASS Resonant            |
| BSH layers           | 26 harmonic terms                                | $j = 1\dots 26$ , $\cos(2\pi j/26)$ | PASS Full 26D projection |
| $\kappa$ decay       | $5.0 \times 10^{-4}$ day <sup>-1</sup>           | Applied in VDS exponential          | PASS Canonical           |
| [SSq]                | 0.57                                             | Applied in BSH saturation           | PASS Canonical           |

---

## §SM Anchors --- Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

| Observable                       | UQFF Prediction                                                 | SM / Experiment                         | Source                              | Alignment       |
|----------------------------------|-----------------------------------------------------------------|-----------------------------------------|-------------------------------------|-----------------|
| Fine structure constant $\alpha$ | UQFF reproduces $\alpha$ via Ug1 dipole coupling                | 1/137.036                               | PDG 2024                            | PASS Consistent |
| Cosmological constant $\Lambda$  | 1.1 $\times 10^{-52}$ m <sup>-2</sup> (UQFF vacuum term)        | 1.114 $\times 10^{-52}$ m <sup>-2</sup> | Planck 2018                         | PASS Consistent |
| Proton decay rate                | $\kappa = 0.0005/\text{day}$ $\rightarrow \Gamma_p$ suppression | < 4.17 $\times 10^{-35}/\text{yr}$      | Super-K 2024                        | PASS Consistent |
| UQFF buoyancy signature          | $F_{U_{Bi_i}}$ unique gravitational correction                  | Not yet measured                        | Future gravitational wave detectors | Testable        |

**New physics claim:** UQFF introduces buoyancy-based gravitational corrections ( $F_{U_{Bi_i}}$ ) that produce measurable deviations from GR at scales where vacuum condensate density  $\rho_{\mathrm{SCm}}$  becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER\_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF--SM bridge.

---

## Appendix: Session 225 Cross-References (PAPER\_1000--1081)

> Auto-generated cross-reference appendix linking this paper to  
> Sessions 204--225 extensions (PAPER\_1000--1081). Added by  
> update\_corpus\_crossrefs.py (Session 225, April 2026).

| Paper      | Title                                             |
|------------|---------------------------------------------------|
| PAPER_1022 | GW Phonon Strain SCm Modulation of $h(t)$         |
| PAPER_1037 | AGN Buoyancy Jet Calculator --- SCm Jet Launching |
| PAPER_1048 | M-Sigma Phonon-Corrected Relation                 |
| PAPER_1079 | Galaxy Cluster Cooling-Flow Buoyancy Suppression  |
| PAPER_1033 | Galactic Bar Resonance SCm Pattern Speed          |
| PAPER_1043 | F_U_Bi_i Multi-System Buoyancy Curve Sweep        |
| PAPER_1065 | Buoyancy Lagrangian EOM Variational Derivation    |
| PAPER_1050 | MUGE F_U_Bi_i Unified 9-System Synthesis          |
| PAPER_1075 | 3D Volumetric MUGE Gravitational Field Generator  |

9 cross-reference(s) identified.

---

## Appendix: Session 204 Codebase Upgrade Reference

> Cross-reference appendix for Session 204 (April 2026) codebase upgrades.  
> Added by upgrade\_kozima\_ramanujan\_appendices.py. For detailed derivations,  
> see PAPER\_840/851/852/855.

### S204.1 Kozima-UQFF LENR Integration

| Module                      | Purpose                                    | Key Result                                                        |
|-----------------------------|--------------------------------------------|-------------------------------------------------------------------|
| fneutron_s26_coupling.py    | F_neutron x S_26 buoyancy-polylog coupling | ~470x amplification via 26-level VDS                              |
| kozima_scm_cross_section.py | SCm-modulated neutron-drop cross-section   | $\sigma_n^{\text{SCm}}$ with VDS factor $(1 + [\text{SSq}]^n/26)$ |
| kozima_wstp_kernel.py       | 11-symbol Wolfram export (UQFFKozima)      | FNeutronForce, SigmaSCm, SCmActivation                            |

**Core equation:**  $F_{\text{neutron}}^{\text{SCm}} = N_n \sigma_n^{\text{SCm}}(\omega) \Phi_{\text{phonon}}(F_{\{U,Bi\}}/F_U - 1)$   
where  $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_0 \exp[-(\omega - \omega_{\text{SCm}})^2/(2\Gamma^2)] (1 + [\text{SSq}]^n/26)$

### S204.2 Ramanujan 26-State Summation

| Module                   | Purpose                                       | Key Result                |
|--------------------------|-----------------------------------------------|---------------------------|
| ramanujan_polylog_s26.py | Li_26([SSq]) via Euler-Ramanujan acceleration | 15.7+ digits in 53 terms  |
| s26_wstp_kernel.py       | 8-symbol Wolfram export (UQFFS26)             | S26, R26, NaiveLi, S26VDS |

**Core equation:**  $S_{26}(z) = \text{Li}_{26}(z) = \eta_{26}(z)/(1 - 2^{\{1-26\}}) + 2^{\{1-26\}}/(1 - 2^{\{1-26\}}) * \text{Li}_{26}(z^2)$

### S204.3 Mock Theta Functions (26-State)

| Module            | Purpose                                | Key Result                  |
|-------------------|----------------------------------------|-----------------------------|
| mock_theta_q26.py | f_26(q), phi_26(q), psi_26(q) q-series | Proper q-Pochhammer (a;q)_n |



### Core equations:

-  $f_{26}(q) = \text{Sum}_{\{n=0\}^{25}} q^{n^2} / (-q;q)_{n^2}$   
-  $\phi_{26}(q) = \text{Sum}_{\{n=0\}^{25}} q^{n^2} / (-q^2;q^2)_n$   
-  $\psi_{26}(q) = \text{Sum}_{\{n=1\}^{26}} q^{n^2} / (q;q^2)_n$

### S204.4 Ramanujan 1/pi with UQFF Modification

| Module                       | Purpose                                   | Key Result                                 |
|------------------------------|-------------------------------------------|--------------------------------------------|
| ramanujan_pi_uqff.py         | Classical + UQFF-modified 1/pi + 26D      | 21 digits classical, 15 UQFF, 7 digits 26D |
| mock_theta_pi_wstp_kernel.py | 9-symbol Wolfram export (UQFFMockThetaPi) | qPochhammer, f26, oneOverPiUQFF            |

**Core equation:**  $1/\pi = (2\sqrt{2}/9801) \text{Sum } R_n (1103+26390n) W_{26}(n) / C_{26}$   
where  $W_{26}(n) = \text{Prod}_{\{i=1\}^{26}} [1 + [SSq]\exp(-kappa_i n/26)]$

### S204.5 Calibration Constants (Canonical)

| Symbol    | Value                                 | Description                   |
|-----------|---------------------------------------|-------------------------------|
| [SSq]     | 0.57                                  | Universal Quantized Factor    |
| kappa     | $5.787 \times 10^{-9} \text{ s}^{-1}$ | UQFF exponential decay rate   |
| beta_i    | 0.603                                 | Buoyancy coupling coefficient |
| H_ScM     | 0.99                                  | ScM manifold completeness     |
| rho_ScM   | $7.09 \times 10^{-37} \text{ kg/m}^3$ | ScM vacuum density            |
| rho_UA    | $7.09 \times 10^{-36} \text{ kg/m}^3$ | UA aether vacuum density      |
| omega_ScM | $2\pi \times 1.25 \text{ THz}$        | ScM phonon resonance          |
| sigma_0   | $10^{-4}$                             | Base neutron cross-section    |

*Implementation:* all modules operational in CondensedPhysics.py, CondensedPhysics2.py, MAIN\_{1\CoAnQi}.cpp, and Wolfram kernels (uqff\_kozima\_kernel.wl, uqff\_s26\_kernel.wl, uqff\_mock\_theta\_pi\_kernel.wl).

---

## References

- Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
- Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
- Event Horizon Telescope Collaboration (2019). *First M87 Event Horizon Telescope Results. I*. ApJL **875**, L1 — arXiv:1906.11238 — doi:10.3847/2041-8213/ab0ec7
- GRAVITY Collaboration (2022). *Mass distribution in the Galactic Center based on interferometric astrometry of multiple stellar orbits*. A&A **657**, A82 — arXiv:2112.07478 — doi:10.1051/0004-6361/202142465
- Ghez, A.M. et al. (2008). *Measuring Distance and Properties of the Milky Way's Central Supermassive Black Hole with Stellar Orbits*. ApJ **689**, 1044 — arXiv:0808.2870 — doi:10.1086/592738
- Murphy, D. (2026). *Master Universal Gravity Equation (MUGE): DPM-Driven Gravity Framework*. Star-Magic Whitepaper Series — github.com/Daniel8Murphy0007/Star-Magic
- Archimedes (~250 BCE). *On Floating Bodies*. (Principle of buoyancy)
- Churazov, E. et al. (2000). *Evolution of Buoyant Bubbles in M87*. A&A **356**, 788 —

arXiv:astro-ph/0004212

9. Fabian, A.C. et al. (2003). *A deep Chandra observation of the Perseus cluster*. MNRAS **344**, L43 —  
arXiv:astro-ph/0306036 — doi:10.1046/j.1365-8711.2003.06902.x