

PAPER_485: UQFF SNR Buoyancy Master Equation — Five Supernova Remnant Systems

Whitepaper | Star-Magic Physics Suite v5.00 Watermark: Copyright — Daniel T. Murphy
| Analyzed: Grok 3 | Date: November 17, 2025

Abstract

This paper presents the Unified Quantum Force Field (UQFF) buoyancy master equation for five supernova remnant (SNR) and high-energy systems: SN 1006 (Type Ia remnant), Eta Carinae (luminous blue variable), Chandra Archive Collection, Galactic Center (near Sgr A*), and Kepler's SNR (SN 1604). The master buoyancy force $F_{U,Bi,i}$ is decomposed into six physically distinct components: LENR (low-energy nuclear reaction), activation energy, dark energy pressure, resonance, neutron production, and relativistic correction. A dual-mode quadratic solver provides SNR expansion mode separation.

1. Master Buoyancy Equation

$$F_{U,Bi,i} = F_{LENR} + F_{act} + F_{DE} + F_{res} + F_{neutron} + F_{rel}$$

1.1 Component Equations

F_{LENR} (Low-Energy Nuclear Reaction Force):

$$F_{LENR} = k_{LENR} \cdot \rho_{vac} \cdot \sin(\omega_{LENR}t) \cdot e^{-r/r_0}$$

where $\omega_{LENR} = 2\pi \times 1.25 \times 10^{12}$ rad/s, $r_0 = 1$ kpc, $F_0 = 1.83 \times 10^{71}$ N.

F_{act} (Activation Energy Force):

$$F_{act} = k_{act} \cdot F_0 \cdot \sin(\omega_{ACT}t) \cdot B \cdot Q_{wave}$$

where $\omega_{ACT} = 2\pi \times 300$ rad/s.

F_{DE} (Dark Energy Pressure):

$$F_{DE} = k_{DE} \cdot \rho_{vac} \cdot r^2 \cdot (1 + z)$$

F_{res} (Resonance Force):

$$F_{res} = k_{res} \cdot F_0 \cdot \cos(\omega_{ACT}t)/r^2$$

$F_{neutron}$ (Neutron Production Force):

$$F_{neutron} = k_n \cdot \eta \cdot m_n c^2 / r^2, \quad \eta = \eta_0 \cdot B / \sqrt{\rho_{vac}}$$

F_{rel} (Relativistic Correction):

$$F_{rel} = k_{rel} \cdot F_{rel,0} \cdot e^{-v_{exp}/c} \cdot (1 + z)^2$$

where $F_{rel,0} = 4.30 \times 10^{33}$ N.

2. Quadratic Buoyancy Mode Solver

For SNR expansion mode separation, the master equation is reformulated as:

$$a \cdot x^2 + b \cdot x + c = 0$$

where x represents the expansion mode amplitude. Both roots are solved:

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For systems with $\Delta = b^2 - 4ac < 0$ (complex modes), the real part $x = -b/(2a)$ is returned, representing the oscillating equilibrium expansion mode.

3. System Parameters

System	Mass (kg)	r (m)	v_exp (m/s)	B (T)	z	t_age (s)
SN 1006	1.989e31	6.17e16	1e6	1e-5	1.0	3.213e10
Eta Carinae	3.978e32	3.09e16	5e5	1e-4	0.0	5.05e14
Chandra Archive	5.967e31	3.09e19	2e5	1e-5	0.1	1.892e16
Galactic Center	1.989e36	2.461e20	1e5	1e-4	0.0	1.577e17
Kepler's SNR	1.989e31	1.852e17	7e3	1e-5	0.0	1.293e10

4. Key Constants

Constant	Symbol	Value
Normalization force	F_0	$1.83 \times 10^{-1} \text{ N}$
Vacuum energy density	ρ_{vac}	$7.09 \times 10^{-3} \text{ J/m}^3$
LENR angular frequency	ω_{LENR}	$2 \times 1.25 \times 10^{12} \text{ rad/s}$
Activation frequency	ω_{ACT}	$2 \times 300 \text{ rad/s}$
Relativistic base	$F_{rel,0}$	$4.30 \times 10^{33} \text{ N}$
g_rt placeholder	—	$F_{total}/(M_{\odot} \cdot Q_{wave})$

5. Physical Motivation

SNR expansion: Supernova remnants expand against interstellar medium. The UQFF buoyancy force models this expansion as a quantum vacuum pressure effect, analogous to physical buoyancy in a fluid but operating via vacuum energy displacement.

LENR component: Models low-energy nuclear reactions triggered by the extreme thermodynamic conditions in SNR shock fronts, contributing to neutron flux.

F_DE (Dark Energy): The $\rho_{\text{vac}} \times r^2$ term captures the cosmological constant's contribution to remnant expansion at large scales (Galactic Center, Chandra Archive).

F_rel: Relativistic ejecta ($v_{\text{exp}} \sim 0.002c$ for SN1006) require the Lorentz-suppression factor $e^{-(v/c)}$, preventing unphysical super-luminal contributions.

6. System Results Preview

System	F_U_Bi_i (N)	Dominant Component
SN 1006	~F_DE-dominated	Dark energy pressure
Eta Carinae	~F_LENRR-dominated	Nuclear reactions
Chandra Archive	~F_DE-dominated	Large-scale dark energy
Galactic Center	~F_LENRR+F_act	Near-nucleus activation
Kepler's SNR	~F_rel-dominated	Relativistic ejecta

7. Integration Reference

- **C++ Implementation:** Core/Modules/UQFFBuoyancySNRModule.cpp
- **Header:** UQFFBuoyancySNRModule.h
- **Related Papers:** PAPER_486 (Cassini buoyancy), PAPER_484 (U_g components)
- **CondensedPhysics2.py class:** UQFFBuoyancySNRCalculator (v4.3.9)