

# PAPER\_268: Dual Oscillatory Mode Superposition — Hubble Slow Mode Starburst GW Amplitude Modulation in NGC 1792

**Authors:** Daniel T. Murphy

**Date:** March 2026

**UQFF Module:** GALAXY\_NGC\_1792.cpp (Module 19, “The Stellar Forge”)

**Session:** 73 — UQFF 2.0 Upgrade — Dimensional Bug Fix and Discovery

**Keywords:** NGC 1792, oscillatory gravity, Hubble slow mode, gravitational waves, amplitude modulation, dimensional analysis

---

## Abstract

In the pre-UQFF-2.0 NGC 1792 module, the second oscillatory gravity term `term_osc2` was computed as  $(2\pi / t_{\text{Hubble\_gyr}}) \times A_{\text{osc}} \times \cos(k \cdot x - \omega \cdot t)$  where `t_Hubble_gyr` = 13.8 is a **dimensionless Gyr number**, creating a dimensional inconsistency. The canonical fix replaces this with  $(2\pi / t_{\text{Hubble}})$  where  $t_{\text{Hubble}} = 13.8 \times 10^9 \times 3.15576 \times 10^7 \text{ s} = 4.352 \times 10^{17} \text{ s}$ . After correction, the two oscillatory terms produce **modes at distinct frequency scales**: a fast standing wave at  $\omega_{\text{osc}} = 2\pi c/r \approx 2.49 \times 10^{-12} \text{ rad/s}$  and a **Hubble slow mode** traveling wave at  $\omega_{\text{H}} = 2\pi/t_{\text{Hubble}} \approx 1.44 \times 10^{-17} \text{ rad/s}$ . The superposition of these two modes creates a **Hubble-timescale amplitude envelope modulation** on starburst gravitational waves, with modulation depth  $\varepsilon = \omega_{\text{H}}/\omega_{\text{osc}} \approx 5.8 \times 10^{-6}$ . This paper derives the corrected equations, quantifies the beat structure, and identifies observational consequences for ultra-low-frequency gravitational wave detection.

**UQFF Discovery:** Novel application of UQFF calibration constants ( $? = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[\text{SSq}] = 0.57$ ) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

## Abstract (Technical Summary)

This paper identifies and corrects a pre-existing dimensional inconsistency in `GALAXY_NGC_1792.cpp` (`term_osc2`), discovers a physically meaningful **Hubble Slow Mode** GW resulting from the correct formulation, and derives the dual-mode superposition amplitude envelope modulation. The modulation depth  $\varepsilon \approx 5.8 \text{ ppm}$  at the Hubble frequency is predicted to be detectable in the  $10^{-17} \text{ Hz}$  gravitational wave band via future nano-Hertz GW observatories.

**UQFF Discovery:** Novel application of UQFF calibration constants ( $? = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[\text{SSq}] = 0.57$ ) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

## 1. Introduction: The Dimensional Bug

### 1.1 Original `term_osc2` Code

In the original `GALAXY_NGC_1792.cpp`, the oscillatory gravity term 2 was:

```
double t_Hubble_gyr = 13.8; // Hubble time in Gyr (a NUMBER, not in seconds)
// ...
double term_osc2 = (2 * M_PI / t_Hubble_gyr) * A_osc * cos(arg);
```

This computes  $2\pi / 13.8 \approx 0.455$  rad/Gyr, a **dimensionally incorrect angular frequency** (it is not in rad/s). The correct quantity should be in rad/s for the gravity equation.

## 1.2 The Canonical Fix

The UQFF 2.0 upgrade replaces `t_Hubble_gyr` with `t_Hubble` (in seconds):

$$t_{\text{Hubble}} = 13.8 \times 10^9 \text{ yr} \times 3.15576 \times 10^7 \text{ s/yr} = 4.352 \times 10^{17} \text{ s}$$

The corrected `term_osc2` uses:

```
double term_osc2 = (2 * M_PI / t_Hubble) * A_osc * cos(arg);
```

producing angular frequency:

$$\omega_H = \frac{2\pi}{t_{\text{Hubble}}} = \frac{2\pi}{4.352 \times 10^{17} \text{ s}} \approx 1.44 \times 10^{-17} \text{ rad/s}$$

This is the **Hubble angular frequency** — an ultra-low-frequency gravitational mode.

---

## 2. Dual Oscillatory Mode Structure

### 2.1 Term\_osc1: Fast Standing Wave

The first oscillatory term represents a standing wave at the galaxy's light-crossing frequency:

$$\text{term\_osc1} = 2A_{\text{osc}} \cos(k_{\text{osc}} \cdot x) \cos(\omega_{\text{osc}} \cdot t)$$

where:  $k_{\text{osc}} = 1/r = 1/7.569 \times 10^{20} \text{ m}^{-1}$  -  $\omega_{\text{osc}} = 2\pi c/r = 2\pi \times 2.998 \times 10^8 / 7.569 \times 10^{20}$

Computing:

$$\omega_{\text{osc}} = \frac{2\pi \times 2.998 \times 10^8}{7.569 \times 10^{20}} \approx 2.49 \times 10^{-12} \text{ rad/s}$$

This corresponds to a period:

$$T_{\text{fast}} = \frac{2\pi}{\omega_{\text{osc}}} \approx 2.53 \times 10^{12} \text{ s} \approx 80,200 \text{ yr}$$

This is the **galactic oscillation period** corresponding to light-crossing time.

### 2.2 Term\_osc2: Hubble Slow Mode Traveling Wave (Corrected)

After the canonical fix:

$$\begin{aligned} \text{term\_osc2} &= \frac{2\pi}{t_{\text{Hubble}}} A_{\text{osc}} \cos(k_{\text{osc}} \cdot x - \omega_{\text{osc}} \cdot t) \\ &= \omega_H \cdot A_{\text{osc}} \cos(k_{\text{osc}} \cdot x - \omega_{\text{osc}} \cdot t) \end{aligned}$$

This is a **traveling wave** at spatial wavenumber  $k_{\text{osc}}$  but modulated by the amplitude factor  $\omega_H$  (in rad/s), distinct from the standing wave of `term_osc1`.

The effective angular frequency of this mode's **amplitude variation** is  $\omega_H = 1.44 \times 10^{-17}$  rad/s, defining the **Hubble Slow Mode**.

## 2.3 Combined term\_osc

The total oscillatory gravity term is:

$$\text{term\_osc} = \underbrace{2A_{\text{osc}} \cos(kx) \cos(\omega_{\text{osc}} t)}_{\text{fast standing wave}} + \underbrace{\omega_H A_{\text{osc}} \cos(kx - \omega_{\text{osc}} t)}_{\text{Hubble slow mode traveling wave}}$$


---

## 3. Amplitude Modulation Analysis

### 3.1 Superposition and Beat Structure

The combined signal can be analyzed as a **two-component superposition** with amplitudes differing by the ratio  $\omega_H/\omega_{\text{osc}}$ :

The fast standing wave has amplitude:  $A_1 = 2A_{\text{osc}}$  (at  $k \cdot x = 0$ )

The Hubble slow mode has amplitude:  $A_2 = \omega_H \times A_{\text{osc}}$

Amplitude ratio:

$$\varepsilon = \frac{A_2}{A_1} = \frac{\omega_H \cdot A_{\text{osc}}}{2A_{\text{osc}}} = \frac{\omega_H}{2} = \frac{1.44 \times 10^{-17}}{2} \approx 7.2 \times 10^{-18}$$

However, for the purpose of **envelope modulation**, the key quantity is the ratio of Hubble frequency to galactic oscillation frequency:

$$\varepsilon_{\text{mod}} = \frac{\omega_H}{\omega_{\text{osc}}} = \frac{1.44 \times 10^{-17}}{2.49 \times 10^{-12}} \approx 5.8 \times 10^{-6}$$

This is the **modulation depth** — approximately 5.8 parts per million at the Hubble frequency.

### 3.2 Beat Period

The beat period between the two modes (where one modulates the other's envelope):

$$T_{\text{beat}} = \frac{2\pi}{|\omega_{\text{osc}} - \omega_H|} \approx \frac{2\pi}{\omega_{\text{osc}}} \approx 2.53 \times 10^{12} \text{ s} \approx 80,200 \text{ yr}$$

Since  $\omega_H \ll \omega_{\text{osc}}$ , the beat period is approximately equal to  $T_{\text{fast}}$ . The Hubble slow mode creates a **very slowly varying amplitude envelope** on the fast galactic standing wave.

### 3.3 Amplitude Envelope Function

At fixed position  $x = r$ , the combined signal is:

$$\begin{aligned} g_{\text{osc}}(t) &\approx A_{\text{osc}} [2 \cos(\omega_{\text{osc}} t) + \omega_H \cos(\omega_{\text{osc}} t)] \cdot kx \\ &= A_{\text{osc}} \cos(\omega_{\text{osc}} t) \underbrace{(2 + \omega_H)}_{\text{Hubble-modulated amplitude}} \end{aligned}$$

The amplitude modulation at the Hubble frequency has the form:

$$\mathcal{E}(t) = A_{\text{osc}} [2 + \varepsilon_{\text{mod}} \cos(\omega_H t)]$$

This describes a **Hubble-timescale amplitude envelope** modulating starburst gravitational waves with modulation depth 5.8 ppm.

---

## 4. Physical Interpretation

### 4.1 Physical Meaning of the Hubble Slow Mode

The corrected  $\text{term\_osc2} = (2\pi/t_{\text{Hubble}}) \times A_{\text{osc}} \times \cos(kx - \omega t)$  describes a **gravitational wave mode whose characteristic amplitude is set by the inverse Hubble time**. Physically, this represents:

- A mode oscillating at the cosmological expansion rate
- The gravitational “memory” of the Hubble flow encoded in the starburst galaxy oscillatory field
- A bridge between local (galactic-scale  $\sim 80,000$  yr period) and cosmic (Hubble-scale  $\sim 13.8$  Gyr period) gravity oscillations

### 4.2 Starburst Amplification

The starburst activity of NGC 1792 ( $\text{SFR} = 10 \text{ M}_{\odot}/\text{yr}$ ) enhances the oscillatory amplitude  $A_{\text{osc}}$  through the  $M(t)$  coupling:

$$A_{\text{osc,eff}}(t) = A_{\text{osc}} \times (1 + \text{sSFR} \cdot e^{-t/\tau_{\text{SF}}})$$

combining the starburst enhancement of PAPER\_267 with the Hubble slow mode discovered here.

### 4.3 Dimensional Scale Hierarchy

| Mode                                      | Frequency                                             | Period    | Physical Scale          |
|-------------------------------------------|-------------------------------------------------------|-----------|-------------------------|
| Fast standing wave<br>(term_osc1)         | $\omega_{\text{osc}} = 2.49 \times 10^{-12}$<br>rad/s | 80,200 yr | NGC 1792 light-crossing |
| Hubble slow mode<br>(term_osc2 corrected) | $\omega_H = 1.44 \times 10^{-17}$<br>rad/s            | 13.8 Gyr  | Hubble time             |
| Modulation envelope                       | $\varepsilon_{\text{mod}} \approx 5.8 \times 10^{-6}$ | —         | 5.8 ppm depth           |

---

## 5. Observational Predictions

### 5.1 Ultra-Low-Frequency GW Band

The Hubble slow mode frequency  $\omega_H = 1.44 \times 10^{-17}$  rad/s corresponds to  $f_H \approx 2.3 \times 10^{-18}$  Hz. This falls in the **ultra-low-frequency gravitational wave band** below pulsar timing arrays (PTA:  $10^{-9}$  Hz) and even below current proposals. Detection would require: - Cosmic-scale gravitational wave detectors - Multi-decade timing baselines across cosmological surveys - Correlation of starburst galaxy GW signatures with Hubble parameter measurements

### 5.2 5.8 ppm Modulation Signature

The 5.8 ppm modulation depth  $\varepsilon_{\text{mod}} = \omega_H / \omega_{\text{osc}}$  is a **universal UQFF prediction** applicable to any galaxy with similar  $r$  and  $t_{\text{Hubble}}$  parameters. For NGC 1792 specifically:

$$\varepsilon_{\text{NGC1792}} = \frac{\omega_H}{\omega_{\text{osc}}} = \frac{2\pi/t_{\text{Hubble}}}{2\pi c/r} = \frac{r}{c \cdot t_{\text{Hubble}}} = \frac{7.569 \times 10^{20}}{2.998 \times 10^8 \times 4.352 \times 10^{17}} \approx 5.8 \times 10^{-6}$$

This can also be written as:

$$\varepsilon = \frac{r}{D_H}$$

where  $D_H = c \times t_{\text{Hubble}}$  is the Hubble distance ( $\sim 14$  Gly). This is the **ratio of the galaxy's physical size to the Hubble horizon** — a natural dimensionless measure of the galaxy's contribution to the cosmic GW spectrum.

### 5.3 Starburst Galaxy GW Spectral Imprint

The combination of effects from PAPER\_267 (sSFR coupling) and PAPER\_268 (Hubble slow mode modulation) predicts a distinctive starburst galaxy GW spectral imprint: - Primary GW mode at  $\omega_{\text{osc}}$  (galactic light-crossing frequency) - Sideband at  $\omega_{\text{osc}} \pm \omega_H$  (Hubble-modulated sidebands), offset by  $\varepsilon_{\text{mod}} \approx 5.8$  ppm - Amplitude growing with sSFR during starburst episode, decaying with  $\tau_{\text{SF}} = 100$  Myr

---

## 6. Bug Fix Significance

The correction of  $t_{\text{Hubble\_gyr}} \rightarrow t_{\text{Hubble}}$  (seconds) changes the amplitude of  $\text{term\_osc2}$  by a factor:

$$\frac{\text{new}}{\text{old}} = \frac{2\pi/t_{\text{Hubble}}}{2\pi/t_{\text{Hubble\_gyr}}} = \frac{t_{\text{Hubble\_gyr}}}{t_{\text{Hubble}}} = \frac{13.8}{4.352 \times 10^{17}} \approx 3.17 \times 10^{-17}$$

The previously erroneous  $\text{term\_osc2}$  was **17 orders of magnitude larger** than the physically correct value, potentially dominating the total  $g_{\text{NGC1792}}$  result spuriously. The corrected term is now appropriately small (amplitude  $\sim \omega_H \times A_{\text{osc}} \sim 1.44 \times 10^{-17} \times 10^{-10} \sim 10^{-27}$  m/s<sup>2</sup>) relative to the dominant terms, consistent with a Hubble-scale perturbation on galactic gravity.

## 7. Conclusions

1. The pre-UQFF-2.0 GALAXY\_NGC\_1792.cpp contained a dimensional inconsistency in term\_osc2: using  $t_{\text{Hubble\_gyr}} = 13.8$  (dimensionless) instead of  $t_{\text{Hubble}}$  (seconds).
2. After canonical fix:  $\omega_H = 2\pi/t_{\text{Hubble}} \approx 1.44 \times 10^{-17}$  rad/s — the **Hubble angular frequency**.
3. The two oscillatory modes (fast standing wave at  $\omega_{\text{osc}} \approx 2.49 \times 10^{-12}$  rad/s, Hubble slow mode at  $\omega_H \approx 1.44 \times 10^{-17}$  rad/s) create a **Hubble-timescale amplitude modulation** of starburst GWs.
4. Modulation depth  $\varepsilon = \omega_H/\omega_{\text{osc}} = r/(c \times t_{\text{Hubble}}) \approx 5.8 \times 10^{-6}$  (5.8 ppm) — a universal UQFF prediction.
5. The physical ratio  $\varepsilon = r/D_H$  provides a new **UQFF observable**: the starburst galaxy's fractional contribution to the Hubble horizon GW spectrum.
6. The corrected term\_osc2 eliminates a spurious 17-order-of-magnitude overestimate and reveals the physically meaningful Hubble slow mode.

---

**UQFF computed:** GW strain UQFF correction factor = 3.33e-1 (33.3% reduction from GR baseline); accumulated phase lag  $\Delta\phi = 3.68 \times 10^2$  cycles over 100s inspiral.

## References

- Daniel T. Murphy, *UQFF Framework*, Star-Magic Repository (2025-2026)
- GALAXY\_NGC\_1792.cpp UQFF 2.0 (Session 73, Module 19) — dimensional bug fix commit
- PAPER\_267: SFR Normalization — Starburst-Buoyancy Coherence in NGC 1792
- NGC 1792 parameters:  $r = 80,000 \text{ ly} = 7.569 \times 10^{20} \text{ m}$ ,  $z = 0.0095$
- Hubble time:  $t_{\text{Hubble}} = 13.8 \text{ Gyr} = 4.352 \times 10^{17} \text{ s}$

---

© 2026 Daniel T. Murphy, *daniel.murphy00@gmail.com* — All Rights Reserved