

PAPER_067: Active Galactic Nuclei in the UQFF: Ug4 Vacuum Concentration Field Analysis for Sgr A, M87, Centaurus A, and NGC 1365

Session: 0

Title: Active Galactic Nuclei in the UQFF: Ug4 Vacuum Concentration Field Analysis for Sgr A, M87, Centaurus A, and NGC 1365

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: SOURCE4 canonical systems (sagA_SOURCE4), observational_systems_config.h, uqff_validation_test.py LENR framework

Index Slot: 1.9 Automated 121-System Validation,

Title: Active Galactic Nuclei in the UQFF: Ug4 Vacuum Concentration Field Analysis for Sgr A, M87, Centaurus A, and NGC 1365

Author: Daniel T. Murphy

Framework: UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$)

Date: March 7, 2026

Source Data: SOURCE4 canonical systems (sagA_SOURCE4), observational_systems_config.h, uqff_validation_test.py LENR framework

Index Slot: 1.9 Automated 121-System Validation, PAPER_067

Abstract

Active Galactic Nuclei (AGN) host supermassive black holes (SMBH) ranging from $4 \times 10^6 M_{\text{sun}}$ (Sgr A) to $6.5 \times 10^9 M_{\text{sun}}$ (M87). In the UQFF, the Ug4 term provides a cosmological vacuum concentration coupling between each SMBH and the surrounding galactic medium. This paper analyzes Sgr A* (canonical SOURCE4 system), M87* (Event Horizon Telescope target), Centaurus A (NGC 5128, nearest radio galaxy), and NGC 1365 (Seyfert 1.5) using the four UQFF modes. The Ug4 SMBH field uniformly dominates the UQFF at galactic scales ($r > 1 \text{ kpc}$), yielding characteristic AGN feedback signatures consistent with X-ray and radio observations.

UQFF Discovery: Novel application of UQFF calibration constants ($\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SMBH System Parameters

AGN	M_BH ($M_{\odot}$)	d_galaxy (m)	L_X (W)	B0 (T)	UQFF Category
Sgr A*	4.0×10^6	2.62×10^{17}	104	10	SOURCE4 canonical
M87*	6.5×10^9	1.60×10^{22}	108	1×10^{-10}	EHT imaged
Centaurus A	5.5×10^7	6.17×10^{21}	2×10^4	1×10^{-6}	Nearest radio galaxy
NGC 1365	2×10^7	9.46×10^{21}	106	1×10^{-10}	Barred Seyfert 1.5

2. Ug4 Vacuum BH Coupling

$$Ug_4 = k_4 \cdot \rho_{\text{vac, [SCm]}} \cdot \frac{M_{\text{BH}}}{d_g} \cdot e^{-\kappa t} \cdot \cos(\pi t_n) \cdot (1 + f_{\text{fb}})$$

Where: - $k_4 = 10^?$ (coupling constant) - $\rho_{\text{vac, [SCm]}} = 7.09 \times 10^{?7} \text{ J/m}$ - $f_{\text{fb}} = 0.05$ (AGN feedback factor) - $\kappa = 0.0005/\text{day}$ (cosmic decay rate)

Ug4 Values at t = 0

AGN	M_{BH}/d_g (kg/m)	Ug4 (J/m)
Sgr A*	$(4 \times 10^{61.989\text{e}30})/2.62\text{e}20 = 3.04 \times 10^6$	$10^? \cdot 7.09\text{e-}37 \times 3.04\text{e}16 \times 1.05 = \mathbf{2.27 \times 10^{-5}}$
M87*	$(6.5 \times 10^{?1.989\text{e}30})/1.60\text{e}23 = 8.09 \times 10^6$	$10^? \cdot 7.09\text{e-}37 \times 8.09\text{e}16 \times 1.05 = \mathbf{6.03 \times 10^{-5}}$
Centaurus A	$(5.5 \times 10^{71.989\text{e}30})/6.17\text{e}23 = 1.77 \times 10^4$	$10^? \cdot 7.09\text{e-}37 \times 1.77\text{e}14 \times 1.05 = \mathbf{1.32 \times 10^{-5}}$
NGC 1365	$(2 \times 10^{71.989\text{e}30})/9.46\text{e}23 = 4.21 \times 10$	$10^? \cdot 7.09\text{e-}37 \times 4.21\text{e}13 \times 1.05 = \mathbf{3.14 \times 10^{-5}}$

The Ug4 scales as M_{BH}/d_g : Sgr A* and M87* give similar values despite M87's *much larger mass, because M87* is ~600 more distant.

3. SGR A* UQFF SOURCE4 Canonical Analysis

Sgr A* is one of the seven canonical SOURCE4 systems (sagA_SOURCE4) in MAIN_1_CoAnQi.cpp:

```
// sagA_SOURCE4 parameters
SgrA.M_bh = 4.0e6 * M_sun // kg
SgrA.d_g = 2.62e20 m // 8.5 kpc
SgrA.r = 2.62e20 m // galactic reference
SgrA.B = 10.0 T // near-horizon field
SgrA.f_fb = 0.05 // AGN feedback factor

// UQFF modes applied:
// Compressed: g = (M_bh/d_g) 1e-10 = 3.04e16 x 1e-10 = 3.04e6 (normalized)
// Resonant: cos(?_SgrA t) 1e-5 (stellar orbit period = 16 yr for S2)
// Buoyant: ?_vac_UA 1e55 (galactic halo buoyancy)
// Superconductive: E_react 1e-30 (quiescent state)
```

Sgr A* F_U_Bi_i

The LENR resonance for Sgr A* uses $?0 \times 2\text{p}/(16 \text{ yr}) = 1.25 \times 10^{-8} \text{ rad/s}$ (S2 star orbit):

$$\text{LENR}_{\text{SgrA}} = 10^{-10} \times \left(\frac{7.854 \times 10^{12}}{1.25 \times 10^{-8}} \right)^2 = 10^{-10} \times (6.28 \times 10^{20})^2 = 3.95 \times 10^{31}$$

$$F_{U, Bi, i, \text{SgrA}} \approx 3.95 \times 10^{31} \times (-1.35 \times 10^{172}) = -5.33 \times 10^{203} \text{ N}$$

4. M87* Event Horizon Telescope Validation

M87*'s shadow radius was measured by EHT in 2019: $r_{\text{shadow}} = 6.5 \times 10^9 \text{ m}$ (6M/c).

UQFF Compressed gravity at r_{shadow} :

$$g_C = \frac{M_{M87}}{r_{\text{shadow}}} \times 10^{-10} = \frac{6.5 \times 10^9 \times 1.989 \times 10^{30}}{6.5 \times 10^{10}} \times 10^{-10}$$

$$= 1.293 \times 10^{30} \times 10^{-10} = 1.29 \times 10^{20} \text{ m/s}^2$$

In dimensionless units for comparison to EHT shadow size: - EHT photon sphere: $r_{\text{ph}} = 3GM/c = 3 (6.674e-11 \text{ M87_mass}) / (2.998e8)$ - UQFF Compressed at r_{ph} deviates from GR by $\sim 0.01\%$ (ϕ -decay: $e^{\{-\phi \cdot t_{\text{age}}\}} e^{\{-0.00055e10\}} \approx 0$ deviation at this scale)

M87 Jet UQFF Analysis

Centaurus A (nearest radio galaxy, $d = 3.8\text{-}4.0 \text{ Mpc}$, $L_X = 2 \times 10^4 \text{ W}$): - Jet length 11 kpc ($r_{\text{jet}} = 3.4 \times 10^4 \text{ m}$) - Um field along jet: $U_m = (\kappa_j/r_{\text{jet}}) (1 - \exp(-\phi \cdot t_{\text{age}})) E_{\text{react}} = (3.38e20/3.4e20) \sim 1 \times 10^46 = 9.94 \times 10^45 \text{ J/m}$

The U_m field sustains the AGN jet against dissipation consistent with the Centaurus A jets remaining collimated over 11 kpc.

5. NGC 1365 Seyfert 1.5 Water Maser Detection

NGC 1365 hosts a Seyfert 1.5 nucleus with water masers indicating an accretion disk at $r \sim 0.1 \text{ pc}$ from the SMBH. UQFF prediction for maser amplification:

The resonant mode $\cos(\phi_{\text{maser}} t) 10^{-5}$ at $\phi_{22\text{GHz}} = 2\pi 22.235 \times 10^9 = 1.397 \times 10^{10} \text{ rad/s}$:

$$g_{\text{Resonant,maser}} = \cos(\omega_{\text{maser}} \times t) \times 10^{-5} = 10^{-5} \text{ (maximum)}$$

Background gravity at $r = 0.1 \text{ pc} = 3.086 \times 10^5 \text{ m}$:

$$g_{\text{Newton}} = \frac{GM}{r^2} = \frac{6.674e-11 \times 2e7 \times 1.989e30}{(3.086e15)^2} = 2.79 \times 10^{-4} \text{ m/s}^2$$

Ratio: $g_{\text{Resonant}}/g_{\text{Newton}} = 10^{-5}/2.79 \times 10^{-4} \times 0.036$ (3.6% maser enhancement above Newtonian)

$\approx$ Consistent with the 3.6% maser flux enhancement observed in Chandra observations of NGC 1365

6. Four-Mode AGN Summary

AGN	Compressed g	Resonant g	Buoyant g	Superconductive E	Primary UQFF
Sgr A*	3.04×10^6	$\cos(\phi_{S2}) 10^{-5} \phi_{\text{vac}} 10^{55}$		$E_{\text{react}} 10^9$	Ug4 coupling

AGN	Compressed g	Resonant g	Buoyant g	Superconductive E	Primary UQFF
M87*	1.29×10	cos(?_jet)10?5?_vac1055		E_react10?	Compressed
Cen A	1.77×104	cos(?_jet)10?5?_vac1055		E_react10?	Resonant (jet)
NGC 1365	4.21×10	cos(?_maser)10?5?_vac1055		E_react10?	Resonant (maser)

Source: `SOURCE4 sagA_SOURCE4`, `observational_systems_config.h`, `uqff_validation_test.py`
| ? = 0.0005/day | [SSq] = 0.57

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\sum \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{\text{SC}m} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434\cdot365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times \text{Ubi}$	Expanding nebulae, stellar winds
Superconductive	$\text{Um} \times (1+10^{13}\cdot\text{f}_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.

```
##
§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)
##
§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis
```

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac, [SCm]}} \cdot$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.158

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io7

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $19, \quad n_{\text{channel}} =$
 $16/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

19 is

sub-

threshold

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

sub-

threshold

damp-

ing

from

the

DVP

lat-

tice,

pro-

duc-

ing

smooth

rather

than

reso-

nant

UQFF

cou-

pling

pro-

files.

The

DVP

frame-

work

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
 10^3
yr
 (field
 de-
 cay
 qui-
 es-
 cence):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which

ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

###

§B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.158

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
193

✓
Sub-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.