

PAPER_204: UQFF Dark Matter — NFW, SIDM, Rotation Curves, and Virial Theorem

Version: 1.0

Date: March 13, 2026

Session: 50 — grok_share_7514fe.txt Full Audit

Author: Star-Magic UQFF Research Framework

Source: grok_share_7514fe.txt lines 6096-6110 (BB_C_Equations items 1326-1340)

$$F_U(r, t) = \sum_{i=1}^4 U_{gi} + U_m + U_A - U_{bi}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, \quad [SSq] = 0.57$$

$$U_{bi}(r) = \kappa \cdot [SSq] \cdot \frac{GM}{r^2}, \quad \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, \quad [SSq] = 0.57, \quad \beta_i = 0.61$$

Abstract

This paper applies the UQFF buoyancy framework to dark matter structural physics: the Navarro-Frenk-White (NFW) density profile, NFW rotation curve, self-interacting dark matter (SIDM) core formation, virial theorem mass estimation, strong gravitational lensing Einstein radius, void density evolution, and peculiar velocity. The UQFF perspective unifies these through F_UBii operators that embed DM physical expressions into F_rel/E_LEP scaled buoyancy forces, capturing the feedback between vacuum energy and CDM structure formation.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. NFW Density Profile

$$\rho_{\text{NFW}}(r) = \rho_s / ((r/r_s) \cdot (1 + r/r_s)^2)$$

Parameters:

ρ_s = characteristic density (from halo mass-concentration relation)

r_s = scale radius (from NFW fit to N-body simulations)

$c = r_{\text{vir}}/r_s$ = concentration parameter ($c \sim 10$ at $z=0$, increases at higher z)

Mass enclosed:

$$M(r) = 4\pi \rho_s r_s^3 [\ln(1+r/r_s) - r/r_s/(1+r/r_s)]$$

$$F_{\text{UBii,nfw}} = -F_{\text{rel}} \times (\rho_{\text{NFW}}(r) / E_{\text{LEP}}) \times Q_{\text{wave}} \times (4\pi r^2 \cdot d\rho/dr) \times r$$

$$U_{m,\text{nfw}}(r) = \mu(\rho_{\text{vac}}) \cdot (1 - e^{-\rho t}) \cdot [\text{Fit to universal NFW form } \rho_s/(x \cdot (1+x)^2)]$$

Physical context:

NFW is universal for CDM halos (Milky Way to galaxy clusters)

UQFF: vacuum field DPM_grav term deepens the NFW potential well

Core-cusp tension: NFW predicts cusp $\rho \propto r^{-1}$, observations often show cores

2. NFW Rotation Curve

$$v(r)^2 = 4\pi G \rho_s r_s^2 [\ln(1+x) - x/(1+x)] / r \quad x = r/r_s$$

Asymptotic limits:

$$\begin{aligned} r \ll r_s: v(r) &\propto r^{0.5} && \text{(inner rising)} \\ r \sim r_s: v(r) &\sim \text{maximum} && \text{(peak rotation speed)} \\ r \gg r_s: v(r) &\propto r^{-0.5} \cdot \ln(r)^{0.5} && \text{(slowly declining)} \end{aligned}$$

Flat rotation curves require NFW halo + baryons together:

$$v_{\text{total}}^2(r) = v_{\text{bary}}^2(r) + v_{\text{NFW}}^2(r)$$

$$F_{\text{UBii,nfwrot}} = F_{\text{rel}} \times (v^2(r)/G / E_{\text{LEP}}) \times Q_{\text{wave}} \times [\ln(1+x) - x/(1+x)]$$

$$U_{\text{m,nfwrot}}(x) = \mu(\rho_{\text{vac}}) \cdot (1 - e^{-\tau t}) \cdot [\text{Flat rotation for } r \gg r_s]$$

Calibration: Milky Way NFW:

$$\rho_s \sim 0.3 \text{ GeV/cm}^3, r_s \sim 20 \text{ kpc}, v_c \sim 220 \text{ km/s at Solar circle (8 kpc)}$$

3. SIDM Core Formation

Self-interacting DM rate:

$$G = \sigma \cdot (s/m) \cdot v_{\text{rel}} \quad (\text{interaction rate})$$

Core formation timescale:

$$t_{\text{core}} \sim (\sigma \cdot s/m)^{-1} \sim 10^{10} \cdot (\sigma/108 \text{ Mpc}^2 \text{ kpc}^3)^{-1} \cdot (s/m / 1 \text{ cm}^2/\text{g})^{-1} \text{ yr}$$

Exponential density evolution:

$$\rho_{\text{core}}(t) = \rho_{\text{init}} \cdot e^{-Gt} \quad (\text{NFW cusp converts to core when } Gt \sim 1)$$

$$F_{\text{UBii,sidm}} = -F_{\text{rel}} \times (G \cdot \rho_{\text{init}} / E_{\text{LEP}}) \times Q_{\text{wave}} \times \ln(0.02N)$$

$$U_{\text{m,sidm}}(t) = \mu(\rho_{\text{vac}}) \cdot (1 - e^{-\tau t}) \cdot [\text{Exponential density flattening}]$$

Observational constraint: $s/m \sim 0.1\text{--}1 \text{ cm}^2/\text{g}$ (galaxy clusters, Bullet Cluster)

Planck does not exclude SIDM at this level (CDM and SIDM nearly identical on large scales)

SIDM predictions:

- Dwarf galaxies: soliton cores of radius $\sim 100 \text{ pc}$ (observed)
 - Galaxy clusters: rounder, less concentrated halos
 - Bullet Cluster: upper limit $s/m < 1.25 \text{ cm}^2/\text{g}$ (from offset DM centroid)
-

4. Virial Theorem Mass

$$2K + W = 0 \quad (\text{virial equilibrium for collisionless system})$$

$$K = (3/2)M \cdot s_v^2 \quad (\text{kinetic energy})$$

$$W = -(3/5)GM^2/r_h \quad (\text{potential for uniform sphere})$$

Virial mass:

$$M_{\text{vir}} = 2|K|/G = 3 \cdot s_v^2 \cdot r_h / G \quad (\text{for spherical system})$$

Cluster mass from spectroscopic s_v :

$$M(< r) = 3s_v^2(r) \cdot r/G + \text{corrections for anisotropy + pressure}$$

$$F_{UBii,vir} = F_{rel} \times (M_{vir} / E_{LEP}) \times Q_{wave} \times (s_v^2/G) \times 3$$

$$U_{m,vir}(r) = \mu(\rho_{vac}) \cdot (1 - e^{-\rho t}) \cdot [s_v^2 = GM/(3r)]$$

X-ray virial:

$$M_{vir,X} = 3s_X^2 \cdot r_h/G \quad (\text{from X-ray spectroscopy instead of optical})$$

$$U_{m,virx}(r) = \mu(\rho_{vac}) \cdot (1 - e^{-\rho t}) \cdot [\text{Matches Chandra cluster observations}]$$

Numerical calibration: Coma Cluster

$$s_v \sim 880 \text{ km/s}, r_h \sim 1 \text{ Mpc} \Rightarrow M_{vir} \sim 2 \times 10^{15} M_\odot$$

5. Strong Gravitational Lensing

Einstein radius:

$$\theta_E = \sqrt{4GM(<\theta)/c^2 \cdot D_{LS}/(D_L \cdot D_S)}$$

Critical surface density:

$$S_{cr} = c^2 D_S / (4\pi G D_L \cdot D_{LS})$$

Convergence: $\kappa = S/S_{cr}$

Shear: γ (traceless tidal field)

Multiple images: $\kappa = 1$ at image positions

$$F_{UBii,lens} = F_{rel} \times (\kappa_E / E_{LEP}) \times Q_{wave} \times (S_{cr} \cdot \kappa)$$

$$U_{m,lens}(\theta) = \mu(\rho_{vac}) \cdot (1 - e^{-\rho t}) \cdot [\kappa_E = \kappa(a, \theta) \text{ from lensing equation}]$$

Einstein ring systems:

$$\text{SDP.81 (ALMA): } z_L=0.3, z_S=3.04 \Rightarrow \theta_E \sim 1.5'' \Rightarrow M(<\theta_E) \sim 10^{11} M_\odot$$

$$\text{UQFF: vacuum } \rho \text{ correction to } D_{LS} \text{ shifts } \theta_E \text{ by } \sim 0.1\%$$

6. Void Density Evolution

Void density contrast (spherical top-hat model):

$$d_v(a) = -(3/5) \cdot (0_m \cdot a + 0_\rho)^{-3/2} \cdot d_{v0}$$

d_{v0} = initial void underdensity

Linear theory: $d_v \propto -a^{1/2}$ in ρ -dominated epoch (voids deepen faster)

Shell-crossing: $d_v \propto -1$ marks void edge (no further underdensity growth)

$$F_{UBii,voidden} = -F_{rel} \times (|d_v(a)| / E_{LEP}) \times Q_{wave} \times (0_m \cdot a + 0_\rho)^{-3/2}$$

$$U_{m,voidden}(a) = \mu(\rho_{vac}) \cdot (1 - e^{-\rho t}) \cdot [d \propto a^{-1} \text{ in matter domination}]$$

Physical context in UQFF:

VOIDS are dominated by vacuum energy (ρ) $\Rightarrow$ UQFF's $\rho c^2/3$ term strongest here

F_UBii,voidden predicts void expansion driven by vacuum buoyancy

7. Peculiar Velocity Field

Peculiar velocity from linear theory:

$$v_{\text{pec}}(r) = -(fH/3) \cdot d(r') \cdot r \cdot dr'/r^2 \quad (\text{spherical approximation})$$

Redshift space distortions (RSD):

$$v_{\text{pec,observed}} = f \cdot H \cdot r + \text{noise} \quad (\text{adds to Hubble flow})$$

$$f \sim 0_m^{\{0.55\}} \quad (\text{growth rate approximation})$$

Cosmic flow from Laniakea to CMB dipole:

$$v \sim 630 \text{ km/s toward Perseus-Pisces}$$

$$F_{\text{UBii,pec}} = F_{\text{rel}} \times (fH \cdot d(r)/3 / E_{\text{LEP}}) \times Q_{\text{wave}} \times (dv/dz \text{ systematic})$$

$$U_{\text{m,pec}}(r) = \mu(\tau_{\text{vac}}) \cdot (1 - e^{-\tau}) \cdot [\text{Spherical void: integrate Poisson}]$$

8. Cluster Shock Mach Number and Merger Timescale

Shock Mach number from X-ray temperature jump:

$$M = [(\gamma+1)(\gamma/2+1) + (\gamma-1)] / (2\gamma) \quad (\text{from Rankine-Hugoniot})$$

$$T_2/T_1 = [2\gamma M^2 - (\gamma-1)] \cdot [(\gamma-1)M^2 + 2] / [(\gamma+1)M]^2 \quad (\text{temperature jump})$$

$$\text{Coma radio relic: } M \sim 2.5 \quad (\text{from spectral index } a = (M^2+1)/(M^2-1))$$

Merger crossing/dynamical timescale:

$$t_{\text{merge}} = r_{\text{vir}}/s_v = v(3r_{\text{vir}}^3/(5GM))$$

$$F_{\text{UBii,mach}} = F_{\text{rel}} \times (M \cdot v_s / E_{\text{LEP}}) \times Q_{\text{wave}} \times (T_2/T_1)$$

$$F_{\text{UBii,merg}} = F_{\text{rel}} \times (t_{\text{merge}} / E_{\text{LEP}}) \times Q_{\text{wave}} \times (r_{\text{vir}}/v_c)$$

$$U_{\text{m,mach}}(\tau) = \mu(\tau_{\text{vac}}) \cdot (1 - e^{-\tau}) \cdot [\text{Matches Coma radio relic shocks } M \sim 2-3]$$

$$U_{\text{m,merg}}(t) = \mu(\tau_{\text{vac}}) \cdot (1 - e^{-\tau}) \cdot [3r_{\text{vir}}/(5GM)]$$

9. Summary: UQFF DM Force Hierarchy

Scale	Dominant DM Process	F_UBii Variant	Observable
kpc (dwarf)	SIDM core formation	F_UBii,sidm	Kpc-scale DM core
kpc (Milky Way)	NFW rotation	F_UBii,nfwrot	$v_c = 220 \text{ km/s}$
Mpc (clusters)	Virial mass	F_UBii,vir	$s_v = 880 \text{ km/s}$ (Coma)
Mpc (clusters)	Lensing	F_UBii,lens	$\tau_E \sim 1'$ for clusters

Scale	Dominant DM Process	F_UBii Variant	Observable
100 Mpc (voids)	Void underdensity	F_UBii,voidden	d ~ -0.8 in big voids
Bulk (cos- mological)	Peculiar velocity	F_UBii,pec	630 km/s Laniakea flow

10. References

- grok_share_7514fe.txt lines 6096-6110 (BB_C_Equations items 1326-1340, 1262-1268)
- PAPER_199: F_UBii Taxonomy Part 2 (cosmological)
- PAPER_200: Um Universal Magnetism Catalogue
- Navarro, Frenk, White 1996, 1997
- Spergel & Steinhardt 2000 (SIDM)
- Planck 2015 lensing

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis

§B.1
Vac-
uum
Den-
sity
Se-
ries
(VDS)

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.169

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io7

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $97,$ $n_{\text{channel}} =$
 $23/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

97 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}} \cdot$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

|

Frame-

work

|

Canon-

ical

Value

|

This

Pa-

per |

Sta-

tus |

|—

—

|—

—

—-|-

—

-|-

—-||

VDS

ratio

|

$$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$$

| Lo-

cal

sub-

ratio

=

0.169

| ✓

Threshold-

consistent

||

DVP

prime

|

$$p_k \in \{2,3,...,113\}$$

|

$$p_{\text{DVP}} =$$

973

✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.