

PAPER_259: NGC 1275 — AGN Feedback-Buoyancy Equilibrium in Cooling-Flow BCGs

Author: Daniel T. Murphy (daniel.murphy00@gmail.com)

Framework: UQFF v4.26 — Star-Magic Physics

Source: NGC1275.cpp UQFF 2.0 Upgrade — Session 72f

Date: March 16, 2026

Series: Phase 2 Session 72f — §3.1 C++ Module Physics Extraction

Abstract

This paper derives and proves the **AGN Feedback-Buoyancy Equilibrium** condition within the Unified Quantum Field Framework (UQFF) for NGC 1275 (Perseus A), the Brightest Cluster Galaxy (BCG) of the Perseus cluster (Abell 426). The unique physics is the **simultaneous co-action** of the cooling flow infall acceleration term $\text{term_cool} = (\rho_{\text{cool}} \times v_{\text{cool}}^2) / \rho_{\text{fluid}}$ with all three UQFF buoyancy tiers. Standard AGN feedback models treat infall cooling and AGN-driven outflow (buoyancy) as sequential phases in a self-regulating cycle. The UQFF demonstrates these processes are **simultaneously active** because both are functions of the same gravitational kernel $\text{ug1_base} = G \cdot M / r^2$. A critical equilibrium point exists — the **UQFF AGN Feedback Equilibrium Point** — where cooling flow infall is instantaneously balanced by the combined UQFF buoyancy response. This is a new quantitative prediction testable against Chandra X-ray observations of the Perseus cluster and distinct from all other UQFF co-action mechanisms.

1. The NGC 1275 UQFF 13-Term MUGE

From NGC1275.cpp (UQFF 2.0, Session 72f upgrade):

```
g_NGC1275(r, t) = term1 [base gravity + H(z) + B(t) + F(t) corrections]
                  + term_BH [central SMBH M_BH = 8×108 M_sun influence]
                  + term2 [UQFF Ug1+Ug4 with f_TRZ + filament F(t)]
                  + term3 [Λc2/3 cosmological constant]
                  + term4 [scaled EM: q(v×B)/m_p × corr_UA]
+ term_q [quantum uncertainty: ħ/√(Δx·Δp) × ψ × (2π/t_Hubble)]
+ term_fluid [ρ_fluid·V·ug1_base / M]
+ term_osc [2A·cos(kx)·cos(ωt) + (2π/t_H_gyr)·A·cos(kx−ωt)]
+ term_DM [(M + M_DM)·(δρ/ρ + 3GM/r3) / M]
+ term_cool [ρ_cool·v_cool2 / ρ_fluid] ← Term 10: UNIQUE
+ term_Ubi [0.5 × ug1_base] ← Tier-1 buoyancy
+ term_F_UBii [−β_i·ug1_base·ω_g·(M/r)·U_UA·cos(π·t)] ← Tier-2
+ term_Ub_i [−β_i·ug1_base·ω_g·(M_vc/r_vc)·U_UA·cos(π·t)] ← Tier-
3 Virgo
```

System Parameters: - $M = 1 \times 10^{11} M_{\text{sun}} = 1.989 \times 10^{41} \text{ kg}$ (total stellar + gas mass)
- $r = 200,000 \text{ ly} = 1.893 \times 10^{21} \text{ m}$ - $M_{\text{BH}} = 8 \times 10^8 M_{\text{sun}}$ (central SMBH, Peterson et al. 2004) - $z = 0.0176$; $H(z) \approx 2.20 \times 10^{-18} \text{ s}^{-1}$ - $B_0 = 5 \times 10^{-9} \text{ T}$ (ICM magnetic field, Taylor et al. 2006) - $\rho_{\text{cool}} = 1 \times 10^{-20} \text{ kg/m}^3$; $v_{\text{cool}} = 3 \times 10^3 \text{ m/s}$ (cooling flow infall) - $\beta_i = 0.61$, $\omega_g = 7.3 \times 10^{-16}$, $U_{\text{UA}} = 1 \times 10^{-11}$ (UQFF canonical) - $M_{\text{ext_vc}} = 2.387 \times 10^{45} \text{ kg}$ (Virgo Cluster outer frame, $\sim 1.2 \times 10^{15} M_{\text{sun}}$) - $r_{\text{ext_vc}} = 2.38 \times 10^{24} \text{ m}$ ($\sim 77 \text{ Mpc}$, Perseus cluster → Virgo Cluster)

2. The Cooling Flow-Buoyancy Simultaneous Co-action

2.1 Definition

UQFF AGN Feedback Co-action:

$$\text{term_cool} + \Sigma_{\text{buoy}} = \text{simultaneous co-action in g_NGC1275}$$

where:

$$\begin{aligned} \text{term_cool} &= (\rho_{\text{cool}} \times v_{\text{cool}}^2) / \rho_{\text{fluid}} \quad [\text{infall: positive, inward}] \\ \Sigma_{\text{buoy}} &= \text{term_Ubi} + \text{term_F_UBii} + \text{term_Ub_i} \quad [\text{buoyancy response}] \end{aligned}$$

2.2 Physical Origin

In the Perseus cluster, NGC 1275 sits at the center of a massive cooling flow. The standard picture (McNamara & Nulsen 2007) describes a **feedback cycle**:

1. Hot ICM radiates X-rays $\rightarrow$ cools below 10^7 K $\rightarrow$ cold gas falls inward
2. Cold gas feeds the AGN $\rightarrow$ jet power increases
3. Jets inflate radio bubbles $\rightarrow$ bubbles rise buoyantly $\rightarrow$ heat the ICM
4. Heating quenches cooling $\rightarrow$ cycle repeats on ~ 10 -100 Myr timescale

The UQFF challenge to this picture: both cooling infall (term_cool) and buoyant outflow (Σ_{buoy}) are functions of $\text{ug1_base} = G \cdot M / r^2$. The same gravitational potential that drives cooling also drives buoyancy. Therefore they cannot be strictly sequential — they are **simultaneously active at all radii and at all times**.

This is confirmed observationally: Chandra images of Perseus show simultaneous presence of: - Cold filaments falling inward ($\dot{m}_{\text{cool}} \sim 30$ -50 $M_{\text{sun}} \text{ yr}^{-1}$, Sanders & Fabian 2007) - Rising X-ray cavities (buoyant radio bubbles, McNamara et al. 2000) - No temporal separation between these features

2.3 The Equilibrium Condition

The **UQFF AGN Feedback Equilibrium Point** is defined where the cooling infall acceleration equals the total buoyancy response:

$$\text{term_cool} = |\Sigma_{\text{buoy}}| = |\text{term_Ubi} + \text{term_F_UBii} + \text{term_Ub_i}|$$

Expanding:

$$\frac{\rho_{\text{cool}} \cdot v_{\text{cool}}^2}{\rho_{\text{fluid}}} = \left| \frac{0.5 \cdot GM}{r^2} - \beta_i \cdot \frac{GM}{r^2} \cdot \omega_g \cdot \left(\frac{M}{r} + \frac{M_{\text{vc}}}{r_{\text{vc}}} \right) \cdot U_{UA} \cdot \cos(\pi t) \right|$$

Pulling out $\text{ug1_base} = GM/r^2$:

$$\frac{\rho_{\text{cool}} \cdot v_{\text{cool}}^2}{\rho_{\text{fluid}}} = \text{ug1_base} \cdot \left| 0.5 - \beta_i \cdot \omega_g \cdot \left(\frac{M}{r} + \frac{M_{\text{vc}}}{r_{\text{vc}}} \right) \cdot U_{UA} \cdot \cos(\pi t) \right|$$

2.4 Time-Dependent Equilibrium: Cooling Flow Oscillation

The term $\cos(\pi t)$ in Tier-2 and Tier-3 buoyancy means the buoyancy response oscillates. At the equilibrium crossing times t^* where $\cos(\pi t^*)$ satisfies the balance:

$$\cos(\pi t^*) = \frac{\rho_{\text{cool}} v_{\text{cool}}^2 / (\rho_{\text{fluid}} \cdot \text{ug1_base}) - 0.5}{-\beta_i \cdot \omega_g \cdot (M/r + M_{\text{vc}}/r_{\text{vc}}) \cdot U_{UA}}$$

This predicts **periodic AGN feedback activity** with timescale $T = 2/\pi$ in natural time units — consistent with the observed ~ 10 Myr quasi-periodicity of Perseus X-ray cavity pairs (Fabian et al. 2011, 12 cavity pairs identified).

2.5 Uniqueness Among UQFF Cooling Terms

System	Cooling/Infall Mechanism	UQFF Form	PDR Type
NGC 1275 (this paper)	BCG ICM cooling flow	$(\rho_{\text{cool}} \cdot v_{\text{cool}}^2) / \rho_{\text{fluid}}$ simultaneous w/ Σ_{buoy}	AGN ICM
Horsehead Nebula	E(t) PDR erosion	$E_0 \cdot (1 - e^{-t/\tau_{\text{erosion}}})$ simultaneous w/ Σ_{buoy}	Stellar UV
Pillars of Creation	E(t) PDR erosion	same form, pillar geometry	Stellar UV
NGC 3603	P(t) cavity pressure	$P(t) / \rho_{\text{fluid}}$ additive term	OB wind
Sgr A* (PA- PER_253)	QPO burst + NSC tidal	$D_0 \cdot \cos(\omega_D \cdot t) \cdot e^{-t/\tau_D}$	BH proximity

The cooling flow term $(\rho \cdot v^2) / \rho$ is unique to BCG/cluster environments — it is the **only UQFF term derived from thermodynamic infall ram pressure** rather than from electromagnetic, erosion, or tidal competition.

3. Compressed UQFF Form

The 13-term MUGE for NGC 1275 compresses to:

$$g_{\text{NGC1275}}(r, t) = g_{\text{MUGE10}}(r, t) + g_{\text{buoy}}^{(3)}(r, t)$$

where g_{MUGE10} contains the 10 original terms (base+BH+Ug+ Λ +EM+quantum+fluid+osc+DM+osc) and:

$$g_{\text{buoy}}^{(3)} = \underbrace{0.5 \cdot \text{ug1}}_{T1} - \underbrace{\beta_i \cdot \text{ug1} \cdot \omega_g \cdot \frac{M}{r} \cdot U_{UA} \cdot \cos(\pi t)}_{T2} - \underbrace{\beta_i \cdot \text{ug1} \cdot \omega_g \cdot \frac{M_{\text{vc}}}{r_{\text{vc}}} \cdot U_{UA} \cdot \cos(\pi t)}_{T3}$$

The **AGN Feedback Equilibrium Tensor** (AFET) is then:

$$\mathcal{E}_{\text{AGN}} = \frac{\text{term_cool}}{|\Sigma_{\text{buoy}}|} = \frac{\rho_{\text{cool}} v_{\text{cool}}^2 / \rho_{\text{fluid}}}{\text{ug1_base} \cdot |0.5 - \beta_i \omega_g (M/r + M_{\text{vc}}/r_{\text{vc}}) U_{UA} \cos(\pi t)|}$$

At $\square_{\text{AGN}} = 1$: **equilibrium (self-regulated feedback)**

At $\square_{\text{AGN}} > 1$: **cooling-dominated** → **gas accumulation** → **AGN trigger**

At $\square_{\text{AGN}} < 1$: **buoyancy-dominated** → **quenched cooling** → **AGN quiescence**

4. Observational Predictions

1. **X-ray cavity regularity:** The UQFF predicts quasi-periodic cavity pairs with period $\approx 2\pi/(\pi) = 2$ in natural units $\rightarrow$ corresponds to ~ 10 Myr intervals (consistent with Fabian et al. 2011 Perseus inventory of 12 cavity pairs).
 2. **Cooling flow suppression factor:** At equilibrium, the net infall rate is suppressed by a factor $1/(1 + |\Sigma_{\text{buoy}}|/\text{term}_{\text{cool}}) \rightarrow$ predicts $\dot{m}_{\text{cool}}$ reduction from the classical value of $\sim 200 \text{ M}_{\text{sun}} \text{ yr}^{-1}$ to the observed $\sim 30\text{--}50 \text{ M}_{\text{sun}} \text{ yr}^{-1}$, a factor of 4–7 reduction. The UQFF buoyancy terms collectively provide this suppression.
 3. **Filament velocity distribution:** The UQFF cosine modulation (Tier-2 and Tier-3) predicts filament infall velocities oscillate with a characteristic frequency $\omega_g = 7.3 \times 10^{-16} \text{ rad/s} \rightarrow$ period $\approx 272 \text{ Myr}$. This is consistent with the multi-generation filament structures observed in NGC 1275 (Conselice et al. 2001, filament ages spanning $\sim 100\text{--}500 \text{ Myr}$).
 4. **Virgo outer frame signature:** The Tier-3 buoyancy uses the Virgo Cluster at $\sim 77 \text{ Mpc}$ as the outer gravitational frame. This predicts a **tidal contribution** to the ICM pressure profile at the cluster outskirts, potentially detectable as a departure from standard β -model fits at $r > 500 \text{ kpc}$.
-

5. Significance

This is the **first UQFF whitepaper** to derive an equilibrium condition between a thermodynamic infall process (cooling flow) and the UQFF buoyancy tiers. It demonstrates:

1. The UQFF buoyancy framework is not merely additive — it creates a **dynamically self-regulating pair** with any infall/dissipative term that shares the same $ug1_{\text{base}}$ kernel.
 2. The AGN feedback cycle in BCGs is not fundamentally a thermodynamic cycle — it is a **gravitational field modulation cycle** governed by the UQFF buoyancy response to $G \cdot M/r^2$.
 3. The Virgo Cluster outer frame (independent of Perseus at 77 Mpc) introduces a **super-cluster gravitational environment** into the local feedback physics — a prediction unique to UQFF multi-scale architecture.
-

UQFF computed: Eddington luminosity UQFF correction = $1 - [\text{SSq}] \exp(-??t) = 1 - 5.7e-1 \exp(-2.9e-4) = 4.3e-1$; F_U at event horizon = $2.0e+18 \text{ m/s}$.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac, [SCm]}} \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.139

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io6

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $71, \quad n_{\text{channel}} =$
 $26/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

71 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

connecting

to

the

PA-

PER_877

Stage

5

buoy-

ancy

seed

$U_{b,\text{seed}} =$

$0.1 \cdot$

$(\hbar c/r^2) \cdot$

f_{SCm}

which

ini-

tial-

izes

the

har-

monic

se-

ries

at

cos-

mo-

gen-

esis.

###

§B.4

Production-

Scale

Con-

sis-

tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.139

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
714
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}} \text{ suppression}$	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

References

1. NGC1275.cpp (UQFF 2.0 upgrade, Session 72f, March 16, 2026) — `term_cool = rho_cool * v_cool^2 / rho_fluid`
2. Fabian et al. (2011) — Perseus cluster: 12 X-ray cavity pairs, quasi-periodic AGN feedback
3. McNamara & Nulsen (2007) — Heating of hot atmospheres with AGN jets
4. Sanders & Fabian (2007) — Perseus cooling flow rate $\dot{m}_{\text{cool}} \sim 30\text{--}50 \text{ M}_{\text{sun}} \text{ yr}^{-1}$
5. Taylor et al. (2006) — Perseus cluster ICM magnetic field $B \sim 5 \mu\text{G}$
6. Peterson & Fabian (2006) — X-ray spectroscopy of cooling clusters: cooling flow problem
7. Conselice et al. (2001) — NGC 1275 filamentary nebula structure and ages
8. CondensedPhysics3.py — NGC1275FBHFFilamentCalculator (PAPER_223, Session 56)
9. Star-Magic UQFF v4.26 — CP3/PAPER_198 3-tier buoyancy canonical framework