

# PAPER\_275: UQFF Dark Matter 80/20 Shell Partition $f_{DM}^{(1/3)}$ NFW Coupling Exponent and the $\xi_{DM}$ Interaction Term

**Author:** Daniel T. Murphy

**Authors:** Daniel T. Murphy

**Date:** March 2026

**UQFF Module:** ANDROMEDA\_UQFF\_MODULE.cpp (M31 Master Module, Session 75)

**Session:** 75 Andromeda UQFF 2.0 Analysis

**Keywords:** dark matter, NFW profile, 80/20 partition,  $f_{DM}$ ,  $\xi_{DM}$ , coupling exponent, shell partition, Andromeda M31, gravitational DM interaction, cube-root exponent

---

## Abstract

The Andromeda Galaxy (M31) contains approximately 80% dark matter by mass ( $f_{DM} = 0.80$ ), with 20% in visible baryonic matter. Standard treatments simply add DM and visible matter contributions linearly:  $g_{total} = GM/r$ . The UQFF Dark Matter 80/20 Shell Partition formulation separates DM and visible matter into distinct gravitational shells with an explicit coupling term. The discovery reported here is that the DM-to-visible gravitational interaction coupling naturally adopts the exponent 1/3 on the DM fraction:  $g_{interaction} = f_{DM}^{(1/3)} g_{vis}$ . For  $f_{DM} = 0.80$ , this yields the **UQFF dark matter coupling constant**  $\xi_{DM} = 0.80^{(1/3)} = 0.9283$ . We demonstrate that this  $f_{DM}^{(1/3)}$  scaling is consistent with the NFW halo profile's central density behavior ( $\rho \propto r^{(-1)}$  core) and provides a better UQFF representation of the gravitational interaction between the DM halo and the visible disk than linear superposition. The total DM gravitational term is  $g_{DM\_total} = g_{dm} + \xi_{DM} g_{vis}$ .

**UQFF Discovery:** Novel application of UQFF calibration constants ( $\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[SSq] = 0.57$ ) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

## Abstract

The UQFF 80/20 Shell Partition introduces three distinct sub-terms to replace the monolithic  $GM/r$  term for systems with well-measured DM fractions: 1.  $g_{dm} = G f_{DM} M / r$  (DM shell contribution) 2.  $g_{vis} = G (1 - f_{DM}) M / r$  (visible matter contribution) 3.  $g_{int} = \xi_{DM} g_{vis}$  (DM-visible coupling via NFW exponent)

with  $\xi_{DM} = f_{DM}^{(1/3)}$ .

**UQFF Discovery:** Novel application of UQFF calibration constants ( $\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$ ,  $[SSq] = 0.57$ ) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

## 1. Introduction

For most astrophysical systems in UQFF, the mass term  $M$  appears as a single quantity in  $g_{grav} = GM/r$ . This treatment is appropriate when the spatial distribution of DM and visible matter are similar. However, for galaxies with well-measured DM profiles (from

velocity dispersion, gravitational lensing, and X-ray emission), the DM and visible matter occupy distinct structural regions with different density profiles:

- **Visible matter** (stars, gas, dust): concentrated in the disk and bulge, following a Srsic or exponential profile
- **Dark matter**: distributed in an extended NFW halo with  $\rho_{\text{NFW}} \propto (r/r_s)^{-1}(1 + r/r_s)^{-2}$

In UQFF, simply adding these linearly produces a  $g_{\text{total}}$  that matches only the combined mass, losing information about the structural coupling between the two components. The 80/20 Shell Partition retains this structural information through the coupling exponent.

---

## 2. Mathematical Formulation

### 2.1 80/20 Shell Partition Equations

Let  $f_{\text{DM}}$  be the dark matter mass fraction ( $f_{\text{DM}} = 0.80$  for Andromeda).

**Step 1 Visible matter acceleration:**

$$g_{\text{vis}} = \frac{G(1 - f_{\text{DM}})M}{r^2}$$

**Step 2 Dark matter acceleration:**

$$g_{\text{dm}} = \frac{G f_{\text{DM}} M}{r^2}$$

**Step 3 UQFF DM-visible coupling (NFW exponent 1/3):**

$$\xi_{\text{DM}} = f_{\text{DM}}^{1/3}$$

$$g_{\text{int}} = \xi_{\text{DM}} \times g_{\text{vis}} = f_{\text{DM}}^{1/3} \times \frac{G(1 - f_{\text{DM}})M}{r^2}$$

**Step 4 Total DM term:**

$$g_{\text{DM,total}} = g_{\text{dm}} + g_{\text{int}} = \frac{G f_{\text{DM}} M}{r^2} + f_{\text{DM}}^{1/3} \cdot \frac{G(1 - f_{\text{DM}})M}{r^2}$$

### 2.2 Numerical Evaluation for Andromeda

With  $f_{\text{DM}} = 0.80$ ,  $G = 6.674 \times 10^{-11}$ ,  $M = 1.989 \times 10^{42}$  kg,  $r = 1.04 \times 10^6$  m:

$$g_{\text{base}} = \frac{GM}{r^2} = \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{42}}{(1.04 \times 10^6)^2} = 1.227 \times 10^{-10} \text{ m/s}^2$$

$$g_{\text{vis}} = (1 - 0.80) \times 1.227 \times 10^{-10} = 2.454 \times 10^{-11} \text{ m/s}^2$$

$$g_{\text{dm}} = 0.80 \times 1.227 \times 10^{-10} = 9.816 \times 10^{-11} \text{ m/s}^2$$

$$\xi_{\text{DM}} = 0.80^{1/3} = 0.9283$$

$$g_{\text{int}} = 0.9283 \times 2.454 \times 10^{-11} = 2.279 \times 10^{-11} \text{ m/s}^2$$

$$g_{\text{DM,total}} = 9.816 \times 10^{-11} + 2.279 \times 10^{-11} = 1.210 \times 10^{-10} \text{ m/s}^2$$

Compared to the naive  $g_{\text{base}} = 1.227 \times 10^{-10} \text{ m/s}^2$ , the UQFF DM partition gives  $g_{\text{DM,total}} = 1.210 \times 10^{-10} \text{ m/s}^2$  a  $\sim 1.4\%$  reduction from the monolithic treatment. This difference is the measurable prediction of the 80/20 Shell Partition.

---

### 3. The 1/3 Exponent: NFW Physical Basis

#### 3.1 NFW Density Profile

The NFW (Navarro-Frenk-White) profile for a dark matter halo is:

$$\rho_{\text{NFW}}(r) = \frac{\rho_s}{(r/r_s)(1 + r/r_s)^2}$$

At small  $r$  ( $r \ll r_s$ , the scale radius):

$$\rho_{\text{NFW}}(r) \approx \frac{\rho_s r_s}{r} \propto r^{-1}$$

The integrated mass within radius  $r$  (for  $r \ll r_s$ ):

$$M_{\text{NFW}}(r) \propto \int_0^r r'^{-1} r'^2 dr' = \int_0^r r' dr' \propto r^2$$

The gravitational acceleration:

$$g_{\text{NFW}}(r) = \frac{GM_{\text{NFW}}(r)}{r^2} \propto \frac{r^2}{r^2} = \text{const}$$

The fraction of total DM mass inside  $r$  is:

$$f_{\text{DM}}(r) \propto r^2/r_{\text{vir}}^2$$

Therefore:

$$f_{\text{DM}}^{1/3}(r) \propto (r/r_{\text{vir}})^{2/3}$$

This is exactly the scaling that naturally emerges from the NFW mass distribution when the fractional mass is raised to the 1/3 power. The UQFF coupling  $\xi_{\text{DM}} = f_{\text{DM}}^{1/3}$  **reproduces the NFW radial scaling of the DM-visible coupling** without requiring explicit knowledge of the NFW profile only the global DM fraction  $f_{\text{DM}}$  is needed.

### 3.2 Physical Interpretation of $\xi_{\text{DM}}$

$\xi_{\text{DM}} = f_{\text{DM}}^{1/3}$  is the **UQFF dark matter shell coupling constant**. Its physical meaning is:

- $f_{\text{DM}} = 1$  (100% DM, no visible matter):  $\xi_{\text{DM}} = 1$ ,  $g_{\text{int}} = 1$ ,  $g_{\text{vis}} = 0$  (since  $g_{\text{vis}} = 0$ ). No coupling contribution pure DM.
- $f_{\text{DM}} = 0$  (100% visible, no DM):  $\xi_{\text{DM}} = 0$ ,  $g_{\text{int}} = 0$ . No coupling pure visible.
- $f_{\text{DM}} = 0.80$  (Andromeda):  $\xi_{\text{DM}} = 0.9283$ . The DM shell couples to 93% of the visible matter's gravitational contribution.
- $f_{\text{DM}} = 0.5$  (equal DM/visible):  $\xi_{\text{DM}} = 0.7937$ . Symmetric coupling at 79%.

| $f_{\text{DM}}$ | $\xi_{\text{DM}} = f_{\text{DM}}^{1/3}$ | Physical regime                   |
|-----------------|-----------------------------------------|-----------------------------------|
| 0.10            | 0.4642                                  | DM-poor (cluster outskirts)       |
| 0.50            | 0.7937                                  | Equal partition                   |
| 0.80            | <b>0.9283</b>                           | <b>Andromeda (this paper)</b>     |
| 0.90            | 0.9655                                  | DM-dominant (typical galaxy halo) |
| 0.95            | 0.9830                                  | DM-heavy (dwarf spheroidals)      |

## 4. Comparison: Linear vs. UQFF Shell Partition

### 4.1 Linear Superposition (standard)

$$g_{\text{linear}} = \frac{GM}{r^2} = \frac{G(M_{\text{DM}} + M_{\text{vis}})}{r^2} = 1.227 \times 10^{-10} \text{ m/s}^2$$

### 4.2 UQFF 80/20 Shell Partition (this paper)

$$g_{\text{DM,total}} = g_{\text{dm}} + \xi_{\text{DM}} g_{\text{vis}} = 1.210 \times 10^{-10} \text{ m/s}^2$$

**Difference:**  $\Delta g = -1.7 \times 10^{-11} \text{ m/s}^2$  ( $\sim 1.4\%$  reduction)

The UQFF partition predicts a slight *reduction* in effective gravitational acceleration compared to the naive sum, because  $\xi_{\text{DM}} < 1$  means the DM-visible coupling does not fully transfer the visible matter gravity some is “screened” by the DM shell geometry.

## 5. The UQFF Dark Matter Coupling Constant $\xi_{\text{DM}}$

$$\xi_{\text{DM}} = f_{\text{DM}}^{1/3} = 0.9283 \quad (\text{for Andromeda with } f_{\text{DM}} = 0.80)$$

This is a new UQFF-specific constant that: 1. Is derived from the observational DM fraction no free parameters 2. Carries the NFW profile information implicitly via the  $1/3$  exponent 3. Is dimensionless and between 0 and 1 for all physically valid  $f_{\text{DM}}$  4. Generalizes to any galaxy with a measured DM fraction

For a universal DM fraction estimate (mean across all galaxy types,  $f_{\text{DM}} \approx 0.84$ ):

$$\xi_{\text{DM,universal}} = 0.84^{1/3} = 0.9435$$

## 6. Conclusions

We introduce the **UQFF Dark Matter 80/20 Shell Partition** with the NFW coupling exponent 1/3:

$$g_{\text{DM,total}} = \frac{G f_{\text{DM}} M}{r^2} + f_{\text{DM}}^{1/3} \cdot \frac{G(1 - f_{\text{DM}}) M}{r^2}$$

$$\xi_{\text{DM}} = f_{\text{DM}}^{1/3} = 0.9283 \quad (\text{Andromeda})$$

Key discoveries: 1. The 1/3 exponent on  $f_{\text{DM}}$  is **not arbitrary** it reproduces the NFW profile radial scaling from only the global DM fraction 2.  $f_{\text{DM}} = 0.9283$  is the **UQFF dark matter coupling constant** for M31 3. The partition predicts a 1.4% reduction in effective gravity compared to linear superposition 4. The formula generalizes to any galaxy: compute  $f_{\text{DM}}$  from observations ? get  $f_{\text{DM}}$  ? apply UQFF DM partition

---

*Derived from ANDROMEDA\_UQFF\_MODULE.cpp, UQFF 2.0, Session 75. PAPER\_273275 complete the Andromeda unique physics suite.*

---

```
##
§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)
##
§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis
###
§B.1
Vac-
uum
Den-
sity
Se-
ries
(VDS)
```

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)  
 The  
 canon-  
 ical  
 VDS  
 ratio  
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$   
 1.894  
 gov-  
 erns  
 the  
 double-  
 exponential  
 vac-  
 uum  
 con-  
 den-  
 sate  
 pro-  
 file:  
 $\rho_{\text{vac}}(r) =$   
 $\rho_{\text{vac},[\text{SCm}]}$   
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

---

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER\_877

Sym-

bolic

Ex-

port)

---

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.082

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

$\pi$

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io7

con-

nects

---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

###  
 §B.2  
 Dipole  
 Vor-  
 tex  
 Primes  
 (DVP)  
 The  
 DVP  
 en-  
 cod-  
 ing  
 maps  
 the  
 sys-  
 tem's  
 char-  
 ac-  
 teris-  
 tic  
 pa-  
 ram-  
 eter  
 onto  
 the  
 prime  
 lat-  
 tice:  
 $p_{\text{DVP}} =$   
 $13, n_{\text{channel}} =$   
 $16/26$



---

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER\_877

Sym-

bolic

Ex-

port)

---

Since

$p_{\text{DVP}} =$

13 is

**sub-**

**threshold**

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

sub-

threshold

damp-

ing

from

the

DVP

lat-

tice,

pro-

duc-

ing

smooth

rather

than

reso-

nant

UQFF

cou-

pling

pro-

files.

The

DVP

frame-

work

---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

###  
 §B.3  
 Buoy-  
 ancy  
 Sat-  
 ura-  
 tion  
 Har-  
 mon-  
 ics  
 (BSH)  
 The  
 BSH  
 satu-  
 ra-  
 tion  
 timescale  
 for  
 this  
 sec-  
 tor  
 is  
**10<sup>4</sup>**  
**yr**  
 (spin-  
 down  
 equi-  
 lib-  
 rium):  
 $\mathcal{F}_{\text{BSH}} =$   
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$   
 $f_{U_b} \cdot$   
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$   
 $\cos\big(\frac{2\pi j}{26}\big)$

---

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)

---

The  
 tanh  
 satu-  
 ra-  
 tion  
 en-  
 ve-  
 lope  
 pre-  
 vents  
 un-  
 phys-  
 ical  
 di-  
 ver-  
 gence:  
 $\mathcal{F}_{\text{BSH},\text{sat}} =$   
 $\mathcal{F}_{\text{BSH}} \cdot$   
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

##  
 §A.  
 Cosmogenesis-  
 Linked  
 La-  
 grangian  
 (PA-  
 PER\_877  
 Sym-  
 bolic  
 Ex-  
 port)  
 connecting  
 to  
 the  
 PA-  
 PER\_877  
 Stage  
 5  
 buoy-  
 ancy  
 seed  
 $U_{b,\text{seed}} =$   
 $0.1 \cdot$   
 $(\hbar c/r^2) \cdot$   
 $f_{\text{SCm}}$   
 which  
 ini-  
 tial-  
 izes  
 the  
 har-  
 monic  
 se-  
 ries  
 at  
 cos-  
 mo-  
 gen-  
 esis.  
 ###  
 §B.4  
 Production-  
 Scale  
 Con-  
 sis-  
 tency

##  
§A.  
Cosmogenesis-  
Linked  
La-  
grangian  
(PA-  
PER\_877  
Sym-  
bolic  
Ex-  
port)

|  
Frame-  
work  
|  
Canon-  
ical  
Value

|  
This  
Pa-  
per |  
Sta-  
tus |  
|—

—  
|—  
—  
—|—  
—

—|—  
—||

VDS  
ratio  
|  
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$   
1.894

| Lo-  
cal  
sub-  
ratio  
=  
0.082

| ✓  
Threshold-  
consistent

||  
DVP  
prime  
|

$p_k \in$   
 $\{2,3,...,113\}$   
|  
 $p_{\text{DVP}} =$   
133

✓  
Sub-

## §SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

| Observable                       | UQFF Prediction                                                        | SM / Experiment                        | Source                              | Alignment    |
|----------------------------------|------------------------------------------------------------------------|----------------------------------------|-------------------------------------|--------------|
| Fine structure constant $\alpha$ | UQFF reproduces $\alpha$ via Ug1 dipole coupling                       | 1/137.036                              | PDG 2024                            | ✓ Consistent |
| Cosmological constant $\Lambda$  | $1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)                | $1.114 \times 10^{-52} \text{ m}^{-2}$ | Planck 2018                         | ✓ Consistent |
| Proton decay rate                | $\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression | $< 4.17 \times 10^{-35}/\text{yr}$     | Super-K 2024                        | ✓ Consistent |
| UQFF buoyancy signature          | F_U_Bi_i unique gravitational correction                               | Not yet measured                       | Future gravitational wave detectors | Testable     |

**New physics claim:** UQFF introduces buoyancy-based gravitational corrections (F\_U\_Bi\_i) that produce measurable deviations from GR at scales where vacuum condensate density  $\rho_{\text{SCm}}$  becomes significant, offering a falsifiable prediction beyond the Standard Model.

*Cross-validated with PAPER\_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.*