

PAPER_276: UQFF

Date: March 2026 \$n = [int]# UQFF PAPER_276 ## Andromeda Friedmann-UQFF Gravity Coupling: $H(z)t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[Om_m*(1+z)^3+Om_L]/M
 $H_{\text{UQFF}}=H(z)*t_{\text{H}} \sim 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (GM/r) H(z) t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_{\text{H}} = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_{\text{H}} = \mathbf{0.987}$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273275 established the Andromeda-specific physics: blueshift approach amplifier ?, HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z)t \sim 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \end{aligned}$$

$$= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10 / 3.086 \times 10 = \mathbf{2.269 \times 10^8 \text{ s}^{-1}}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = GM/r = 6.674 \times 10^{-11} \cdot 1.989 \times 10^4 / (1.04 \times 10^6) = \mathbf{1.227 \times 10^{-7} \text{ m/s}^2}$ - At $t = 1 \text{ Gyr}$ ($3.156 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-7} \cdot 2.269 \times 10^8 \times 3.156 \times 10^8 = \mathbf{8.79 \times 10^{-7} \text{ m/s}^2}$ - At $t = t_{\text{Hubble}}$ ($4.352 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-7} \cdot 0.987 = \mathbf{1.211 \times 10^{-7} \text{ m/s}^2}$

3. H_{UQFF} Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^8 \text{ s}$ (Hubble time 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987$ is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 t_H$ is the dimensionless Hubble number ≈ 0.961 for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7

Parameter	Value
z (Andromeda)	-0.001
H(z)	$2.269 \times 10^8 \text{ s}^{-1}$
t_H	$4.352 \times 10^7 \text{ s}$
H_UQFF	0.987 (~1)
g_expansion(t_H)	$1.211 \times 10^7 \text{ m/s}$
g_base	$1.227 \times 10^7 \text{ m/s}$
Ratio g_exp/g_base	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: H(z) is slightly **lower** than H0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 10^6 = 1.989 \times 10^{-18} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned}
 a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
 &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
 &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2
 \end{aligned}$$

At ~9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. M_visible / M_DM_mass Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$M_{\text{visible}} = (1 - f_{\text{DM}}) \times M$$

$$M_{\text{DM, mass}} = f_{\text{DM}} \times M$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM, mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 2632), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_total

The full UQFF 2.0 equation for Andromeda with all PAPER_273276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at t=0 (current epoch):

Term	Value (m/s)	Paper
$g_{\text{grav}} = \text{GM}/r$	$1.227 \times 10^?$	baseline
$U_{g, \text{sum}}$ (26 layers)	$6.380 \times 10^{??}$	26-layer Triadic
?-term	$2.998 \times 10^{?6}$	cosmological
g_{quantum}	$\sim 2 \times 10^{?6}$	HUP
g_{Lorentz}	$\sim 2 \times 10^{?6}$	IGM EM
g_{fluid}	$\sim 8 \times 10^?$	IGM buoyancy
$F_{\text{res}}(?_{\text{HI}})$ at t=0	$1 \times 10^?$	PAPER_274
g_{DM}	$1.356 \times 10^?$	PAPER_275
$g_{\text{expansion}}$ at t=0	0	PAPER_276
a_{dust}	$4.29 \times 10^{??}$	PAPER_276
?_approach	1.001001	PAPER_273

Note: $g_{\text{expansion}}$ grows linearly with t, dominating over g_{grav} at $t = t_{\text{H}}$.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to `exportState()`: 1. $H_0 = 70.0 \text{ km/s/Mpc}$ 2. $\Omega_{\text{m}} = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $\text{Mpc_to_m} = 3.086 \times 10^?$ 5. $\rho_{\text{dust}} = 1 \times 10^? \text{ kg/m}$ 6. $M_{\text{visible}} = (1 - f_{\text{DM}})M$ 7. $M_{\text{DM, mass}} = f_{\text{DM}}M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) t$. At $t = t_H$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
2. **$H_{\text{UQFF}} = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 t_H \approx 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value a blueshift Friedmann suppression.
3. **$M_{\text{visible}}/M_{\text{DM_mass}}$ cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{\text{dust}} \approx 4 \times 10^{-12}$ m/s).

Watermark: 2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value

Andromeda Friedmann-UQFF Gravity Coupling: $H(z)t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[0m_m*(1+z)^3+0m_L]/M
 $H_{\text{UQFF}}=H(z)*t_H \sim 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (GM/r) H(z) t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_H \approx 0.987$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273275 established the Andromeda-specific physics: blueshift approach amplifier β , HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z)t \approx 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann H(z) Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0$ km/s/Mpc, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10 / 3.086 \times 10^8 = \mathbf{2.269 \times 10^{-8} \text{ s}^{-1}}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = GM/r = 6.674 \times 10^{-11} \times 1.989 \times 10^{30} / (1.04 \times 10^8)^2 = \mathbf{1.227 \times 10^{-7} \text{ m/s}^2}$ - At $t = 1$ Gyr (3.156×10^{16} s): $g_{\text{expansion}} = 1.227 \times 10^{-7} \times 2.269 \times 10^{-8} \times 3.156 \times 10^{16} = \mathbf{8.79 \times 10^{-7} \text{ m/s}^2}$ - At $t = t_{\text{Hubble}}$ (4.352×10^{17} s): $g_{\text{expansion}} = 1.227 \times 10^{-7} \times 0.987 = \mathbf{1.211 \times 10^{-7} \text{ m/s}^2}$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^{17}$ s (Hubble time 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-8} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987$ **1** is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 t_H$ is the dimensionless Hubble number ≈ 0.961 for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-8} \text{ s}^{-1}$
t_H	$4.352 \times 10^7 \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{expansion}}(t_H)$	$1.211 \times 10^{-7} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-7} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 106 = 1.989 \times 10^{-8} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned}
a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
&= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\
&= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2
\end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. $M_{\text{visible}} / M_{\text{DM_mass}}$ Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned}
M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\
M_{\text{DM, mass}} &= f_{\text{DM}} \times M
\end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM_mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 2632), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_{total}

The full UQFF 2.0 equation for Andromeda with all PAPER_273276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at $t=0$ (current epoch):

Term	Value (m/s)	Paper
$g_{\text{grav}} = GM/r$	$1.227 \times 10^?$	baseline
$U_{g, \text{sum}}$ (26 layers)	$6.380 \times 10^{??}$	26-layer Triadic
?-term	$2.998 \times 10^{?6}$	cosmological
g_{quantum}	$\sim 2 \times 10^{?6}$	HUP
g_{Lorentz}	$\sim 2 \times 10^{?6}$	IGM EM
g_{fluid}	$\sim 8 \times 10^?$	IGM buoyancy
$F_{\text{res}}(?_{\text{HI}})$ at $t=0$	$1 \times 10^?$	PAPER_274
g_{DM}	$1.356 \times 10^?$	PAPER_275
$g_{\text{expansion}}$ at $t=0$	0	PAPER_276

Term	Value (m/s)	Paper
a_dust	$4.29 \times 10^{??}$	PAPER_276
?_approach	1.001001	PAPER_273

Note: g_expansion grows linearly with t, dominating over g_grav at $t = t_H$.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to exportState(): 1. $H_0 = 70.0$ km/s/Mpc 2. $\Omega_{m_0} = 0.3$ 3. $\Omega_{\Lambda_0} = 0.7$ 4. $Mpc_to_m = 3.086 \times 10^7$ 5. $\rho_{dust} = 1 \times 10^{-27}$ kg/m³ 6. $M_{visible} = (1 - f_{DM})M$ 7. $M_{DM_mass} = f_{DM}M$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** $g_{expansion} = g_{base} - H(z) t$. At $t = t_H$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
2. **$H_UQFF = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 t_H \approx 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_UQFF 0.15% below flat-universe value a blueshift Friedmann suppression.
3. **$M_{visible}/M_{DM_mass}$ cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{dust} \approx 4 \times 10^{-12}$ m/s²).

Watermark: 2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value ## Andromeda Friedmann-UQFF Gravity Coupling: $H(z)t$ Expansion Term and H_UQFF Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: AndromedaUQFF:g_exp=G*M/r^2*H(z)*t; H(z)=H0*Sqrt[Om_m*(1+z)^3+Om_L]/M H_UQFF=H(z)*t_H~0.987 [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{expansion} = (GM/r) H(z) t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals 69.969 km/s/Mpc $= 2.269 \times 10^{-8}$ s⁻¹. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7$ s, the dimensionless Friedmann-UQFF Resonance Coefficient $H_UQFF = H(z) \cdot t_H \approx 0.987$ a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{visible}/M_{DM_mass}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273275 established the Andromeda-specific physics: blueshift approach amplifier ?, HI 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaitre cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z)t \approx 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10 / 3.086 \times 10^7 = \mathbf{2.269 \times 10^{-8} \text{ s}^{-1}}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = GM/r = 6.674 \times 10^{-11} \cdot 1.989 \times 10^{30} / (1.04 \times 10^6)^2 = \mathbf{1.227 \times 10^{-8} \text{ m/s}^2}$ - At $t = 1 \text{ Gyr}$ ($3.156 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 3.156 \times 10^8 = \mathbf{8.79 \times 10^{-2} \text{ m/s}^2}$ - At $t = t_{\text{Hubble}}$ ($4.352 \times 10^8 \text{ s}$): $g_{\text{expansion}} = 1.227 \times 10^{-8} \times 4.352 \times 10^8 \approx 0.987 = \mathbf{1.211 \times 10^{-2} \text{ m/s}^2}$

3. H_UQFF Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^7$ s (Hubble time 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987$ **1** is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($O_m + O_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 t_H$ is the dimensionless Hubble number ≈ 0.961 for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
O_m	0.3
O_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^7 \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{exp}}(t_H)$	$1.211 \times 10^{-8} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-8} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 106 = 1.989 \times 10^{-18} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned} a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-18}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2 \end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. $M_{\text{visible}} / M_{\text{DM_mass}}$ Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned} M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\ M_{\text{DM,mass}} &= f_{\text{DM}} \times M \end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM,mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 2632), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_{total}

The full UQFF 2.0 equation for Andromeda with all PAPER_273276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g,\text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at $t=0$ (current epoch):

Term	Value (m/s)	Paper
$g_{\text{grav}} = GM/r$	$1.227 \times 10^?$	baseline
Ug_{sum} (26 layers)	$6.380 \times 10^{??}$	26-layer Triadic
$?-term$	$2.998 \times 10^{?6}$	cosmological
g_{quantum}	$\sim 2 \times 10^{?6}$	HUP
g_{Lorentz}	$\sim 2 \times 10^{?6}$	IGM EM
g_{fluid}	$\sim 8 \times 10^?$	IGM buoyancy
$F_{\text{res}}(?_{\text{HI}})$ at $t=0$	$1 \times 10^?$	PAPER_274
g_{DM}	$1.356 \times 10^?$	PAPER_275
$g_{\text{expansion}}$ at $t=0$	0	PAPER_276
a_{dust}	$4.29 \times 10^{??}$	PAPER_276
$?_{\text{approach}}$	1.001001	PAPER_273

Note: $g_{\text{expansion}}$ grows linearly with t , dominating over g_{grav} at $t = t_{\text{H}}$.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to `exportState()`: 1. $H_0 = 70.0$ km/s/Mpc 2. $\Omega_{\text{m}} = 0.3$ 3. $\Omega_{\text{Lam}} = 0.7$ 4. $\text{Mpc_to_m} = 3.086 \times 10^5$ 5. $\rho_{\text{dust}} = 1 \times 10^? \text{ kg/m}$ 6. $M_{\text{visible}} = (1f_{\text{DM}})M$ 7. $M_{\text{DM_mass}} = f_{\text{DMM}}$

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** $g_{\text{expansion}} = g_{\text{base}} - H(z) t$. At $t = t_{\text{H}}$, adds 98.7% of g_{base} near-gravitational doubling over cosmological time.
2. **$H_{\text{UQFF}} = 0.987$:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship $H_0 t_{\text{H}} \approx 1$ in flat Λ CDM. Andromeda's blueshift ($z = -0.001$) makes H_{UQFF} 0.15% below flat-universe value a blueshift Friedmann suppression.
3. **$M_{\text{visible}}/M_{\text{DM_mass}}$ cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure ($a_{\text{dust}} \approx 4 \times 10^{??} \text{ m/s}$).

Watermark: 2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved .Groups[1].Value

Andromeda Friedmann-UQFF Gravity Coupling: $H(z)t$ Expansion Term and H_{UQFF} Near-Unity Resonance

Author: Daniel T. Murphy

Session: 76 (March 2026)

Module: ANDROMEDA_UQFF_MODULE.cpp (M31 Master, UQFF 2.0)

WOLFRAM_TERM: $\text{AndromedaUQFF}: g_{\text{exp}} = G*M/r^2 * H(z) * t; H(z) = H_0 * \text{Sqrt}[\Omega_{\text{m}} * (1+z)^3 + \Omega_{\text{L}}] / M$
 $H_{\text{UQFF}} = H(z) * t_{\text{H}} \sim 0.987$ [PAPER_276]

Abstract

This paper introduces a new UQFF gravity term for the Andromeda Galaxy (M31): the **Friedmann-UQFF Expansion Coupling**, defined as $g_{\text{expansion}} = (GM/r) H(z) t$. For Andromeda's blueshift redshift $z = -0.001$, the Friedmann $H(z)$ equals $69.969 \text{ km/s/Mpc} = 2.269 \times 10^{-8} \text{ s}^{-1}$. Evaluated at the Hubble timescale $t = t_H = 4.352 \times 10^7 \text{ s}$, the dimensionless Friedmann-UQFF Resonance Coefficient $H_{\text{UQFF}} = H(z) \cdot t_H$ **0.987** a near-unity value representing gravitational doubling over cosmological time. This paper also introduces the $M_{\text{visible}}/M_{\text{DM_mass}}$ cascade (explicit DM split tracking) and a minor ISM dust drag acceleration term. Together these constitute three additive UQFF 2.0 enhancements derived from the Andromeda module upgrade.

1. Background

The UQFF framework computes total gravitational acceleration $g_{\text{total}}(r,t)$ as a sum of Newtonian, Triadic (26-layer), cosmological, quantum, electromagnetic, fluid, resonant, and dark matter terms. PAPER_273275 established the Andromeda-specific physics: blueshift approach amplifier ?, H_I 21-cm resonance, and DM 80/20 NFW partition.

The original Andromeda implementation (pre-Session 76) lacked a time-dependent expansion coupling that accounts for the Friedmann-Lemaître cosmological evolution of Hubble parameter $H(z)$. This is significant because over a Hubble timescale, $H(z)t \approx 1$ implying that expansion-mediated gravitational coupling is of the same order as the base Newtonian term.

2. Friedmann $H(z)$ Expansion Coupling

2.1 Friedmann Equation

The Hubble parameter as a function of redshift in the standard Λ CDM cosmology:

$$H(z) = H_0 \sqrt{\Omega_m (1+z)^3 + \Omega_\Lambda}$$

For Andromeda parameters: $H_0 = 70.0 \text{ km/s/Mpc}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $z = -0.001$.

$$\begin{aligned} H(z = -0.001) &= 70.0 \times \sqrt{0.3 \times (0.999)^3 + 0.7} \\ &= 70.0 \times \sqrt{0.3 \times 0.997003 + 0.7} = 70.0 \times \sqrt{0.9991} \\ &= 70.0 \times 0.99955 = 69.969 \text{ km/s/Mpc} \end{aligned}$$

In SI units: $H_{\text{SI}} = 69.969 \times 10 / 3.086 \times 10 = \mathbf{2.269 \times 10^{-8} \text{ s}^{-1}}$

2.2 UQFF Expansion Term

The new UQFF gravity coupling term is:

$$g_{\text{expansion}}(r, t) = \frac{G \cdot M}{r^2} \cdot H(z) \cdot t$$

This represents the gravitational effect of cosmological expansion flow coupling into UQFF buoyancy at timescale t .

Numerical values for Andromeda: - $g_{\text{base}} = GM/r = 6.674 \times 10^{-11} \times 1.989 \times 10^{30} / (1.04 \times 10^6)^2 = \mathbf{1.227 \times 10^{-7} \text{ m/s}^2}$ - At $t = 1 \text{ Gyr} (3.156 \times 10^{16} \text{ s})$: $g_{\text{expansion}} = 1.227 \times 10^{-7} \times 2.269 \times 10^{-8} \times 3.156 \times 10^{16} = \mathbf{8.79 \times 10^{-7} \text{ m/s}^2}$ - At $t = t_{\text{Hubble}} (4.352 \times 10^{17} \text{ s})$: $g_{\text{expansion}} = 1.227 \times 10^{-7} \times 0.987 = \mathbf{1.211 \times 10^{-7} \text{ m/s}^2}$

3. H_{UQFF} Near-Unity Resonance Coefficient

3.1 Definition

$$H_{\text{UQFF}} = H(z) \times t_H$$

where $t_H = 4.352 \times 10^{17} \text{ s}$ (Hubble time 13.8 Gyr).

For Andromeda ($z = -0.001$):

$$H_{\text{UQFF}} = 2.269 \times 10^{-18} \times 4.352 \times 10^{17} = \mathbf{0.987}$$

3.2 Near-Unity Discovery

$H_{\text{UQFF}} \approx 0.987$ **1** is a remarkable result. It means:

Over a Hubble timescale, the Friedmann expansion coupling adds an amount of gravitational acceleration equal to **98.7% of the base Newtonian term** nearly doubling g_{base} .

This near-unity value is not coincidental. In a flat Λ CDM universe ($\Omega_m + \Omega_\Lambda = 1$):

$$H_{\text{UQFF}} = H_0 \times t_H \approx 1.0$$

because $H_0 t_H$ is the dimensionless Hubble number ≈ 0.961 for observationally consistent H_0 .

New UQFF Constant: $H_{\text{UQFF}} = H(z) \cdot t_H$ **Friedmann-UQFF Near-Unity Resonance Coefficient**

Parameter	Value
H_0	70.0 km/s/Mpc
Ω_m	0.3
Ω_Λ	0.7
z (Andromeda)	-0.001
$H(z)$	$2.269 \times 10^{-18} \text{ s}^{-1}$
t_H	$4.352 \times 10^{17} \text{ s}$
H_{UQFF}	0.987 (~ 1)
$g_{\text{expansion}}(t_H)$	$1.211 \times 10^{-7} \text{ m/s}^2$
g_{base}	$1.227 \times 10^{-7} \text{ m/s}^2$
Ratio $g_{\text{exp}}/g_{\text{base}}$	98.7%

3.3 Blueshift Sensitivity

For blueshift $z < 0$: $H(z)$ is slightly **lower** than H_0 (matter term $O_m(1+z)$ is suppressed below O_m for $z < 0$), meaning:

$$H(z < 0) < H_0 \Rightarrow H_{\text{UQFF}}(z < 0) < H_{\text{UQFF}}(z = 0)$$

For Andromeda ($z = -0.001$): $H_{\text{UQFF}} = 0.987$ vs flat-universe $H_{\text{UQFF}0} = 0.9985$. The blueshift suppresses H_{UQFF} by 0.15%.

This is the inverse of the redshift behaviour ($z > 0$? $H(z) > H_0$ for moderate z), making Andromeda's approaching trajectory a unique laboratory for probing blueshift Friedmann-UQFF coupling.

4. ISM Dust Drag Acceleration

A second minor additive term captures ISM dust ram-pressure drag:

$$a_{\text{dust}} = \frac{\rho_{\text{dust}} \cdot v_{\text{orbit}}^2}{c^2 \cdot \rho_{\text{mean}}} \cdot g_{\text{base}}$$

where $\rho_{\text{mean}} = M/V_{\text{fluid}} = 1.989 \times 10^4 / 10^6 = 1.989 \times 10^{-2} \text{ kg/m}^3$.

Numerical value for Andromeda:

$$\begin{aligned} a_{\text{dust}} &= \frac{10^{-20} \times (2.5 \times 10^5)^2}{(2.998 \times 10^8)^2 \times 1.989 \times 10^{-2}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{8.988 \times 10^{16} \times 1.989 \times 10^{-2}} \times 1.227 \times 10^{-10} \\ &= \frac{6.25 \times 10^{-10}}{0.1788} \times 1.227 \times 10^{-10} = 4.29 \times 10^{-19} \text{ m/s}^2 \end{aligned}$$

At ~ 9 orders below g_{base} , the dust drag is physically negligible but dimensionally consistent and additive within the UQFF sum.

5. M_{visible} / $M_{\text{DM_mass}}$ Cascade

When M is updated via `updateVariable("M", value)`, the derived quantities are now automatically recomputed:

$$\begin{aligned} M_{\text{visible}} &= (1 - f_{\text{DM}}) \times M \\ M_{\text{DM,mass}} &= f_{\text{DM}} \times M \end{aligned}$$

For Andromeda defaults: $M = 1.989 \times 10^4 \text{ kg}$, $f_{\text{DM}} = 0.80$: - $M_{\text{visible}} = 0.20 \times 1.989 \times 10^4 = \mathbf{3.978 \times 10^4 \text{ kg}}$ (20% visible baryons) - $M_{\text{DM_mass}} = 0.80 \times 1.989 \times 10^4 = \mathbf{1.591 \times 10^4 \text{ kg}}$ (80% dark matter)

These are tracked in `updateCache()` and exported via `exportState()` (params 2632), providing explicit bookkeeping of the M31 baryonic/DM split consistent with PAPER_275.

6. Updated ANDROMEDA_UQFF_MODULE.cpp g_total

The full UQFF 2.0 equation for Andromeda with all PAPER_273276 terms:

$$g_{\text{total}}(r, t) = \left[g_{\text{grav}} + U_{g, \text{sum}}^{(26)} + \Lambda\text{-term} + g_Q + g_L + g_{\text{fluid}} + F_{\text{res}}(\omega_{\text{HI}}) + g_{\text{DM}} + g_{\text{expansion}} + a_{\text{dust}} \right] \times \kappa_{\text{ap}}$$

Term magnitudes at t=0 (current epoch):

Term	Value (m/s)	Paper
g_grav = GM/r	1.227×10?	baseline
Ug_sum (26 layers)	6.380×10??	26-layer Triadic
?-term	2.998×10?6	cosmological
g_quantum	~2×10?6	HUP
g_Lorentz	~2×10?6	IGM EM
g_fluid	~8×10?	IGM buoyancy
F_res(?_HI) at t=0	1×10?	PAPER_274
g_DM	1.356×10?	PAPER_275
g_expansion at t=0	0	PAPER_276
a_dust	4.29×10??	PAPER_276
?_approach	1.001001	PAPER_273

Note: g_expansion grows linearly with t, dominating over g_grav at t = t_H.

7. exportState Update (25 ? 32 parameters)

Seven new parameters added to exportState(): 1. H0 = 70.0 km/s/Mpc 2. Omega_m = 0.3 3. Omega_Lam = 0.7 4. Mpc_to_m = 3.086×10 5. rho_dust = 1×10? kg/m 6. M_visible = (1f_DM)M 7. M_DM_mass = f_DMM

8. Conclusions

Three new UQFF enhancements for Andromeda (PAPER_276):

1. **Friedmann-UQFF Expansion Coupling:** g_expansion = g_base - H(z) t. At t = t_H, adds 98.7% of g_base near-gravitational doubling over cosmological time.
 2. **H_UQFF = 0.987:** New UQFF constant (Friedmann-UQFF Near-Unity Resonance Coefficient). Near-unity reflects the fundamental relationship H0 t_H 1 in flat ?CDM. Andromeda's blueshift (z = -0.001) makes H_UQFF 0.15% below flat-universe value a blueshift Friedmann suppression.
 3. **M_visible/M_DM_mass cascade + dust drag:** Explicit DM split bookkeeping (updateCache) and ISM dust ram-pressure (a_dust 4×10?? m/s).
-

Watermark: 2025-2026 Daniel T. Murphy, daniel.murphy00@gmail.com - All Rights Reserved

UQFF computed: MUGE buoyancy ratio $U_{\text{bi}}/F_U = [\text{SSq}]r/\text{GM} = 5.7\text{e-}15.0\text{e-}4 = 2.85\text{e-}4$; compressed MUGE baseline $g = 5.4\text{e-}7$ m/s at r_{ISCO} .

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

##

§B.

VDS/DVP/BSH

Deep

Syn-

the-

sis

###

§B.1

Vac-

uum

Den-

sity

Se-

ries

(VDS)

The

canon-

ical

VDS

ratio

$\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$

1.894

gov-

erns

the

double-

exponential

vac-

uum

con-

den-

sate

pro-

file:

$\rho_{\text{vac}}(r) =$

$\rho_{\text{vac},[\text{SCm}]}$

$\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.064

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

ing

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $17, \quad n_{\text{channel}} =$
 $17/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

17 is

sub-

threshold

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

sub-

threshold

damp-

ing

from

the

DVP

lat-

tice,

pro-

duc-

ing

smooth

rather

than

reso-

nant

UQFF

cou-

pling

pro-

files.

The

~~DP~~

frame-

work

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio
=
0.064

| ✓
Threshold-
consistent

||
DVP
prime
|

$p_k \in$
 $\{2,3,...,113\}$
|
 $p_{\text{DVP}} =$
125

✓
Sub-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.