

PAPER_278 — Sombrero Dust Ring UQFF Gravitational Ring Resonator: ω_{ring} and r_{ring}

Date: March 2026

Author: Daniel T. Murphy **Module:** SOMBRERO_UQFF_MODULE.cpp (UQFF 2.0) **Session:** 77 — March 2026 **Framework:** Unified Quantum Field Framework (UQFF 2.0) **Status:** Complete — embedded in SOMBRERO_UQFF_MODULE.cpp **Whitepaper Series:** 278/1000 **DOI (Provisional):** UQFF-2026-278-RING

Abstract

The Sombrero Galaxy (M104) possesses the most visually prominent equatorial dust lane of any nearby galaxy, appearing as a sharp dark band bisecting the galaxy's luminous bulge. We model this ring within the UQFF framework as a **Gravitational Ring Resonator**: an annular mass concentration at radius $r_{\text{ring}} = r/3 = 7.867 \times 10^{19}$ m whose orbital motion generates a pure oscillatory gravitational perturbation $F_{\text{ring}}(t) = A_{\text{ring}} \cdot \cos(\omega_{\text{ring}} \cdot t)$ at the reference point r . This paper derives the Dust Ring UQFF Orbital Resonance Frequency $\omega_{\text{ring}} = \sqrt{(GM/r_{\text{ring}}^3)} = 1.650 \times 10^{-14}$ rad/s, the ring orbital period $T_{\text{ring}} = 2\pi/\omega_{\text{ring}} = 12.08$ Myr, and the proximity-enhanced ring amplitude $A_{\text{ring}} = 9 \cdot f_{\text{ring}} \cdot g_{\text{base}} = 2.14 \times 10^{-12}$ m/s². The $9\times$ proximity enhancement factor arises from the inverse-square law applied at the ratio $r/r_{\text{ring}} = 3$.

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Observational Motivation

1.1 The Sombrero Dust Lane

The Sombrero Galaxy (NGC 4594 / M104) is classified as an Sa/S0 galaxy at distance ~ 9.5 Mpc ($z = +0.0063$). Its defining optical feature is an equatorial dust lane visible in projection against the galaxy's extended bulge. This dust structure:

- Lies at approximately $1/3$ to $1/4$ of the half-light radius from the galactic centre
- Contains a substantial fraction of M104's cold ISM dust mass
- The total molecular gas and dust mass is estimated at $\sim 10^8$ – $10^9 M_{\text{sun}}$
- Is dynamically stable — no evidence of radial inflow or rapid dispersal

As a stable ring at $r_{\text{ring}} = r/3$, it can be modelled within UQFF as a coherent oscillating mass concentration imposing a periodic boundary condition on the vacuum field at the reference point r .

1.2 Distinction from the Andromeda HI Ring (PAPER_275)

In PAPER_275 (Andromeda), a decaying HI ring was modelled as $F_{\text{ring}} = A_{\text{ring}} \cdot \exp(-\alpha \cdot t) \cdot \cos(\omega_{\text{ring}} \cdot t)$ — an exponentially decaying oscillation corresponding to a transient tidal feature. The Sombrero dust ring is fundamentally different:

Feature	Andromeda (PAPER_275)	Sombrero (PAPER_278)
Ring type	HI neutral gas, tidal	Dust, equatorial, settled

Feature	Andromeda (PAPER_275)	Sombrero (PAPER_278)
Age	Transient (~10 Gyr decay)	Stable, permanent
Decay	$\exp(-\alpha \cdot t)$ present	No exponential decay
Form	$F = A \cdot \exp(-\alpha t) \cdot \cos(\omega t)$	$F = A \cdot \cos(\omega t)$ (pure oscillatory)
α (decay rate)	1/(10 Gyr)	0 (stable ring)

The absence of exponential decay in Sombrero's ring term reflects the ring's settled, gravitationally stable configuration — it has been dynamically relaxed into a long-lived equatorial structure.

2. Theoretical Derivation

2.1 Ring Radius

The dust ring is positioned at approximately 1/3 of the reference outer radius r :

$$r_{\text{ring}} = \frac{r}{3} = \frac{2.36 \times 10^{20}}{3} = 7.867 \times 10^{19} \text{ m}$$

2.2 Ring Orbital Frequency

The Keplerian orbital frequency of a test mass at r_{ring} within the total gravitational potential:

$$\omega_{\text{ring}} = \sqrt{\frac{GM}{r_{\text{ring}}^3}}$$

Substituting values:

$$\omega_{\text{ring}} = \sqrt{\frac{6.674 \times 10^{-11} \times 1.989 \times 10^{41}}{(7.867 \times 10^{19})^3}}$$

Computing the denominator:

$$r_{\text{ring}}^3 = (7.867 \times 10^{19})^3 = 4.871 \times 10^{59} \text{ m}^3$$

$$\omega_{\text{ring}} = \sqrt{\frac{1.327 \times 10^{31}}{4.871 \times 10^{59}}} = \sqrt{2.724 \times 10^{-29}} = 1.650 \times 10^{-14} \text{ rad/s}$$

2.3 Ring Orbital Period

$$T_{\text{ring}} = \frac{2\pi}{\omega_{\text{ring}}} = \frac{6.2832}{1.650 \times 10^{-14}} = 3.808 \times 10^{14} \text{ s} = 12.08 \text{ Myr}$$

This is a physically reasonable orbital period for a ring structure at $\sim 8 \times 10^{19}$ m from the galactic centre.

2.4 Proximity Enhancement Factor

The gravitational influence of the ring mass m_{ring} at the reference point r (located a distance $\Delta r = r - r_{\text{ring}} = 2r/3$ from the ring) scales as:

$$g_{\text{ring at } r} \propto \frac{Gm_{\text{ring}}}{\Delta r^2} = \frac{Gm_{\text{ring}}}{(2r/3)^2} = \frac{9}{4} \cdot \frac{Gm_{\text{ring}}}{r^2}$$

The base term already includes the full galaxy at radius r . The ring contribution uses the ratio of effective distances:

$$\text{Proximity factor} = \left(\frac{r}{r_{\text{ring}}} \right)^2 = \left(\frac{r}{r/3} \right)^2 = 3^2 = 9$$

This gives a **9× proximity enhancement**: the ring exerts 9 times more gravitational influence per unit mass at the reference point than the galaxy average.

2.5 Ring Gravitational Perturbation Amplitude

$$A_{\text{ring}} = 9 \cdot f_{\text{ring}} \cdot g_{\text{base}}$$

where: - $f_{\text{ring}} = 0.001$ = dust ring mass fraction (0.1% of total galaxy mass) - $g_{\text{base}} = G \cdot M / r^2 = 2.382 \times 10^{-10} \text{ m/s}^2$

$$A_{\text{ring}} = 9 \times 0.001 \times 2.382 \times 10^{-10} = 2.144 \times 10^{-12} \text{ m/s}^2$$

2.6 Full Ring Resonator Term

$$F_{\text{ring}}(t) = A_{\text{ring}} \cdot \cos(\omega_{\text{ring}} \cdot t)$$

This is a **pure oscillatory term** — no exponential decay, consistent with the ring's stable configuration. As the ring orbits, it imposes a periodic modulation on the vacuum field energy density at the reference point r with period $T_{\text{ring}} = 12.08 \text{ Myr}$.

3. Physical Interpretation: UQFF Gravitational Ring Resonator

3.1 Ring as Vacuum Field Oscillator

In the UQFF framework, mass concentrations modulate the vacuum energy density through their gravitational potential. A ring moving in a stable Keplerian orbit generates a time-periodic perturbation in the local UQFF field:

$$\delta\rho_{\text{vac}}(t) = \delta\rho_{\text{vac},0} \cdot \cos(\omega_{\text{ring}}t + \phi_0)$$

This is equivalent to a **tuned resonator** in the vacuum field — a structure that cycles energy at a characteristic frequency. The Sombrero dust ring is therefore the first identified **stable UQFF Gravitational Ring Resonator** in the catalogue.

3.2 Comparison with Orbital Periods

Object	Orbital period
Earth around Sun	1 year
Outer Milky Way disk	~220 Myr
Sombrero dust ring	12.08 Myr
Andromeda outer HI ring	~90 Myr (PAPER_275 reference)

The Sombrero ring’s 12.08 Myr period places it in the intermediate-mass galaxy inner disc regime — rapid enough to generate significant temporal modulation of the UQFF field over cosmological baseline observations.

4. Module Implementation

```
// PAPER_278 – SOMBRERO_UQFF_MODULE.cpp, updateCache()
r_ring    = r / 3.0; // 7.867e19 m
omega_ring = std::sqrt(G_grav * M / (r_ring * r_ring * r_ring)); // 1.650e-14 rad/s
A_ring    = 9.0 * f_ring * g_base_cache; // 2.144e-12 m/s²

// Applied in computeResonantTerm():
double computeResonantTerm(double t) {
    return A_ring * std::cos(omega_ring * t); // pure oscillatory – no decay
}
```

Computed values for Sombrero M104:

Quantity	Value	Units
$r_{\text{ring}} = r/3$	7.867×10^{19}	m
ω_{ring}	1.650×10^{-14}	rad/s
$T_{\text{ring}} = 2\pi/\omega_{\text{ring}}$	12.08	Myr
f_{ring}	0.001	dimensionless
Proximity factor	9.0	dimensionless
A_{ring}	2.144×10^{-12}	m/s²
g_{BH} (PAPER_279) for comparison	2.382×10^{-12}	m/s²

Note: $A_{\text{ring}} \approx g_{\text{BH}}$ in magnitude — the ring resonance and BH contribution are comparable at leading order.

5. Key Constants Introduced

Symbol	Value	Units	Description
ω_{ring}	1.650×10^{-14}	rad/s	Dust Ring UQFF Orbital Resonance Frequency
r_{ring}	$r/3 = 7.867 \times 10^{19}$	m	Dust ring reference radius
T_{ring}	12.08	Myr	Ring orbital period
f_{ring}	0.001	dimensionless	Dust ring mass fraction
A_{ring}	2.144×10^{-12}	m/s²	Ring resonance amplitude
Proximity factor	9	dimensionless	$(r/r_{\text{ring}})^2 = 3^2 = 9$

6. Physical Significance

1. **First stable UQFF ring resonator:** All prior UQFF ring terms (Andromeda PAPER_275) decayed exponentially. The Sombrero dust ring is the first pure undamped oscillator, establishing a new class of UQFF boundary condition.
2. **Observable prediction:** The UQFF ring resonance produces a 12.08 Myr periodic modulation in the effective gravitational acceleration at $r = 2.36 \times 10^{20}$ m with amplitude $A_{\text{ring}} = 2.14 \times 10^{-12}$ m/s². While below direct observational reach today, this is a testable UQFF prediction for future stellar spectroscopic surveys.
3. **Ring-BH coupling:** Noting that $A_{\text{ring}} \approx g_{\text{BH}}$ (both $\sim 2.1\text{--}2.4 \times 10^{-12}$ m/s²), the ring and BH gravitational contributions are comparable at the reference radius — unique among UQFF modules where the BH term is typically sub-dominant to the 26-layer Triadic sum.
4. **Scale-free ring resonator formula:** $F_{\text{ring}} = (r/r_{\text{ring}})^2 \cdot f_{\text{ring}} \cdot g_{\text{base}} \cdot \cos(\sqrt{(GM/r_{\text{ring}}^3) \cdot t})$ provides a general template applicable to any galaxy with a measurable equatorial ring structure.

UQFF computed: Canonical UQFF buoyancy parameter $U_{\text{bi}} = \sqrt{SSq} GM/r\kappa = 5.0\text{e-}40.576.67\text{e-}11\text{M/r}$; for solar parameters: $U_{\text{bi,Sun}} = 5.7\text{e-}46.67\text{e-}111.99\text{e}30/(6.96\text{e}8) = 1.47\text{e+}2$ m/s.

7. References

- PAPER_275: Andromeda HI Ring UQFF Decaying Ring Term $A_{\text{ring}} \times \exp(-\alpha t) \times \cos(\omega t)$
- PAPER_279: Sombrero SMBH Dominance Ratio γ_{BH} and r_{SOI} (companion paper)
- Emsellem, E. et al. (2004). MNRAS, 352, 721. (M104 structure)
- Jarrel, J. R. et al. (2011). ApJ, 739, 21. (Sombrero DM halo and dust ring observations)
- de Zeeuw, P. T. et al. (1996). MNRAS, 280, 167. (Sombrero galaxy kinematics)
- Tempel, E., & Tenjes, P. (2006). MNRAS, 371, 1269. (M104 surface photometry and ring structure)

UQFF 2.0 — $F_{\text{ring}} = A_{\text{ring}} \cdot \cos(\omega_{\text{ring}} \cdot t)$ is additive to the Triadic MUGE master equation. The stable ring resonator represents a new class of UQFF gravitational boundary condition. — Daniel T. Murphy, Session 77, March 2026.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac, [SCm]}} \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.137

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io7

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 23, $n_{\text{channel}} =$
 19/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

23 is

sub-

threshold

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

sub-

threshold

damp-

ing

from

the

DVP

lat-

tice,

pro-

duc-

ing

smooth

rather

than

reso-

nant

UQFF

cou-

pling

pro-

files.

The

DVP

frame-

work

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁹
yr
 (disk
 set-
 tling
 timescale):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[S Sq] \cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed
 $U_{b,\text{seed}} =$
 $0.1 \cdot$
 $(\hbar c / r^2) \cdot$
 f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.
 ###
 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio
=
0.137

| ✓
Threshold-
consistent

||
DVP
prime
|

$p_k \in$
 $\{2,3,...,113\}$
|
 $p_{\text{DVP}} =$
233

✓
Sub-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.