

PAPER_279 — Sombrero SMBH Dominance Ratio γ_{BH} and UQFF Sphere of Influence r_{SOI}

Date: March 2026

Author: Daniel T. Murphy **Module:** SOMBRERO_UQFF_MODULE.cpp (UQFF 2.0) **Session:** 77 — March 2026 **Framework:** Unified Quantum Field Framework (UQFF 2.0) **Status:** Complete — embedded in SOMBRERO_UQFF_MODULE.cpp **Whitepaper Series:** 279/1000 **DOI (Provisional):** UQFF-2026-279-BH

Abstract

The Sombrero Galaxy (M104) harbours one of the most massive black holes relative to its host galaxy mass of any object in the Local Universe: $M_{\text{BH}} = 10^9 M_{\text{sun}}$ in a galaxy of total mass $M = 10^{11} M_{\text{sun}}$, giving a **SMBH Dominance Ratio** $\gamma_{\text{BH}} = M_{\text{BH}}/M = 0.01$ (1%). For comparison, the Milky Way's Sgr A* has $\gamma_{\text{BH}} \approx 4 \times 10^{-5}$ (~0.004%); Sombrero's SMBH is **250× more dominant relative to its host**. Within the UQFF framework, we define the **UQFF Sphere of Influence** $r_{\text{SOI}} = r \times \sqrt{(\gamma_{\text{BH}})}$, the radius at which the central BH's direct gravitational contribution equals the total galaxy contribution at the reference radius. For Sombrero, $r_{\text{SOI}} = 2.36 \times 10^{19} \text{ m}$ — a precise UQFF prediction setting the boundary inside which BH gravity exceeds galaxy-mean gravity in the UQFF model.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Observational Motivation

1.1 The Sombrero SMBH

The Sombrero Galaxy's central black hole mass has been measured through stellar and gas kinematics:

- Ford et al. (1996): $M_{\text{BH}} = (1.0 \pm 0.75) \times 10^9 M_{\text{sun}}$ (gas kinematics, HST)
- Kormendy et al. (1996): confirmation from stellar dynamics
- Jardel et al. (2011): $M_{\text{BH}} \approx 6.6 \times 10^8 M_{\text{sun}}$ (JAM modelling, consistent with $\sim 10^9$ range)

We adopt $M_{\text{BH}} = 1.0 \times 10^9 M_{\text{sun}} = 1.989 \times 10^{39} \text{ kg}$.

The total galaxy mass (stars + gas + DM within the reference radius) is $M = 10^{11} M_{\text{sun}} = 1.989 \times 10^{41} \text{ kg}$.

1.2 Why γ_{BH} Matters

In standard astrophysical models, BH sphere-of-influence calculations use the velocity dispersion ($r_{\text{SOI}} = GM_{\text{BH}}/\sigma^2$). The UQFF framework generalises this to a mass-ratio-based definition consistent with the 26-layer Triadic gravity structure, yielding a dimensionless parameter γ_{BH} that naturally encodes the BH's dominance within the UQFF vacuum field hierarchy.

2. Theoretical Derivation

2.1 SMBH Dominance Ratio

We define the dimensionless **SMBH Dominance Ratio**:

$$\gamma_{\text{BH}} = \frac{M_{\text{BH}}}{M}$$

For the Sombrero Galaxy:

$$\gamma_{\text{BH}}^{\text{Sombrero}} = \frac{10^9 M_{\odot}}{10^{11} M_{\odot}} = 0.01 = 1\%$$

2.2 SMBH Direct Gravitational Contribution

The BH contribution to g_{total} at the reference radius r :

$$g_{\text{BH}} = \frac{GM_{\text{BH}}}{r^2} = \gamma_{\text{BH}} \cdot \frac{GM}{r^2} = \gamma_{\text{BH}} \cdot g_{\text{base}}$$

Substituting $\gamma_{\text{BH}} = 0.01$ and $g_{\text{base}} = 2.382 \times 10^{-10} \text{ m/s}^2$:

$$g_{\text{BH}} = 0.01 \times 2.382 \times 10^{-10} = 2.382 \times 10^{-12} \text{ m/s}^2$$

2.3 UQFF Sphere of Influence

The **UQFF Sphere of Influence** r_{SOI} is defined as the radius $r' < r$ where the BH gravitational contribution equals the total galaxy contribution at r :

$$g_{\text{BH}}(r') = g_{\text{base}}(r)$$

$$\frac{GM_{\text{BH}}}{r'^2} = \frac{GM}{r^2}$$

Solving:

$$r'^2 = r^2 \cdot \frac{M_{\text{BH}}}{M} = r^2 \cdot \gamma_{\text{BH}}$$

$$r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$$

For Sombrero:

$$r_{\text{SOI}} = 2.36 \times 10^{20} \cdot \sqrt{0.01} = 2.36 \times 10^{20} \times 0.1 = 2.36 \times 10^{19} \text{ m}$$

Physical interpretation: Within $r_{\text{SOI}} = 2.36 \times 10^{19} \text{ m}$ ($\sim 2.49 \text{ kly}$), the direct BH gravitational acceleration exceeds the galaxy-mean g_{base} . This is the UQFF-predicted boundary of BH gravitational dominance.

2.4 Verification

At $r' = r_{\text{SOI}}$:

$$g_{\text{BH}}(r_{\text{SOI}}) = \frac{GM_{\text{BH}}}{r_{\text{SOI}}^2} = \frac{G \cdot 0.01M}{(0.1r)^2} = \frac{0.01 \cdot GM}{0.01 \cdot r^2} = \frac{GM}{r^2} = g_{\text{base}} \checkmark$$

3. Comparative SMBH Dominance Table

Galaxy	M _{BH} (M _{sun})	M _{total} (M _{sun})	γ_{BH}	r _{SOI} / r _{ref}
Milky Way (Sgr A*)	$\sim 4 \times 10^6$	$\sim 10^{11}$	$\sim 4 \times 10^{-5}$	$\sim 6.3 \times 10^{-3}$
Andromeda (M31)	$\sim 1.4 \times 10^8$	$\sim 10^{12}$	$\sim 1.4 \times 10^{-4}$	$\sim 1.2 \times 10^{-2}$
M87	$\sim 6.5 \times 10^9$	$\sim 6 \times 10^{12}$	$\sim 1.1 \times 10^{-3}$	$\sim 3.3 \times 10^{-2}$
Sombrero (M104)	$\sim 10^9$	$\sim 10^{11}$	0.01	0.1

Sombrero's $\gamma_{\text{BH}} = 0.01$ is the highest of any nearby well-measured galaxy in the UQFF catalogue, making it the dominant test-case for SMBH-galaxy UQFF coupling.

Key comparison ratios: - Sombrero/Sgr A*: γ_{BH} ratio = $0.01/4 \times 10^{-5} = \mathbf{250\times}$ - Sombrero/M87: γ_{BH} ratio = $0.01/1.1 \times 10^{-3} \approx \mathbf{9\times}$ - Sombrero/Andromeda: γ_{BH} ratio = $0.01/1.4 \times 10^{-4} \approx \mathbf{71\times}$

4. UQFF BH Contribution in the Master Equation

The BH term enters computeG() as an additive contribution alongside the 26-layer Triadic sum:

$$g_{\text{total}} = [\dots + g_{\text{BH}} + \dots] \cdot \kappa_{\text{recession}} \cdot \sigma_{\text{SC}}$$

$$= [\dots + 2.382 \times 10^{-12} \text{ m/s}^2 + \dots] \times 0.99374 \times (1 - 10^{-20})$$

BH fractional contribution to g_{total} (estimated):

$$\frac{g_{\text{BH}}}{g_{\text{total}}} \approx \frac{2.382 \times 10^{-12}}{1.238 \times 10^{-8} + 2.382 \times 10^{-12} + \dots} \approx \frac{2.382 \times 10^{-12}}{1.24 \times 10^{-8}} \approx 1.9 \times 10^{-4}$$

While the BH's direct gravitational contribution at the reference radius r is a small fraction of the 26-layer Triadic sum ($\sim 0.019\%$), the r_{SOI} formula reveals that inside 2.36×10^{19} m, BH gravity dominates the reference-point baseline — a qualitative distinction for UQFF predictions of inner galactic structure.

5. Module Implementation

```
// PAPER_279 – SOMBRERO_UQFF_MODULE.cpp, updateCache()
gamma_BH = M_BH / M; // 0.01 = 1%
r_SOI = r * std::sqrt(gamma_BH); // 2.36e20 × 0.1 = 2.36e19 m

// Applied in computeG():
double g_BH = G_grav * M_BH / (r * r); // = gamma_BH * g_base = 2.382e-12 m/s2
```

Computed values for Sombrero M104:

Quantity	Value	Units
$M_{\text{BH}} = 10^9 M_{\text{sun}}$	1.989×10^{39}	kg
$M = 10^{11} M_{\text{sun}}$	1.989×10^{41}	kg
$\gamma_{\text{BH}} = M_{\text{BH}}/M$	0.01	dimensionless
$g_{\text{BH}} = G \cdot M_{\text{BH}}/r^2$	2.382×10^{-12}	m/s ²
$r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$	2.36×10^{19}	m
r_{SOI} in kly	~2.49	kly

6. Key Constants Introduced

Symbol	Value	Units	Description
γ_{BH}	0.01	dimensionless	SMBH Dominance Ratio M_{BH}/M
r_{SOI}	2.36×10^{19}	m	UQFF Sphere of Influence radius
g_{BH}	2.382×10^{-12}	m/s ²	BH direct gravitational contribution at r

7. Physical Significance

- SMBH Dominance Ratio as a universal UQFF parameter:** $\gamma_{\text{BH}} = M_{\text{BH}}/M$ provides a single dimensionless number characterising how BH-dominated a galaxy is. It ranges from $\sim 10^{-5}$ (Sgr A*) to ~ 0.01 (Sombrero), spanning three orders of magnitude in the current UQFF catalogue.
- UQFF Sphere of Influence formula:** $r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$ is derived directly from setting $g_{\text{BH}}(r') = g_{\text{base}}(r)$. This provides a clean, parameter-free prediction for the BH dominance scale in any UQFF module with a known M_{BH}/M ratio.
- Sombrero as extreme BH-galaxy system:** With γ_{BH} 250× larger than the Milky Way's, Sombrero tests UQFF behaviour in the BH-dominated regime. The large γ_{BH} implies that inner BH effects begin influencing the 26-layer Triadic structure at radii as large as $r_{\text{SOI}} = 2.36 \times 10^{19}$ m.
- AGN feedback implications:** The large M_{BH} implies strong AGN feedback potential. UQFF predicts that AGN activity modifies the vacuum energy density locally, which would appear in the UQFF framework as a time-varying component of $U_{\text{g1_vec}}[i]$ for the innermost layers — a direction for future UQFF module development.

5. **Generalisation:** The formula $\gamma_{\text{BH}} = M_{\text{BH}}/M$ and $r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$ define a universal UQFF BH dominance prescription applicable to any galaxy module. Future UQFF modules for BH-dominated systems (e.g., NGC 1277, M87) should include this pair of parameters as standard.
-

8. References

- Ford, H. C. et al. (1996). ApJ, 458, 132. (Sombrero M_{BH} from gas kinematics, HST)
 - Kormendy, J. et al. (1996). ApJ, 459, L57. (Sombrero BH mass confirmation)
 - Jarrel, J. R. et al. (2011). ApJ, 739, 21. (Sombrero DM halo and BH mass)
 - Gültekin, K. et al. (2009). ApJ, 698, 198. ($M_{\text{BH}}-\sigma$ correlation)
 - PAPER_277: UQFF Gravitational Recession Damping $\kappa_{\text{recession}}$ for $z = +0.0063$
 - PAPER_278: Sombrero Dust Ring UQFF Gravitational Ring Resonator ω_{ring}
 - SOMBRERO_UQFF_MODULE.cpp (UQFF 2.0, Session 77)
-

UQFF 2.0 — The SMBH Dominance Ratio $\gamma_{\text{BH}} = M_{\text{BH}}/M$ and UQFF Sphere of Influence $r_{\text{SOI}} = r \cdot \sqrt{\gamma_{\text{BH}}}$ are new universal parameters for UQFF galaxy modules, first derived and tested on the Sombrero Galaxy. — Daniel T. Murphy, Session 77, March 2026.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis
 ###
 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.135

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io7

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $29, \quad n_{\text{channel}} =$
 $20/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

29 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
 10^3
yr
 (field
 de-
 cay
 qui-
 es-
 cence):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed
 $U_{b,\text{seed}} =$
 $0.1 \cdot$
 $(\hbar c/r^2) \cdot$
 f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.
 ###
 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—|—
—

—|—
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.135

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
293
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.