

PAPER_554: Buoyancy-Stratified Factorial Geometry — Riemann Curvature of the Aether Metric

Author: Daniel T. Murphy — Star Magic / UQFF Framework

Session: 148 | **Source:** Composed from CP4 #43, #66, #67 (Sessions 107-110)

CP4 Class: BSFGriemannCurvatureAetherMetricCalculator (#149)

Date: 2026-03-27

Context note: The Aether metric $A_{\mu\nu}$ was introduced in PAPER_392 (CP4 #43) with coupling $\eta = 10^{-22}$ and scalar amplitude T_{s00} . PAPER_416 (CP4 #66) provided the five-component spatial decomposition of $T_{s00}(r)$. This paper combines those results to derive for the first time the full Riemann curvature tensor of the BSFG geometry — the first paper in the BSFG Complete Geometric System series (PAPER_554-558).

Abstract

This paper presents a UQFF analysis of Stratified Factorial Geometry — Riemann Curvature of the Aether Metric, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

§1 Abstract

The Buoyancy-Stratified Factorial Geometry (BSFG) is defined on a 26-dimensional pseudo-Riemannian manifold $\mathcal{M}^{26}$ equipped with the Aether-perturbed metric $A_{\mu\nu}(r) = g_{\mu\nu} + \varepsilon(r)\delta_{\mu\nu}$, where $\varepsilon(r) = \eta T_{s00}(r) \cos(\pi t_n)$ and $T_{s00}(r)$ is the five-component stellar stress-energy density from PAPER_416. This paper derives the Christoffel symbols, Riemann curvature tensor, Ricci tensor, and Ricci scalar for the 4D slice of BSFG at a field point r . The key result is:

$$R^r_{0r0} \approx \frac{\varepsilon''}{2} = \frac{6\eta \cos(\pi t_n) C_{\text{num}}}{r^5}$$

with $C_{\text{num}} = (M_s c^2 + L_s/c^2)/(4\pi/3)$. At the solar surface $r = R_\odot$: $R^r_{0r0} \approx 1.56 \times 10^{-19} \text{ m}^{-2}$, which is $\approx 4 \times 10^{-26}$ times the Schwarzschild curvature — confirming the Aether contribution is a genuine but ultra-weak geometric perturbation.

§2 The Aether Metric — Previous Results

From PAPER_392 (CP4 #43), the BSFG metric is:

$$A_{\mu\nu} = g_{\mu\nu} + \eta T_{s00} \cos(\pi t_n) \delta_{\mu\nu}$$

In components (4D Minkowski background with signature $+---$):

$$A_{\mu\nu} = \text{diag}(1 + \varepsilon, -1 + \varepsilon, -1 + \varepsilon, -1 + \varepsilon)$$

where $\varepsilon = \eta T_{s00}(r) \cos(\pi t_n)$. From PAPER_416 (CP4 #66), the five-component T_{s00} with dominant radial term:

$$T_{s00}(r) = \underbrace{\frac{M_s c^2}{\frac{4}{3}\pi r^3}}_{T_1 \propto r^{-3}} + \underbrace{\frac{L_s}{c^2 \cdot \frac{4}{3}\pi r^3}}_{T_2 \propto r^{-3}} + \underbrace{\frac{\rho_{SCm} v_{SCm}^2}{c^2} + \frac{\rho_A v_{UA}^2}{c^2} + \rho_{sw} v_{sw}^2}_{\text{constant terms}}$$

The radially-dependent numerator is $C_{\text{num}} = (M_s c^2 + L_s/c^2)/(4\pi/3) \approx 4.27 \times 10^{46} \text{ J} \cdot \text{m}$.

§3 Christoffel Symbols

Step 1. Compute $\varepsilon(r)$ and its radial derivatives:

$$\varepsilon'(r) = \frac{d\varepsilon}{dr} = -\frac{3\eta \cos(\pi t_n) C_{\text{num}}}{r^4}, \quad \varepsilon''(r) = +\frac{12\eta \cos(\pi t_n) C_{\text{num}}}{r^5}$$

Step 2. For a diagonal metric $A_{\mu\mu}(r)$ depending only on $r = x^1$, the non-zero Christoffel symbols of the Levi-Civita connection are:

$$\Gamma_{\mu\mu}^r = -\frac{\partial_r A_{\mu\mu}}{2 A_{rr}} = -\frac{\varepsilon'}{2(-1 + \varepsilon)} \quad (\text{no sum on } \mu)$$

$$\Gamma_{\alpha r}^\alpha = \Gamma_{r\alpha}^\alpha = \frac{\partial_r A_{\alpha\alpha}}{2 A_{\alpha\alpha}} \quad (\text{no sum on } \alpha)$$

Explicitly at leading order in ε :

Symbol	Exact	Leading order
Γ_{00}^r	$-\varepsilon'/(2(-1 + \varepsilon))$	$+\varepsilon'/2$
Γ_{rr}^r	$+\varepsilon'/(2(-1 + \varepsilon))$	$-\varepsilon'/2$
Γ_{0r}^0	$+\varepsilon'/(2(1 + \varepsilon))$	$+\varepsilon'/2$
Γ_{ir}^i	$+\varepsilon'/(2(-1 + \varepsilon))$	$-\varepsilon'/2$
Γ_{ii}^r	$-\varepsilon'/(2(-1 + \varepsilon))$	$+\varepsilon'/2$

§4 Riemann Curvature Tensor

Step 3. Apply the Riemann tensor formula:

$$R^\rho_{\sigma\mu\nu} = \partial_\mu \Gamma_{\nu\sigma}^\rho - \partial_\nu \Gamma_{\mu\sigma}^\rho + \Gamma_{\mu\lambda}^\rho \Gamma_{\nu\sigma}^\lambda - \Gamma_{\nu\lambda}^\rho \Gamma_{\mu\sigma}^\lambda$$

The dominant component (tidal force in the radial-temporal plane):

$$R^r_{0r0} = \partial_r \Gamma_{00}^r + \Gamma_{rr}^r \Gamma_{00}^r - \Gamma_{00}^r \Gamma_{0r}^0$$

$$= \frac{\varepsilon''}{2} - \frac{(\varepsilon')^2}{2} + O(\varepsilon^3) \approx \frac{\varepsilon''}{2}$$

Step 4. Substituting $\varepsilon'' = 12\eta \cos(\pi t_n) C_{\text{num}}/r^5$:

$$R^r_{0r0} = \frac{6\eta \cos(\pi t_n) C_{\text{num}}}{r^5}$$

The higher-order $(\varepsilon')^2$ correction is of order $\eta^2 T_{s00}^2 \sim 10^{-30}$, negligible.

§5 Ricci Tensor and Ricci Scalar

Ricci tensor — applying $R_{\mu\nu} = R^\rho_{\mu\rho\nu}$ with $SO(3)$ spherical symmetry ($x^2 = y$, $x^3 = z$ contribute equally to $x^1 = r$):

$$R_{00} = R^r_{0r0} + R^\theta_{0\theta0} + R^\phi_{0\phi0} = 3 R^r_{0r0}$$

$$R_{rr} = -R^r_{0r0} + 2\left(\frac{\varepsilon''}{2} - \frac{(\varepsilon')^2}{4}\right)$$

Ricci scalar:

$$R = A^{\mu\nu} R_{\mu\nu} = \frac{R_{00}}{A_{00}} + \frac{R_{rr}}{A_{rr}} + \frac{2R_{\theta\theta}}{A_{\theta\theta}}$$

Kretschner scalar (leading order):

$$K = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \approx 12 (R^r_{0r0})^2$$

§6 Numerical Values at the Solar Surface

At $r = R_\odot = 6.96 \times 10^8$ m, $t_n = 0$ (maximum coupling), $\eta = 10^{-22}$:

Quantity	Value	Notes
$\varepsilon(R_\odot)$	1.27×10^{-2}	Not small — linearization is qualitative at surface
$\varepsilon'(R_\odot)$	$-5.47 \times 10^{-11} \text{ m}^{-1}$	Aether gradient
$\varepsilon''(R_\odot)$	$+3.13 \times 10^{-19} \text{ m}^{-2}$	Curvature driver
R^r_{0r0}	$+1.56 \times 10^{-19} \text{ m}^{-2}$	BSFG tidal curvature
R_{scalar}	$\approx +3.0 \times 10^{-19} \text{ m}^{-2}$	de Sitter (gravity-dominant)
K	$\approx 2.9 \times 10^{-37} \text{ m}^{-4}$	Kretschner scalar

For comparison, the Schwarzschild curvature at $R_\odot$: $R^r_{0r0}|_{\text{GR}} \approx GM_\odot/R_\odot^3 \approx 3.95 \times 10^{-7} \text{ m}^{-2}$, giving:

$$\frac{R^r_{0r0}|_{\text{BSFG}}}{R^r_{0r0}|_{\text{GR}}} \approx \frac{1.56 \times 10^{-19}}{3.95 \times 10^{-7}} \approx 3.9 \times 10^{-13}$$

The BSFG Aether curvature is $\sim 10^{12}$ times weaker than GR at the solar surface, consistent with the perturbative assumption $\eta \sim 10^{-22}$ being a tiny coupling.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis
 ###
 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.184

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io6

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 47, $n_{\text{channel}} =$
 9/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

47 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
**Hub-
 ble
 time**
 (super-
 Hubble
 satu-
 ra-
 tion):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos(\frac{2\pi j}{26})$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed
 $U_{b,\text{seed}} =$
 $0.1 \cdot$
 $(\hbar c / r^2) \cdot$
 f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.
 ###
 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—-|-

—|-
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.184

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
472
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
GR Schwarzschild metric recovery	BSFG line element $\rightarrow g_{tt} = -(1-2GM/rc^2) \equiv \text{GR}$ in $\varepsilon_{\text{BSFG}} \rightarrow 0$ limit	Schwarzschild metric (GR exact)	PDG 2024 / MTW	✓ BSFG reduces to GR
Shapiro time delay	BSFG geodesic $\rightarrow \Delta t_{\text{BSFG}} \approx \Delta t_{\text{GR}} \times (1 + \varepsilon_{\text{correction}})$	Cassini: $\Delta t / \Delta t_{\text{GR}} = 1 \pm 2.3\text{e-}5$	Cassini/GR 2003	✓ Within Shapiro bound
Gravitational wave speed v_{GW}	BSFG: $v_{\text{GW}} = c \times (1 + k_{\eta}^2) \approx c + 10^{-226} \text{ m/s}$	GW150914 / GW170817:	$v_{\text{GW}}/c - 1$	$< 10^{-15}$
Perihelion precession (Mercury)	BSFG adds buoyancy correction $\delta\varphi = \kappa \times \varphi_{\text{GR}} \sim 10^{-6} \text{ arcsec/century}$	GR prediction: 43.03"/century; observed: 43.1"	GR + obs.	UQFF correction undetectable at current precision

New physics claim: BSFG (Buoyancy-Stratified Factorial Geometry) reproduces all tested GR predictions in the classical limit, while adding a vacuum buoyancy correction $\Delta g \sim 10^{-6} \text{ arcsec/century}$ to Mercury's perihelion. This is a falsifiable GR extension testable with future LISA or BepiColombo precision gravitational measurements.

Cite PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

§7 Connection to UQFF Framework

The Riemann tensor R^r_{0r0} of the BSFG geometry provides the **geometric encoding of tidal UQFF forces**. The buoyancy force F_U^{bi} represents the differential curvature between interior ($r < R_*$) and exterior ($r > R_*$) regions — a curvature discontinuity at the stellar boundary. The Aether field $\varepsilon(r) \propto r^{-3}$ creates a genuine non-flat geometry whose curvature $\propto r^{-5}$ decays rapidly away from the source, consistent with the UQFF fifth-force measurements reported in PAPER_413-418.

Hub: PAPER_554 (#149) is the first paper in the BSFG series. See PAPER_558 (#153) for the complete geometric system definition and PAPER_555 (#150) for metric compatibility and geodesic equation.