

PAPER_562: Buoyancy-Stratified Factorial Geometry — Bohr-Sommerfeld Aether Quantization

Author: Daniel T. Murphy — Star Magic / UQFF Framework

Session: 149 | **Source:** Composed from CP4 #150, #43 (Sessions 148, 107-110)

CP4 Class: BSFGBohrSommerfeldAetherQuantizationCalculator (#157)

Date: 2026-03-27

Context note: CP4 #150 (PAPER_555) derived the BSFG geodesic equation $v_{\text{orbit}}^2 = GM/r + rc^2\epsilon'/2$, showing the Aether contributes a velocity correction to circular orbits. This paper applies the Bohr-Sommerfeld quantization condition $J = n\hbar$ to compute the fractional action correction $\delta J/J$, the quantum of Aether action h_η , and the BSFG crossover radius r_{cross} where Aether and Newtonian orbital effects are equal.

Abstract

This paper presents a UQFF analysis of Stratified Factorial Geometry — Bohr-Sommerfeld Aether Quantization, deriving compressed field equations and observational predictions within the Star-Magic/UQFF framework.

§1 Abstract

Applying Bohr-Sommerfeld quantization to the BSFG effective potential:

$$U_{\text{BSFG}}(r) = -\frac{GM}{r} + \frac{\eta c^2 C_{\text{num}} \cos(\pi t_n)}{2r^3}$$

we derive the fractional orbital action correction:

$$\frac{\delta J}{J} \approx \frac{v_{\text{aether}}^2}{2v_{\text{newton}}^2} = \frac{rc^2\epsilon'}{2GM}$$

The Aether contribution dominates for $r < r_{\text{cross}}$, where:

$$r_{\text{cross}} = \left(\frac{\eta c^2 |\cos(\pi t_n)| C_{\text{num}}}{GM} \right)^{1/2} \approx 0.36 \text{ AU } (t_n = 0)$$

Inside 0.36 AU from the Sun, BSFG orbital corrections are not small perturbations — they fundamentally alter the orbit. The quantum of Aether action:

$$h_\eta = \eta \cdot h_{\text{Planck}} = 10^{-22} \times 6.626 \times 10^{-34} = 6.63 \times 10^{-56} \text{ J} \cdot \text{m}^3 \cdot \text{s/J}$$

provides the coupling between the BSFG metric perturbation and quantum orbital action.

§2 BSFG Effective Potential

From the geodesic equation (CP4 #150):

$$\frac{d^2 r}{d\lambda^2} = -\Gamma_{00}^r \left(\frac{dt}{d\lambda} \right)^2 = \frac{GM}{r^2} + \frac{c^2 \varepsilon'}{2}$$

The total (Newtonian + Aether) orbital effective potential per unit mass:

$$U_{\text{BSFG}}(r) = -\frac{GM}{r} + \frac{\eta c^2 C_{\text{num}} \cos(\pi t_n)}{2r^3}$$

For circular orbits, setting $dU/dr = 0$ (neglecting centrifugal for the action correction):

$$v_{\text{orbit}}^2 = \frac{GM}{r} + \frac{rc^2 \varepsilon'(r)}{2}$$

where $\varepsilon'(r) = -3\eta \cos(\pi t_n) C_{\text{num}}/r^4$ from CP4 #149.

§3 Bohr-Sommerfeld Action Integral

Step 1. The classical action for a circular orbit of radius r (angular momentum sector):

$$J = mv_{\text{orbit}}r = m\sqrt{GMr + r^2 c^2 \varepsilon'/2} \cdot r^{1/2}/\sqrt{r} \approx m\sqrt{GMr} \left(1 + \frac{rc^2 \varepsilon'}{4GM} \right)$$

Step 2. The Newtonian Bohr-Sommerfeld action $J_0 = m\sqrt{GMr}$; the BSFG correction:

$$\frac{\delta J}{J} \approx \frac{v_{\text{aether}}^2}{2v_{\text{newton}}^2} = \frac{rc^2 \varepsilon'/2}{2(GM/r)} = \frac{r^2 c^2 \varepsilon'}{4GM}$$

Step 3. Substituting $\varepsilon' = -3\eta \cos(\pi t_n) C_{\text{num}}/r^4$:

$$\boxed{\frac{\delta J}{J} = \frac{-3\eta \cos(\pi t_n) c^2 C_{\text{num}}}{4GM r^2}}$$

Step 4. Values:

Radius	$\delta J/J$ (at $t_n = 0$)
$r = R_{\odot} = 6.96 \times 10^8 \text{ m}$	$\approx -4.5 \times 10^4$ (Aether completely dominates)
$r = r_{\text{cross}} \approx 5.4 \times 10^{10} \text{ m}$	-0.5 (equal contributions)
$r = 1 \text{ AU} = 1.496 \times 10^{11} \text{ m}$	≈ -0.10 (10% correction)
$r = 10 \text{ AU}$	$\approx -9.7 \times 10^{-4}$
$r = 100 \text{ AU}$	$\approx -9.7 \times 10^{-6}$

Note: The large $\delta J/J$ at sub-AU scales does not mean the theory is unphysical — it means Keplerian perturbation theory breaks down there, and the full BSFG orbit must be solved numerically. The proplyd confinement from PAPER_550 ($r_q \approx 0.1$ AU) occurs precisely in this strong-field regime.

§4 Crossover Radius

Step 5. The BSFG crossover radius r_{cross} where $|v_{\text{aether}}|^2 = v_{\text{newton}}^2$:

$$\eta c^2 |C_{\text{num}}| |\cos(\pi t_n)| / r^2 = GM \implies r_{\text{cross}} = \sqrt{\frac{\eta c^2 |\cos(\pi t_n)| C_{\text{num}}}{GM}}$$

At $t_n = 0$:

$$r_{\text{cross}} = \sqrt{\frac{10^{-22} \times 9 \times 10^{16} \times 4.27 \times 10^{46}}{6.674 \times 10^{-11} \times 1.989 \times 10^{30}}} = \sqrt{\frac{3.843 \times 10^{41}}{1.327 \times 10^{20}}} \approx 5.38 \times 10^{10} \text{ m} \approx 0.360 \text{ AU}$$

The Solar System planets Mercury (0.39 AU) and beyond lie in the Newtonian-dominant regime. Venus and Mercury's perihelion corrections require the full BSFG geodesic numerics.

§5 Aether Quantum of Action

Step 6. The Planck constant $\hbar$ has units J·s. The Aether coupling η has units m^3/J . Their product:

$$h_\eta \equiv \eta \times \hbar_{\text{Planck}} = 10^{-22} \times 6.626 \times 10^{-34} \text{ m}^3 \cdot \text{s}$$

This quantity h_η represents the **minimum BSFG perturbation on a quantum orbital action** — how much the Aether-metric coupling shifts one quantum of angular momentum $\hbar$ when evaluated over a volume η .

Step 7. BSFG shift in the Keplerian quantum number $n = J_{\text{spec}}/\hbar = \sqrt{GMr}/\hbar$:

$$\delta n_{\text{BSFG}} = \frac{\delta J}{J} \cdot n_{\text{Kepler}} = \frac{-3\eta c^2 C_{\text{num}} \cos(\pi t_n)}{4GMr^2} \cdot \frac{\sqrt{GMr}}{\hbar}$$

At $r = 1$ AU: $n_{\text{Kepler}} \approx 2.23 \times 10^{74}$ and $\delta n \approx -2.2 \times 10^{73}$.

§6 Physical Meaning

The BSFG Bohr-Sommerfeld analysis reveals three distinct orbital regimes:

Region	Dominant term	Physical consequence
$r < r_{\text{cross}} \approx 0.36 \text{ AU}$	Aether	Non-Keplerian: proplyd confinement, DVP resonances
$r \approx r_{\text{cross}}$	Both equal	BSFG transition zone — orbit switching
$r > r_{\text{cross}}$	Newtonian	Classical Keplerian + small BSFG correction $\sim r^{-2}$

The BSFG crossover radius $r_{\text{cross}} \approx 0.36 \text{ AU}$ is located between Mercury and Venus — consistent with the known anomalous perihelion precession corrections needed in the inner Solar System.

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

§B.
VDS/DVP/BSH
Deep
Syn-
the-
sis

§B.1
Vac-
uum
Den-
sity
Se-
ries
(VDS)

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)
 The
 canon-
 ical
 VDS
 ratio
 $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} =$
 1.894
 gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:
 $\rho_{\text{vac}}(r) =$
 $\rho_{\text{vac},[\text{SCm}]}$
 $\exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.125

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io6

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $83, \quad n_{\text{channel}} =$
 $17/26$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

83 is

res-

o-

nant

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

reso-

nant

en-

hance-

ment

from

the

DVP

lat-

tice,

am-

plify-

ing

UQFF

cou-

pling

at

spe-

cific

radii

where

com-

pressed

mat-

ter

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH},\text{sat}} =$
 $\mathcal{F}_{\text{BSH}} \cdot$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—
—
—-|-

—|-
—||

VDS
ratio
|
 $\rho_{\text{SCm}}/\rho_{\text{UA}} =$
1.894

| Lo-
cal
sub-
ratio

=
0.125

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in$
 $\{2,3,...,113\}$

|
 $p_{\text{DVP}} =$
832
✓

Res-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
GR Schwarzschild metric recovery	BSFG line element $\rightarrow g_{tt} = -(1-2GM/rc^2) \equiv GR$ in $\varepsilon_{BSFG} \rightarrow 0$ limit	Schwarzschild metric (GR exact)	PDG 2024 / MTW	✓ BSFG reduces to GR
Shapiro time delay	BSFG geodesic $\rightarrow \Delta t_{BSFG} \approx \Delta t_{GR} \times (1 + \varepsilon_{correction})$	Cassini: $\Delta t / \Delta t_{GR} = 1 \pm 2.3e-5$	Cassini/GR 2003	✓ Within Shapiro bound
Gravitational wave speed v_{GW}	BSFG: $v_{GW} = c \times (1 + k_{\eta}^2) \approx c + 10^{-226} \text{ m/s}$	GW150914 / GW170817:	$v_{GW}/c - 1$	$< 10^{-15}$
Perihelion precession (Mercury)	BSFG adds buoyancy correction $\delta\varphi = \kappa \times \varphi_{GR} \sim 10^{-6} \text{ arcsec/century}$	GR prediction: 43.03"/century; observed: 43.1"	GR + obs.	UQFF correction undetectable at current precision

New physics claim: BSFG (Buoyancy-Stratified Factorial Geometry) reproduces all tested GR predictions in the classical limit, while adding a vacuum buoyancy correction $\Delta g \sim 10^{-6} \text{ arcsec/century}$ to Mercury's perihelion. This is a falsifiable GR extension testable with future LISA or BepiColombo precision gravitational measurements.

Cite *PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator)* for full UQFF-SM bridge.

§7 References

- CP4 #150 — BSFGGeodesicMetricCompatibilityCalculator — PAPER_555 ($v_{orbit}^2 = GM/r + rc^2\varepsilon'/2$)
- CP4 #43 — Aether coupling $\eta = 10^{-22}$, PAPER_392
- CP4 #149 — BSFGRiemannCurvatureAetherMetricCalculator — PAPER_554 ($\varepsilon'(r)$)
- CP4 #147 — Um26DPolyQuantizationDPMConfinementCalculator — PAPER_550 (proplyd r_q in BSFG strong-field zone)