

PAPER_785: Spirograph Nebula IC 418 — UQFF Planetary Nebula Fast Wind

Author: Daniel T. Murphy

Framework: UQFF (Universal Quantum Field Superconductive Framework)

Session: 181 | v5.42

Date: 2026

CP4 Class: #369 — SpirographNebulaIC418UQFFCalculator

Abstract

IC 418, the “Spirograph Nebula,” is one of the most intricately structured planetary nebulae, located $\sim 2,000$ light-years away ($z \approx 0.0007$) in the constellation Lepus. Hubble WFPC2 imaging reveals a complex pattern of nested shells and radial spokes resembling a spirograph drawing. The central white dwarf ($T_{\text{eff}} \sim 36,000$ K) drives a fast stellar wind at $\sim 1,500$ km/s ($v = 1.5 \times 10^6$ m/s), ionizing the ejected AGB envelope. Under UQFF, the fast wind velocity ($v = 1.5 \times 10^6$ m/s) with standard PN B-field ($B = 10^{-5}$ T) yields $g_{\text{IC418}} \approx 1.580 \times 10^{-2}$ m/s².

1. Introduction

IC 418 represents a late-stage AGB star ($\sim 1\text{--}3 M_{\odot}$ progenitor) that ejected its outer envelope $\sim 3,000$ years ago, creating the current planetary nebula shell. The Hubble image reveals the “spirograph” interference pattern from multiple overlapping AGB pulsation shells that were ejected at slightly different angles. The central star’s fast wind at $\sim 1,500$ km/s (confirmed by UV spectroscopy) is the highest drive velocity in the slow/fast wind PN interaction model. UQFF encodes this through $v = 1.5 \times 10^6$ m/s, which is $1.5\times$ the LBV eruptive velocity (and thus $15\times$ the standard HII velocity), yielding $g = 1.580 \times 10^{-2}$ m/s².

2. Master UQFF Gravity Equation

$$g_{\text{IC418}}(r, t) = (G \times M) / r^2 \times (1 + H(z) \times t) \times (1 - E_{\text{rad}}) \times (1 + f_{\text{TRZ}}) + a_{\text{EM}}$$

2.1 Parameters

Parameter	Symbol	Value	Source
Nebula mass (envelope)	M	$\sim 0.6 M_{\odot} = 1.193 \times 10^{30}$ kg	Hubble
Nebula radius	r	1×10^{16} m (~ 0.1 pc)	Hubble
Age	t	$\sim 3,000$ yr = 9.468×10^{10} s	Expansion age
E_rad	—	0.20	EUV photoionization loss
Redshift	z	0.0007	Distance
v_EM	v	1.5×10^6 m/s	Central star fast wind
B_EM	B	10^{-5} T	PN field
f_TRZ	—	0.05	UQFF

3. Long-Form Derivation

Step 1: Base Gravitational Term

$$g_{\text{grav}} = 6.6743\text{e-}11 \times 1.193\text{e}30 / (1\text{e}16)^2 = 7.962\text{e-}13 \text{ m/s}^2$$

Step 2: Cosmic Expansion Factor

$$H(z) \times t \approx \text{negligible} (2.268\text{e-}18 \times 9.468\text{e}10 \approx 2.15\text{e-}7); \text{ factor} \approx 1.0000002$$

Step 3: Radiation Energy Loss (EUV Photoionization)

$$E_{\text{rad}} = 0.20; 1 - E_{\text{rad}} = 0.80$$

Step 4: Time-Reversal Correction

$$f_{\text{TRZ}} = 0.05; 1 + f_{\text{TRZ}} = 1.05$$

Step 5: Gravitational Total

$$g_{\text{grav_total}} = 7.962\text{e-}13 \times 1.0 \times 0.80 \times 1.05 = 6.689\text{e-}13 \text{ m/s}^2$$

Step 6: Aether EM Correction (Fast Wind)

$$v = 1.5 \times 10^6 \text{ m/s}, B = 10^{-5} \text{ T}$$
$$a_{\text{EM}} = (1.602\text{e-}19 \times 1.5\text{e}6 \times 1\text{e-}5 / 1.673\text{e-}27) \times 11 \times 1\text{e-}12 = 1.580\text{e-}2 \text{ m/s}^2$$

Step 7: Final Solution

$$g_{\text{IC418}} = 6.689\text{e-}13 + 1.580\text{e-}2 \approx 1.580\text{e-}2 \text{ m/s}^2$$

4. Physical Interpretation

IC 418's result ($g = 1.580 \times 10^{-2}$) falls precisely between the standard HII (1.053×10^{-3}) and LBV eruptive (1.053×10^{-2}) regimes. The $1.5\times$ velocity multiplier (1.5×10^6 vs. 1.0×10^6 m/s) directly yields a $1.5\times$ larger result. IC 418 establishes the "fast wind PN" UQFF subcategory at $v = 1.5 \times 10^6$ m/s and will be shared with NGC 6307+7027 (PAPER_788) and M57 (PAPER_791) as the canonical planetary nebula fast-wind value.

5. Conclusions

UQFF applied to IC 418 Spirograph Nebula yields $g_{\text{IC418}} \approx 1.580 \times 10^{-2} \text{ m/s}^2$, establishing the planetary nebula fast-wind UQFF parameter class. The 1.5×10^6 m/s central star wind velocity is directly observed in UV spectroscopy, providing an observational anchor for the PN fast-wind UQFF subcategory.

PAPER_785, CP4 class #369. v5.42.

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.
 VDS/DVP/BSH
 Deep
 Syn-
 the-
 sis

 §B.1
 Vac-
 uum
 Den-
 sity
 Se-
 ries
 (VDS)
 The
 canon-
 ical
 VDS
 ratio

$$\rho_{\text{vac, [SCm]}}/\rho_{\text{UA}} = 1.894$$

gov-
 erns
 the
 double-
 exponential
 vac-
 uum
 con-
 den-
 sate
 pro-
 file:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac, [SCm]}} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right)$$

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

For

this

sys-

tem,

the

local

VDS

sub-

ratio

is

0.097

(near-

threshold

regime),

plac-

ing

it in

the

$t \rightarrow$

π

col-

lapse

zone

where

the

double-

exponential

tran-

si-

tions

sharply

from

con-

densed

to di-

lute

vac-

uum.

This

thresh-

old

be-

hav-

io4

con-

nects

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.2
 Dipole
 Vor-
 tex
 Primes
 (DVP)
 The
 DVP
 en-
 cod-
 ing
 maps
 the
 sys-
 tem's
 char-
 ac-
 teris-
 tic
 pa-
 ram-
 eter
 onto
 the
 prime
 lat-
 tice:
 $p_{\text{DVP}} =$
 $13,$ $n_{\text{channel}} =$
 6/26

##

§A.

Cosmogenesis-

Linked

La-

grangian

(PA-

PER_877

Sym-

bolic

Ex-

port)

Since

$p_{\text{DVP}} =$

13 is

sub-

threshold

(thresh-

old

at

$p >$

26),

the

sys-

tem's

vac-

uum

topol-

ogy

in-

her-

its

sub-

threshold

damp-

ing

from

the

DVP

lat-

tice,

pro-

duc-

ing

smooth

rather

than

reso-

nant

UQFF

cou-

pling

pro-

files.

The

DVP

frame-

work

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

 §B.3
 Buoy-
 ancy
 Sat-
 ura-
 tion
 Har-
 mon-
 ics
 (BSH)
 The
 BSH
 satu-
 ra-
 tion
 timescale
 for
 this
 sec-
 tor
 is
10⁴
yr
 (spin-
 down
 equi-
 lib-
 rium):
 $\mathcal{F}_{\text{BSH}} =$
 $\sum_{j=1}^{26} \frac{1}{j} \cdot$
 $f_{U_b} \cdot$
 $(1 - e^{-[SSq] \cdot m/M_{\odot}}).$
 $\cos\big(\frac{2\pi j}{26}\big)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

The
 tanh
 satu-
 ra-
 tion
 en-
 ve-
 lope
 pre-
 vents
 un-
 phys-
 ical
 di-
 ver-
 gence:
 $\mathcal{F}_{\text{BSH,sat}} =$
 $\mathcal{F}_{\text{BSH}}$
 $\left(1 - \tanh\left(\frac{t-t_{\text{sat}}}{\tau_{\text{BSH}}}\right)\right)$

 §A.
 Cosmogenesis-
 Linked
 La-
 grangian
 (PA-
 PER_877
 Sym-
 bolic
 Ex-
 port)

connecting
 to
 the
 PA-
 PER_877
 Stage
 5
 buoy-
 ancy
 seed

$$U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2).$$

f_{SCm}
 which
 ini-
 tial-
 izes
 the
 har-
 monic
 se-
 ries
 at
 cos-
 mo-
 gen-
 esis.

 §B.4
 Production-
 Scale
 Con-
 sis-
 tency

§A.
Cosmogenesis-
Linked
La-
grangian
(PA-
PER_877
Sym-
bolic
Ex-
port)

|
Frame-
work
|
Canon-
ical
Value

|
This
Pa-
per |
Sta-
tus |
|—

—
|—

—
—-|-

—|-

—||

VDS
ratio
|

$$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$$

| Lo-
cal
sub-
ratio

$$= 0.097$$

| ✓
Threshold-
consistent

||
DVP
prime

|
 $p_k \in \{2,3,...,113\}$
|

$$p_{\text{DVP}} = 130$$

✓
Sub-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	✓ Consistent
Proton decay rate	$\kappa = 0.0005/\text{day} \rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMPParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.