

PAPER #66 ■ Magnetar Systems: UQFF Predictions for SGR1745, Crab, Vela

Author: Daniel T. Murphy

PAPER #66 ■ Magnetar Systems: UQFF Predictions for SGR1745, Crab, Vela

Title: Magnetar Systems in the UQFF: Field Predictions for SGR1745, Crab, Vela, and ASKAP J1832-0911

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic ($\dot{\phi} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$) **Date:** March 7, 2026 **Source Data:** uqff_validation_test.py, observational_systems_config.h, SOURCE4 (SGR1745), MAIN_1_CoAnQi.cpp **Index Slot:** ■1.9 Automated 121-System Validation, Paper #66

Abstract

Magnetars are neutron stars with surface magnetic fields exceeding 10^{10} T ($B_{\text{crit,magnetar}} = 4.4 \times 10^{10}$ T), classifying them as the most extreme electromagnetic environments in the observable universe. The UQFF assigns each magnetar system all four operational modes (Compressed, Resonant, Buoyant, Superconductive) plus the Ug1 magnetic dipole enhancement. This paper presents UQFF predictions for SGR1745-2900 (canonical), the Crab Pulsar (PSRB0531+21), the Vela Pulsar, and ASKAP J1832-0911. The magnetic Ug1 dominates over standard Newtonian gravity by factors of 10^{10} to 10^5 , consistent with magnetar X-ray timing observations.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{\phi} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. System Parameters

System	M (kg)	r (m)	B0 (T)	$\dot{\phi}$ (rad/s)	Period
SGR1745-2900	2.785×10^{30}	2.62×10^4	2.3×10^{10}	1.671	3.76 s
Crab Pulsar	1.0×10^{30}	4.73×10^6	5.0×10^8	2.0×10^{-4}	~33 ms
Vela Pulsar	2.8×10^{30}	1.7×10^7	3.0×10^8	1.0×10^{-4}	~89 ms
ASKAP J1832	2.785×10^{30}	4.63×10^6	1.0×10^{10}	2.38×10^{-4}	44 min

2. Ug1 Magnetic Dipole Term

For each magnetar, the Ug1 term amplifies standard gravity:

$$Ug_1 = \frac{GM}{r^2} \cdot (1 + \delta_t) \cdot \frac{\mu_0 B_0^2}{8\pi}$$

System	g_Newton (m/s ²)	$\mu_0 B_0^2 / 8\pi$	Ug1 (m/s ²)	Amplification
SGR1745-2900	2.71×10^{-2}	1.33×10^4	3.60×10^2	0.13 (weak field region)

System	g_{Newton} (m/s ²)	B_0 /8p	U_{g1} (m/s ²)	Amplification
Crab Pulsar	2.99×10^7	3.14×10^7	9.37×10^7	negligible
Vela Pulsar	6.45×10^7	1.41×10^7	9.09×10^7	negligible
ASKAP J1832	8.67×10^4	5.00×10^6	4.34×10^8	5×10^6

ASKAP J1832-0911 has a magnetar-class surface field of $B_0 = 10^{10}$ T in the `uqff_validation_test.py` parameters, yielding an enormous U_{g1} enhancement. This represents the ultra-compact source (sub- 10^6 m scale) field contribution from the neutron star core.

3. FUBi_i Computations

LENR Resonance Term

The dominant driver of $FUBi_i$ at high (ω_{LENR}/ω_0) ratios:

$$\omega_{LENR} = k_{LENR} \left(\frac{\omega_{LENR}}{\omega_0} \right)^2, \quad \omega_{LENR} = 2\pi \times 1.25 \text{ THz} = 7.854 \times 10^{12} \text{ rad/s}$$

System	ω_0 (rad/s)	ω_{LENR}/ω_0	LENR term	$FUBi_i$ (N)
Vela	1.0×10^7	7.85×10^4	6.17×10^7	$\sim 8.3 \times 10^7$
Crab	2.0×10^7	3.93×10^4	1.54×10^5	$\sim 2.1 \times 10^7$
ASKAP J1832	2.38×10^7	3.30×10^5	1.09×10^8	$\sim 1.5 \times 10^7$
SGR1745	1.671	4.70×10^4	2.21×10^5	$\sim 3.0 \times 10^8$

Physical interpretation

The LENR term captures the resonance between the UQFF THz vacuum field ($\omega_{LENR} = 7.85 \times 10^{12}$ rad/s) and the astrophysical system's own oscillation frequency (ω_0). For slowly rotating or long-period systems (Vela, Crab), the ratio is enormous representing the extreme mismatch between the quantum vacuum oscillation timescale ($\sim 10^{-12}$ s) and the stellar spin period ($\sim 10^{-2}$ s to seconds). This gives the largest UQFF forces for slowly rotating compact objects.

4. SOURCE4 SGR1745 Canonical System

SGR1745-2900 is one of seven pre-defined astrophysical systems in the SOURCE4 namespace of MAIN1CoAnQi.cpp:

```
// SOURCE4 magnetar parameters (sgr1745_SOURCE4)
SGR1745.M = 2.785e30 kg // 1.4 M_sun neutron star
SGR1745.B = 2.3e10 T // Surface field
SGR1745.P = 3.76 s // Spin period
SGR1745.r = 2.62e20 m // Distance from SgrA* (~8.5 kpc)
// UQFF: Ug1 = (GM/r^2) * (1+d) * (B0/8p)
// F_U = SOURCE4::compute_FU_SOURCE4(sgr1745, r, t, tn, theta)
```

UQFF prediction for SGR1745:

- Ug1:** G-gravity $\ll (2.3 \times 10^{10})/8p$ = G-gravity $\ll 6.64 \times 10^8$? dominates over Newtonian
- Ug4 (vacuum BH coupling):** linked to SgrA* ($M_{BH} = 4 \times 10^6 M_{\text{sun}}$) at $d_g = 2.62 \times 10^{20}$ m

F_UQFF: Combined Compressed + Superconductive modes (nearest to BH uses Ug4 strongly)

5. Crab Pulsar Energy Budget

B_crit,magnetar = 4.4e10 T from index.js constants. The Crab surface field (~10? T) is sub-critical:

Quantity	Value
Crab B0 (surface)	~10? T
Bcrit/BCrab	~4.4e104 (sub-critical)
L_X (Crab total)	10? W
?_0 (33 ms pulsar)	~190 rad/s
UQFF Mode	Resonant dominant (33 ms pulse ? 190 Hz)

The Crab's fast spin (33 ms, ?0 ~ 190 rad/s, not the 2e10? rad/s in the config which is the orbital frequency) produces a lower LENR ratio than slower pulsars, meaning the Crab's FUBi_i is smaller in magnitude than Vela's consistent with the Crab being younger and more energetic (higher spin-down luminosity from Resonant mode, not static Compressed mode).

6. Vela Pulsar: UQFF Supernova Kick Prediction

Vela's very small ?0 = 10? rad/s in the config represents the orbital barycenter frequency of the PWN system. This produces the largest UQFF FUBi_i in the magnetar set: -8.3e10? N (comparable to the ensemble mean from Paper #63).

UQFF kick velocity prediction:

$$v_{\rm kick} = \frac{F_{U,Bi,i} \times \Delta t}{M} = \frac{8.3 \times 10^{219} \times 10^{-35}}{2.8 \times 10^{30}} \approx 296 \text{ km/s}$$

(using ?t ~ 10?5 s Planck-epoch impulse duration) ? Observation: Vela kick velocity ~ 60 km/s (range 60-350 km/s) ? UQFF is consistent with pulsar kick observations

Summary

System	B0 (T)	FUBi_i (N)	Dominant Mode	UQFF Status
Vela	3e10?8	-8.3e10??	Resonant	STABLE ?
Crab	5e10?8	-2.1e10??7	Resonant	STABLE ?
ASKAP J1832	10??	-1.5e10??	Compressed	STABLE ?
SGR1745	2.3e10??	-3.0e10?87	Compressed + Ug4	SOURCE4 validated ?

Source: uqffvalidationtest.py, observationalsystemsconfig.h, MAIN1CoAnQi.cpp SOURCE4 | ? = 0.0005/day | [SSq] = 0.57

See also: PAPER_065 | Part of the Star-Magic UQFF Whitepaper Series.

UQFF computed: Eddington luminosity UQFF correction = 1 - [SSq]exp(-?t) = 1 - 5.7e-1 exp(-2.9e-4) = 4.3e-1; F_U at event horizon = 2.0e+18 m/s.

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgradeearlywhitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-10} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{10} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{\{g1\}} + U_{\{g2\}} + U_{\{g3\}} + U_{\{g4\}} + U_{\{bi\}} + U_m - \sum_{i=1}^4 \lambda_i \text{ig}l[\lambda_i \cdot U_i(r,t) \cdot E_{\text{react}}] \text{igr}$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\sum \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420): $\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 U_m Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imes} \text{ig}l(1 + 10^{13} \backslash, \backslash \Theta(\text{ho}_{\text{SCm}} - \text{ho}_c) \text{igr}) \text{imes} \text{ig}l(1 + A_q \cos(\Delta \omega a, t) \text{igr})$$

Symbol	Value	Description
ρ_c	10^1 kg/m^3	SCm critical superconducting density

Symbol	Value	Description
A _q	0.1	Quasi-periodic beating amplitude (10%)
Δω	2π/(434·365.25) rad/day	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	U _g _sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	β _i × U _{bi}	Expanding nebulae, stellar winds
Superconductive	U _m × (1+10 ¹³ ·f _H)	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.