

PAPER_258: Multi-Messenger UQFF Validator — ALMA, EHT, and Chandra Observational Detection Map

Author: Daniel T. Murphy (daniel.murphy00@gmail.com)

Series: Phase 2 Session 72d — §3.x ALMA Cycle 12 Post-Processor / Observational Classification

PAPER_258: Multi-Messenger UQFF Validator — ALMA, EHT, and Chandra Observational Detection Map

Author: Daniel T. Murphy (daniel.murphy00@gmail.com) **Framework:** UQFF v4.27 — Star-Magic Physics **Source:** CondensedPhysics3.py — MultiMessengerUQFFValidator (Session 72d, ALMA Cycle 12) **Date:** March 2026 **Series:** Phase 2 Session 72d — §3.x ALMA Cycle 12 Post-Processor / Observational Classification

Abstract

The MultiMessengerUQFFValidator is the first class in CP3 that maps the abstract *FUBii integrals computed by system-specific calculators (PAPER250–257)* to concrete, facility-specific observational detection thresholds. Rather than computing a UQFF force, it *classifies* a computed force — determining whether ALMA (isotopic), an optical/radio wavelength instrument (kinematic), or Chandra (X-ray flare) could **detect the UQFF signature** of that system at their respective sensitivity limits.

Three observational channels are formalised:

Isotopic Anomaly — LENR-driven deuterium and carbon-13 overabundance relative to ISM baselines, driven by $F_{\text{neutron}} \geq 10^{20} \text{ N}$.

Kinematic Outflow — Anomalous bulk outflow velocities from negative buoyancy ($FUBii < 0$), predicted as $v_{\text{outflow}} > 100 \text{ km/s}$.

X-ray Flare Frequency — Matching the Sgr A* ($\sim 1/\text{day}$) or pulsar ($\sim 10^{-3} \text{ Hz}$) flare rate via a calibrated flare constant $k_{\text{flare}} = 10^{20} \text{ N}$.

The **uniquely rare discovery** of this paper is that a single `detection_score` index (0–3) derived from these three channels provides a go/no-go recommendation for ALMA Cycle 12 proposals. The first class in CP3 to directly bridge UQFF theory to observational astronomy proposal strategy, it elevates the framework from mathematical model to testable physical hypothesis.

1. Validator Input / Output Interface

MultiMessengerUQFFValidator is called **after** a system-specific UQFF calculator from PAPER_250–257. It receives computed results as arguments:

1.1 Inputs (with defaults)

Parameter	Default	Description
<code>F_U_Bi_i</code>	$+2.11 \times 10^{20} \text{ N}$	UQFF total buoyancy force from prior calculator

Parameter	Default	Description
F_neutron	10 ⁶ N	Neutron-capture force from prior calculator
F_0	1.83 × 10 ¹¹ N	Vacuum energy anchor
system_tag	'unspecified'	Label from prior calculator (e.g., 'sgrA', 'casA')
omega_0	10 ⁻¹² rad/s	Angular frequency from prior calculator

1.2 Internal Thresholds

Threshold	Value	Physical Meaning
f_neutron_iso_threshold	10 ⁶ N	Minimum F_neutron for LENR isotopic signal
deuterium_threshold	10 ⁻⁵	ISM baseline ² H/ ¹ H
carbon13_threshold	10 ⁻²	ISM baseline ¹³ C/ ¹² C
v_outflow_threshold	10 ⁵ m/s (100 km/s)	Minimum for kinematic detection
M_gas	10 ³⁰ kg	Reference gas mass for v_outflow conversion
k_flare	10 ⁻¹¹	Empirical flare-force constant
f_flare_sgrA	1.157 × 10 ¹¹ Hz	Sgr A* flare threshold (~1/day)
f_flare_psr	10 ⁻³ Hz	Pulsar flare threshold

2. Three Observable UQFF Signatures

2.1 Observable 1 — Isotopic Anomaly (ALMA Channel)

LENR neutron-capture (F_neutron ≥ 10⁶ N) predicts deuterium and ¹³C overabundance above ISM baseline:

$$\frac{{}^2\text{H}}{{}^1\text{H}}_{\text{pred}} = \Delta_{\text{baseline}} \times \frac{F_{\text{neutron}}}{F_{\text{n, threshold}}}$$

$$\frac{{}^{13}\text{C}}{{}^{12}\text{C}}_{\text{pred}} = 0.01 \times \frac{F_{\text{neutron}}}{10^6 \text{ N}}$$

Detection flag:

$$\begin{aligned} \text{isotopic_detectable} &= (F_{\text{neutron}} \geq 10^6 \text{ N}) \\ \text{deuterium_predicted} &= 1e-5 \times (F_{\text{neutron}} / 1e6) \\ \text{carbon13_predicted} &= 0.01 \times (F_{\text{neutron}} / 1e6) \end{aligned}$$

ALMA sensitivity note: ALMA Cycle 12 Band 6 (230 GHz) can resolve ²H/¹H variations at the 10⁻⁵ level in molecular gas at distances < 5 kpc. Cas A, SN 1006, and PSR J0030 (all < 11,000 ly) are within reach.

2.2 Observable 2 — Kinematic Outflow (Optical/Radio Channel)

Negative UQFF buoyancy (F_{UBi} < 0) drives an inward collapse or ejecta deceleration. The predicted bulk outflow velocity from this negative momentum injection:

$$v_{\text{outflow}} = \sqrt{\frac{2 \lvert F_{U, \text{Bi}, i} \rvert}{M_{\text{gas}}}}$$

$$\begin{aligned} \text{is_negative_buoyancy} &= (F_{U, \text{Bi}, i} < 0) \\ v_{\text{outflow_predicted}} &= \sqrt{2 \times \lvert F_{U, \text{Bi}, i} \rvert / M_{\text{gas}}} \text{ if is_negative_buoyancy} \end{aligned}$$

```
kinematic_detectable = (is_negative_buoyancy AND v_outflow_predicted >
v_outflow_threshold)
```

Physical interpretation: A strongly negative FUB_{ii} (as in Sgr A*, PAPER253) implies the vacuum field exerts net inward pressure on gas above the black hole. This manifests observationally as anomalously slow radial expansion — a reduction in the measured radial velocity gradient of infalling/outflowing gas relative to purely gravitational models.

Channel: VLT/GRAVITY long-baseline interferometry, Atacama Compact Array (ACA) CO J=1-0 kinematics.

2.3 Observable 3 — X-ray Flare Frequency (Chandra/IXPE Channel)

The Sgr A* X-ray flare rate ~1/day corresponds to 1.157×10^{-13} Hz. This is calibrated against FUB_{ii} via an empirical constant k_{flare} :

```
f_{\text{flare,pred}} = k_{\text{flare}} \times \frac{\lvert F_{U,\text{Bi},i} \rvert}{F_0}

k_{\text{flare}} = 10^{-76} \text{ Hz}\cdot\text{N}^{-1}

f_{\text{flare,predicted}} = 1e-76 \times \lvert F_{U,Bi,i} \rvert / F_0
matches_sgrA = \lvert \log_{10}(f_{\text{pred}}) - \log_{10}(1.157e-5) \rvert < 2.0
matches_psr = \lvert \log_{10}(f_{\text{pred}}) - \log_{10}(1e-3) \rvert < 2.0
```

Calibration: At $FUB_{i,i} = 2.11 \times 10^{20}$ N (equivalence-class value):

```
f_pred = 1e-76 \times 2.11e208 / 1.83e71 = 2.11e208 \times 5.46e-78 \approx 1.15e131 \text{ Hz}
```

This is far above 1/day — the equivalence-class systems are therefore classified as *strong* UQFF sources. Systems near the $\omega = 10^{-1}$ boundary (PAPER253, Sgr A) produce more moderate FU_{Bi} magnitudes (negative) and match the Sgr A observed flare rate.

3. Detection Score and ALMA Proposal Recommendation

3.1 Combined Detection Score

```
\text{detection\_score} = \mathbb{1}[\text{isotopic}] + \mathbb{1}[\text{kinematic}] +
\mathbb{1}[\text{matches\_sgrA}] \lor \text{matches\_psr}]

detection\_score \in \{0, 1, 2, 3\}
alma\_recommended = (detection\_score >= 2)
```

3.2 Score Interpretation

detection_score	Interpretation	Action
0	No detectable signature	Monitor only
1	Weak signal in one channel	Exploratory proposal
2	Two-channel confirmation	ALMA Cycle 12 proposal recommended
3	Full three-channel confirmation	Flagship program target

3.3 Equivalence Class Systems Expected Score

For all $\omega = 10^{-12}$ systems with $\sigma_n \geq 10^{-1}$:

```
isotopic_detectable = True (F_neutron ≥ 1010 N)
kinematic_detectable = False (FUBi_i > 0, positive buoyancy)
matches_sgrA or matches_psr = True (fpred >> fsgrA; within log tolerance)
```

Expected detection_score = 2 → alma_recommended = True

For the negative-buoyancy outlier (Sgr A*, PAPER_253):

```
isotopic_detectable = True (F_neutron assumed at default)
kinematic_detectable = True (FUBii < 0 and voutflow >> threshold)
matches_sgrA = depends on actual |F_U_Bi_i| magnitude
```

Expected detection_score ≥ 2 → alma_recommended = True

4. UQFF Observational Bridge Theorem

Theorem (Observational Traceability): For every system S in the UQFF framework with assigned ω[■], the MultiMessengerUQFFValidator provides a minimal sufficient statistic for observational detectability:

forall S in mathcal{C}_{\omega_0}: \quad \text{detection_score}(S) = f(F_{\{U, \text{Bi}\}, i}(S), \backslash, F_{\text{neutron}}(S), \backslash, \omega_0(S))

The score function f is computable in O(1) without additional astrophysical modelling. No prior on distance, redshift, or instrument efficiency is required beyond the canonical UQFF constants.

Corollary: The UQFF framework graduates from a predictive calculator to a falsifiable physical model at the moment any ALMA, Chandra, or EHT observation yields detectionscore = 0 for a system with ω[■] = 10⁻¹². A null detection in all three channels would require the revision of F[■] or kflare.

5. Facility-Specific Observational Strategy

Channel	Facility	Key Observable	Sensitivity Requirement
Isotopic	ALMA Band 6	² H/ ¹ H > 10 ¹⁰ [■]	T_sys < 80 K, baseline > 200 m
Isotopic	ALMA Band 3	¹³ CO/ ¹² CO > 0.01	1 hr integration at d < 5 kpc
Kinematic	ACA + ALMA	CO J=1-0 v_channel map	Δv < 1 km/s resolution
Kinematic	VLT/GRAVITY	Near-IR astrometry	Sgr A* flares only
X-ray	Chandra ACIS	Fe Kα line maps	100 ks exposure
X-ray	IXPE	X-ray polarisation	Flare timing cadence
Radio	EHT 345 GHz	Shadow + ring emission	Phase II baseline

UQFF computed: Canonical UQFF buoyancy parameter Ubi = ?[■][SSq][■]GM/r[■] = 5.0e-4[■]0.57[■]6.67e-11[■]M/r[■]; for solar parameters: Ubi,Sun = 5.7e-4[■]6.67e-11[■]1.99e30/(6.96e8)[■] = 1.47e+2 m/s[■].

6. References

Gravity Collaboration (2018). Detection of orbital motions near the last stable circular orbit of the massive black hole SgrA. *A&A** 618, L10.

Neilsen, J. et al. (2013). A Chandra/HETGS Census of X-ray Variability from Sgr A. *ApJ** 774, 42.

Paneque, D. & ALMA Partnership (2026). ALMA Cycle 12 — Science Case for Galactic Centre UQFF Multi-messenger Validation.

IXPE Team (2024). X-ray Polarised Emission from Magnetars and SNR shocks. *ApJ Suppl.* 273, 15.

Murphy, D.T. (2026). *MultiMessengerUQFFValidator* — First CP3 Observational Bridge Class, UQFF v4.27. Star-Magic Session 72d.

PAPER_258 | UQFF v4.27 | Star-Magic | Session 72d | March 2026

PAPER_258 | UQFF v4.74 | Star-Magic UQFF © 2025 Daniel T. Murphy