

PAPER_285: M16 Eagle Nebula UQFF #8212; Erosion Saturation Half-Time and Δg_{Max}

Author: Daniel T. Murphy | Date: March 2026

Session: 80 | Module: M16UQFFMODULE.cpp (22nd C++ UQFF module)

PAPER_285: M16 Eagle Nebula UQFF — Erosion Saturation Half-Time and Δg_{Max}

Photoevaporation Asymptotic Saturation: $t_{\text{half}} = \tau \cdot \ln 2 = 2.079 \text{ Myr}$

Classification: UQFF 2.0 Gravitational Physics — Nebular Erosion Dynamics **System:** M16 Eagle Nebula (IC 4703), Eagle Nebula Star-Forming Region **Session:** 80 | **Module:** M16UQFFMODULE.cpp (22nd C++ UQFF module) **Author:** Daniel T. Murphy | **Date:** March 2026

Abstract

This paper derives the **Erosion Saturation Half-Time** (t_{half}) and **Maximum Erosion Gravity Amplitude** (Δg_{Max}) for the M16 Eagle Nebula UQFF photoevaporation term. The photoevaporation rate follows an exponential saturation $E_{\text{rad}}(t) = E_{\text{0}} \times (1 - \exp(-t/\tau))$ with e-folding time $\tau = 3 \text{ Myr}$. The half-erosion time $t_{\text{half}} = \tau \cdot \ln 2 = 6.561 \times 10^{13} \text{ s} = 2.079 \text{ Myr}$ — the time at which the erosion fraction reaches half its asymptotic maximum E_{0} . The maximum gravity perturbation is $\Delta g_{\text{Max}} = E_{\text{0}} \times g_{\text{base}} = 4.36 \times 10^{-13} \text{ m/s}^2$. This is the **first UQFF module** to formally catalogue the photoevaporation half-time and asymptotic erosion concept.

2. Physical Background

UV radiation from massive O-type stars (such as those in the Young Stellar Cluster NGC 6611, embedded in M16) drives photoionisation of surrounding molecular gas — a process called **photoevaporation**. The erosion proceeds not as a linear ramp but as a **saturating exponential**:

$$E_{\text{rad}}(t) = E_{\text{0}} \left(1 - e^{-t/\tau}\right)$$

where:

$E_{\text{0}} = 0.3$ is the **asymptotic maximum fraction** (30% of mass eventually eroded)

$\tau = 3 \text{ Myr}$ is the **e-folding time** (photoevaporation efficiency timescale)

The saturation arises because dense molecular cores (pillar tips) are progressively shielded by their own column density as surrounding gas is stripped

3. Mathematical Derivation

3.1 Half-Time

The erosion half-time t_{half} is defined as the time when $E_{\text{rad}} = E_{\text{0}}/2$:

$$E_{\text{0}} \left(1 - e^{-t_{\text{half}}/\tau}\right) = \frac{E_{\text{0}}}{2}$$

$$1 - e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$e^{-t_{\text{half}}/\tau} = \frac{1}{2}$$

$$t_{\text{half}} = \tau \ln(2)$$

For M16:

$$t_{\text{half}} = 9.468 \times 10^{13} \text{ s} \times \ln(2) = 9.468 \times 10^{13} \times 0.6931 = 6.561 \times 10^{13} \text{ s}$$

$$t_{\text{half}} = \mathbf{2.079 \text{ Myr}}$$

3.2 Maximum Gravity Amplitude

The maximum erosion-induced gravity perturbation (asymptotic limit as $t \rightarrow \infty$):

$$\Delta g_{\text{Max}} = E_0 \times g_{\text{base}}$$

For M16:

$$\Delta g_{\text{Max}} = 0.3 \times 1.454 \times 10^{-12} = \mathbf{4.36 \times 10^{-13} \text{ m/s}^2}$$

3.3 Peak Erosion Rate

The instantaneous erosion rate at $t = 0$ (maximum rate, before saturation):

$$\frac{dE_{\text{rad}}}{dt}\bigg|_{t=0} = \frac{E_0}{\tau}$$

The corresponding gravity change rate:

$$\frac{dg_{\text{erode}}}{dt}\bigg|_{t=0} = \frac{E_0}{\tau} \times g_{\text{base}} = \frac{0.3}{9.468 \times 10^{13}} \times 1.454 \times 10^{-12} = \mathbf{4.61 \times 10^{-27} \text{ m/s}^2/\text{s}}$$

4. Saturation Profile

Time	t (s)	E_rad / E■	E_rad	g_erode (m/s ²)
0 Myr	0	0%	0	0
t_half = 2.079 Myr	6.561×10 ¹³	50%	0.150	2.18×10⁻¹³
τ = 3 Myr	9.468×10 ¹³	63.2%	0.190	2.76×10 ⁻¹³
5 Myr	1.578×10 ¹⁴	81.1%	0.243	3.54×10 ⁻¹³
∞ (asymptote)	→ ∞	100%	0.300	4.36×10⁻¹³

Key insight: At $\tau = 3$ Myr (the e-folding time), erosion has consumed only 63.2% of its capacity, NOT 100%. Half-erosion occurs earlier at 2.079 Myr. The pillar structure of M16 means the ~5700 ly "Pillars of Creation" are still observed today because erosion saturates — it cannot fully strip the densest pillar cores within observable timescales.

5. UQFF 2.0 Context

In the full M16 g_total equation, the erosion half-time governs the **temporal shape of g_dyn(t)**:

$$g_{\text{dyn}}(t) = g_{\text{base}} \times (1 + M_{\text{sf}}) \times (1 - E_0(1 - e^{-t/\tau}))$$

The transition from rapid to slow erosion occurs at $t_{half} = 2.079 \text{ Myr}$. For the UQFF simulation (t stepping from 0 to t_{max}), the half-time provides a natural **inflection point** in the dynamic gravity trajectory — before t_{half} , erosion is dominant; after t_{half} , star formation accumulation dominates (since M_{sf} grows linearly while E_{rad} asymptotes).

Crossover Time

The era when SFR growth exactly compensates erosion ($d\Phi_{dm}/dt = 0$ — maximum Φ_{dm}):

$$\frac{d\Phi_{dm}}{dt} = \text{SFR_rate} \times (1 - E_{rad}) - (1 + M_{sf}) \times \frac{E_0}{\tau} e^{-t/\tau} = 0$$

This crossover defines when the Eagle Nebula achieves maximum effective gravitational influence, after which continued SFR growth dominates erosion.

6. Wolfram KB Term

```
M16UQFF:t_half=tau*Log[2]=6.561e13s=2.079Myr; DeltaGMax=E_0*g_base=4.36e-13 m/s^2
[PAPER_285]
```

7. Cross-References

****PAPER_284:**** Dual Mass Co-Action Product ($\Phi_{dm} = (1+M_{sf}) \times (1-E_{rad})$)

****PAPER_286:**** Nebular Friedmann Redshift (κ_{neb} , $z=0.0015$)

M16UQFFMODULE.cpp: Full UQFF 2.0 C++ implementation (22nd module)

CondensedPhysics3.py: M16ErosionSaturationHalfTimeCalculator

Copyright — Daniel T. Murphy, Session 80, March 2026. UQFF 2.0.