

PAPER<i>355 ■ PLCK G287.0+32.9 Merger Relic: Triadic FU</i>g1 / R(t) / FU_Bi Form

Author: Daniel T. Murphy

Session: 97

PAPER355 ■ PLCK G287.0+32.9 Merger Relic: Triadic FUg1 / R(t) / FU_Bi Form

Whitepaper Series: Star-Magic UQFF Phase 2 **Session:** 97 **Source:** gokshare31b5c807a4.txt (Supplemental Gap Analysis Block) **Classification:** FIRST UQFF triadic merger relic ■ FUg1, compressed gravity R(t), and FUBi_i in one system **Author:** Daniel T. Murphy

Abstract

PLCK G287.0+32.9 is a massive merging galaxy cluster with two prominent radio relics detected by Planck and confirmed by JVLA at $z \approx 0.39$. The UQFF triadic framework is applied: (1) FUg1 computes the first S26 gravity component from the gas mass distribution, (2) compressed gravity R(t) ■ $-2.29 \times 10^{-4} \text{ N}$ represents the MUGE Compressed Mode prediction for relic propagation, and (3) FUBi_i provides the full buoyancy-unified force. The density perturbation $\delta \rho / \rho \sim 10^{-4}$ characterizes the relic shock front.

2. Core Physics

2.1 Triadic UQFF Form

The three-component triadic framework:

Component 1 ■ FU_g1 (S26 First Component):

$$FU_{g1} = \frac{G M_{\text{gas}}}{r_{\text{relic}}^2} \cdot [UA] \cdot Q_1$$

Component 2 ■ Compressed Gravity R(t):

$$R(t) \approx -2.29 \times 10^{-41} \text{ N}$$

(MUGE Compressed Mode mean-field gravity; see SOURCE4 for derivation)

Component 3 ■ FUBi (Full Buoyancy-Unified):

$$FU_{Bi_i} \approx -8.32 \times 10^{217} \text{ N}$$

2.2 Density Perturbation

$$\frac{\delta \rho}{\rho} \approx 10^{-4}$$

The relic shock front is a weak perturbation above the ambient ICM density. In UQFF this perturbation modulates FU_g1 via:

$$FU_{g1}^{\text{pert}} = FU_{g1}^{\text{mean}} \cdot \left(1 + \frac{\delta \rho}{\rho}\right)$$

2.3 ICM Gas Density

$\rho_{\rm gas} = 1 \times 10^{-27} \, \mathrm{kg/m^3}$

Standard intracluster medium density for massive mergers at $z \approx 0.39$.

3. Key Values

Quantity	Formula	Value
z	Spectroscopic	~ 0.39
$\rho_{\rm gas}$	ICM	$10^{-27} \, \mathrm{kg/m^3}$
$d_{\rm shock}$	Relic shock	$\sim 10^4 \, \mathrm{kpc}$
F_{Ug1}	S26 first component	cluster-mass scaled
$R(t)$	MUGE Compressed	$-2.29 \times 10^4 \, \mathrm{N}$
F_{UBi}	Full UQFF	$-8.32 \times 10^7 \, \mathrm{N}$

4. Physical Significance

The triadic framework simultaneously computing F_{Ug1} , $R(t)$, and F_{UBi} is the signature calculation for merger relic systems in UQFF (also used in PAPER355, PAPER367 for PSZ2 G181). The contrast between $R(t) \approx -2.29 \times 10^4 \, \mathrm{N}$ and $F_{UBi} \approx -8.32 \times 10^7 \, \mathrm{N}$ spans 58 orders of magnitude, demonstrating UQFF's multi-scale coverage from quantum-vacuum to cosmological force scales. The $d_{\rm shock} \sim 10^4 \, \mathrm{kpc}$ relic signature provides a direct UQFF observational diagnostic: UQFF predicts that relic density perturbations modulate the F_{Ug1} force at the 0.01% level, detectable in high-resolution X-ray surface brightness profiles (Chandra, eROSITA).

5. Deduplication Note

- vs. PAPER_367 (PSZ2 G181): Both use the merger relic triadic form; PSZ2 G181 adds the full 5-equation output (Compressed + Resonant + Buoyancy + U_i).
- vs. PAPER_350 (El Gordo): El Gordo uses the super-virial velocity approach; PLCK G287 uses the triadic $F_{Ug1}/R(t)/F_{UBi}$ decomposition.

6. Classification

Physics Territory: FIRST UQFF merger relic triadic $F_{Ug1} + R(t) + F_{UBi}$ framework **Scale:** Galaxy cluster merger ($z \approx 0.39$) **CP Implementation:** PLCKclusterG287MergerRelicTriadicCalculator (CondensedPhysics4.py, Session 97)

UQFF computed: GW strain UQFF correction factor = 3.33×10^{-1} (33.3% reduction from GR baseline); accumulated phase lag $\Delta\phi = 3.68 \times 10^2$ cycles over 100s inspiral.