

PAPER403 — Ubi: 4-Term Solar Wind Buoyancy Decomposition with ϵ_{sw}

Session: 108 (groksharecfdcad2f5.txt construction file re-analysis)

PAPER403 — Ubi: 4-Term Solar Wind Buoyancy Decomposition with ϵ_{sw}

Source: groksharecfdcad2f5.txt, lines 277–1600 ("Star Magicconstruction file04Oct2025.docx" C++ implementation) **Section:** C++ source — compute_Ubi() function as 4-term sum over all Ugi components **Session:** 108 (groksharecfdcad2f5.txt construction file re-analysis) **CP4 Class:** Ubi4TermSolarWindBuoyancyEpsilonSwCalculator (#52)

1. Overview

PAPER394 defined the buoyancy force $U_{\{bi\}}$ applied to a single Ugi term. PAPER403 extracts the **complete 4-term Ubi decomposition** from the construction file:

$$U_{\{bi\}} = U_{\{bi,1\}} + U_{\{bi,2\}} + U_{\{bi,3\}} + U_{\{bi,4\}}$$

where each term applies the buoyancy formula independently to $U_{\{g1\}}$, $U_{\{g2\}}$, $U_{\{g3\}}$, $U_{\{g4\}}$.

The novel addition is the **solar wind buoyancy modulation** $(1 + \epsilon_{sw} \cdot \rho_{sw})$, a first-principles coupling between solar wind plasma density and UQFF buoyancy response.

2. Formula

2.1 Individual Ubi Term

$$U_{\{bi,k\}} = -\beta_i \cdot U_{\{g,k\}} \cdot \Omega_g \cdot \frac{M_{\{bh\}}}{d_g} \cdot (1 + \epsilon_{sw} \cdot \rho_{sw}) \cdot U_{\{UA\}} \cdot \cos(\pi t_n)$$

where $k = 1, 2, 3, 4$ and $U_{\{g,k\}}$ is the corresponding $U_{\{g1\}}$, $U_{\{g2\}}$, $U_{\{g3\}}$, $U_{\{g4\}}$.

2.2 Total Buoyancy

$$U_{\{bi,\text{total}\}} = \sum_{k=1}^4 U_{\{bi,k\}} = -\beta_i \cdot \Omega_g \cdot \frac{M_{\{bh\}}}{d_g} \cdot (1 + \epsilon_{sw} \cdot \rho_{sw}) \cdot U_{\{UA\}} \cdot \cos(\pi t_n) \cdot \left(U_{\{g1\}} + U_{\{g2\}} + U_{\{g3\}} + U_{\{g4\}} \right)$$

3. Parameters

Symbol	Value	Notes
β_i	0.6	Buoyancy coefficient (PAPER_394)
Ω_g	7.3×10^{-16} rad/s	Galactic angular frequency
$M_{\{bh\}}$	8.155×10^{36} kg	Sgr A*
d_g	2.62×10^{20} m	GC distance
ϵ_{sw}	10^{-3}	Solar wind buoyancy coupling coefficient

Symbol	Value	Notes
ρ_{sw}	$8 \times 10^{-21} \text{ kg/m}^3$	Solar wind plasma density at 1 AU
U_{UA}	1.0	Unified Aether field (default)
t_n	normalized time	$\cos(\pi t_n)$ oscillation

4. Novel Physics

4.1 Solar Wind Buoyancy Modulation ($1 + \epsilon_{sw} \cdot \rho_{sw}$)

The modulation factor:

$$1 + \epsilon_{sw} \cdot \rho_{sw} = 1 + (10^{-3})(8 \times 10^{-21}) = 1 + 8 \times 10^{-24} \approx 1.000000000000000000000008$$

At standard solar wind conditions, this is near-unity (0.8 parts in 10^{23} amplification). However, during **solar energetic particle events** and **coronal mass ejections**, ρ_{sw} can increase by 6-8 orders: $\rho_{sw, \text{CME}} \sim 10^{-13} \text{ kg/m}^3$.

At CME density: $1 + \epsilon_{sw} \cdot \rho_{sw, \text{CME}} = 1 + 10^{-16}$ — still small but potentially measurable in precision Ubi experiments.

4.2 4-Term Ubi Architecture

Prior formulations used a single U_{bi} applied to the total U_g . The construction file reveals that **each Ugi component generates its own buoyancy response independently**:

$$U_{bi,k} \propto U_{g,k}$$

This creates a buoyancy **cascade**: large U_{g2} (from E_{react} amplification) generates the dominant buoyancy contribution, while U_{g4} (scale-invariant at $4.219 \times 10^{-10} \text{ m/s}^2$) generates a universal background buoyancy.

4.3 Comparison: Sum vs Previous Single-Term

With the 4-term sum, total U_{bi} at Sun:

$$U_{bi, \text{total}} \approx -\beta_i \cdot \Omega_g \cdot \frac{M_{bh}}{d_g} \cdot 1.0 \cdot U_{UA} \cdot (U_{g1} + U_{g2} + U_{g3} + U_{g4})$$

The dominant term will be $U_{bi, 2}$ (from U_{g2}) due to E_{react} amplification. $U_{bi, 4}$ provides the universal solar-wind-modulated vacuum background.

5. Solar Wind Density Scaling

Condition	ρ_{sw} (kg/m ³)	$(1 + \epsilon_{sw} \cdot \rho_{sw})$
Calm solar wind (1 AU)	8×10^{-21}	$1 + 8 \times 10^{-24}$
Active solar wind	8×10^{-20}	$1 + 8 \times 10^{-23}$
Solar storm (CME)	$\sim 10^{-17}$	$1 + 10^{-20}$
Solar minimum (outer helio)	$\sim 10^{-23}$	$1 + 10^{-26}$

The ϵ_{sw} term provides UQFF **solar weather sensitivity** — a direct modulation of galactic-scale buoyancy by local solar wind conditions.

6. C++ Source

```
// grok_share_cfdcad2f5.txt construction file
double beta_i = 0.6;
double epsilon_sw = 1e-3;
double rho_sw = 8e-21; // solar wind density at 1 AU
double wind_mod = 1.0 + epsilon_sw * rho_sw; // ≈ 1 + 8e-24
double Omega_g = 7.3e-16;
double UUA = 1.0;
double cos_factor = cos(M_PI * t_n);

// 4-term buoyancy sum
double Ubi1 = -beta_i * Ug1 * Omega_g * (Mbh / dg) * wind_mod * UUA * cos_factor;
double Ubi2 = -beta_i * Ug2 * Omega_g * (Mbh / dg) * wind_mod * UUA * cos_factor;
double Ubi3 = -beta_i * Ug3 * Omega_g * (Mbh / dg) * wind_mod * UUA * cos_factor;
double Ubi4 = -beta_i * Ug4 * Omega_g * (Mbh / dg) * wind_mod * UUA * cos_factor;
double Ubi_total = Ubi1 + Ubi2 + Ubi3 + Ubi4;
```

7. Relationship to Prior Papers

Paper	Ubi Form	Notes
PAPER_198	$U_{\text{bi}} = -\beta_i \cdot U_{\text{g1}} \cdot \Omega_g \cdot (M/r) \cdot [UA] \cdot \cos(\pi t_n)$	Single-term, compact object form
PAPER_394	FU master sum includes total U_{bi}	Simplified sum
PAPER_403	4-term U_{bi} with ϵ_{sw} solar wind	NEW — FIRST ϵ_{sw} solar wind buoyancy

Whitepaper generated Session 108. Source: groksharecfdcad2f5.txt lines 277-1600.