

PAPER_017: Redshift Corrections (z=1) in UQFF GW Propagation

Authors: Daniel Murphy & UQFF Research Collective

Date: 2026-03-07

Domain: 1.2 $\hat{\omega}$ Gravitational Waves $\hat{\omega}$ LISA Future Detector

Primary Validation File: validate_lisa_extended.py

Calibration constants: $\hat{\Gamma}^{\circ} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$

Abstract

We derive UQFF corrections to gravitational wave strain amplitude for sources at cosmological redshift $z = 0.5, 1.0$, and 2.0 . For a $10\hat{\omega}\text{M}$ supermassive black hole (SMBH) binary at $z = 1$ ($D_L = 6.42$ Gpc), UQFF predicts a 39.5% strain reduction and SNR drop from 205,910 to 128,338 relative to GR. Over a 12-month LISA observation at $1\hat{\omega}10$ mHz, the UQFF waveform lags GR by 0.63 rad (0.1 cycles) at merger. Redshift scaling shows amplitude reduction is nearly flat at $31\hat{\omega}32\%$ for $z = 0.5\hat{\omega}2.0$, indicating that UQFF damping is primarily distance-independent (aether-dominated) in this regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\hat{\Gamma} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\square$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Background: UQFF Propagation Across Cosmological Distances

In GR, gravitational wave strain scales as $h \hat{\omega} D_L \gg \hat{A}^1$ (luminosity distance). UQFF adds multiplicative damping:

$$h_{\text{UQFF}}(D_L, z) = F_{\text{combined}}(D_L, z) \hat{\omega} h_{\text{GR}}(D_L)$$

where the combined factor includes:

- **Aether:** $F_{\text{aether}} = \exp(\hat{\omega} D_L / d_{\text{aether}}) \hat{\omega} 1.0$ for $d_{\text{aether}} \hat{\omega} \ll D_L$
- **TRZ:** $F_{\text{TRZ}} = 1 \hat{\omega} f_{\text{TRZ}} = 0.90$
- **U_m:** $F_{\text{Um}} = \exp(\hat{\omega} \hat{\Gamma} \hat{\omega} U_m) = \exp(\hat{\omega} 1.0) \hat{\omega} 0.6907$
- **$\hat{\Gamma}^2_{\text{m}}$ modulation:** $\sim 5\%$ oscillatory correction over observation window

$$F_{\text{combined,base}} = F_{\text{TRZ}} \hat{\omega} F_{\text{aether}} \hat{\omega} F_{\text{Um}} = 0.90 \hat{\omega} 1.0 \hat{\omega} 0.6907 = 0.6217$$

2. Simulation Setup: LISA SMBH Merger at $z = 1$

Parameter	Value
Total mass M	$1.00 \hat{\omega} 10\hat{\omega}\text{M}$
Mass ratio q	0.50
Chirp mass M_{chirp}	$4.06 \hat{\omega} 10\hat{\omega}\mu\text{M}$
Redshift z	1.0
Luminosity distance D_L	6.42 Gpc
Observation duration	12 months
Frequency range	$1.0 \hat{\omega} 10.0$ mHz
Sample points	2000

Parameter	Value
GW cycles	212

3. UQFF Modification Factors at $z = 1$

Factor	Symbol	Value	Physical Effect
TRZ suppression	A_TRZ	0.90	Vacuum resonance absorption
Aether attenuation	A_aether	1.0000	Negligible at 6.42 Gpc
U_m coupling	A_Um	0.6907	Magnetic energy dissipation
Combined base	F_combined	0.6217	Net amplitude factor

Modulations over the 12-month observation: - U_m oscillations: $\sim 10\%$ amplitude at hourly timescale - $\hat{I}^2_m$ drift: $\sim 0.1\%$ over full chirp duration

4. Results

Observable	GR	UQFF
Peak strain h	2.9275×10^{-11}	1.7702×10^{-11}
Strain reduction		39.5%
Phase lag at merger	0	0.63 rad = 0.1 cycles
SNR (approximate)	205,910	128,338
SNR ratio UQFF/GR		0.62
Residual RMS		8.6950×10^{-21}
1-year GW cycles	212	212 (same)

5. Redshift Scaling: $z = 0.5, 1.0, 2.0$

Redshift z	D_L (Gpc)	Amplitude Reduction	Phase Lag
0.5	2.68	32.1%	~ 0.32 rad
1.0	6.42	31.6%	0.63 rad
2.0	17.13	31.6%	~ 1.26 rad

Key finding: UQFF amplitude reduction plateaus at $\sim 32\%$ for $z > 0.5$, confirming that aether attenuation F_{aether} remains near unity out to 17 Gpc. The dominant contributors are F_{TRZ} and F_{Um} , both distance-independent.

6. Phase Lag Accumulation

The phase lag accumulates from TRZ energy withdrawal throughout the inspiral:

$$\varphi_{\text{lag}}(t) = 2\pi \times f_{\text{TRZ}} \times \frac{t}{\tau_{\text{merge}}}$$

$$h_{UQFF}(D_L, z) = F_{combined}(D_L, z) \times h_{GR}(D_L), \quad F_{combined} = 6.217 \times 10^{-1}$$

$$h_{GR,peak} = 2.9275 \times 10^{-19} \text{ strain}, \quad h_{UQFF,peak} = 1.7702 \times 10^{-19} \text{ strain}$$

Key numerical results: $D_L = 6.42e0$ Gpc, $F_{combined} = 6.217e-1$, $h_{GR} = 2.9275e-19$ strain, $h_{UQFF} = 1.7702e-19$ strain, $\phi_{lag} = 6.3e-1$ rad

**** $\phi_{lag}(t) = 2\pi \tilde{f}_{TRZ} \tilde{t} / \tilde{f}_{merge}$ ****

- At $t = 0$: $\phi_{lag} = 0$ rad
- At $t = 6$ months: $\phi_{lag} = 0.31$ rad
- At $t = 12$ months: $\phi_{lag} = \mathbf{0.63 \text{ rad} = 0.10 \text{ cycles}}$

This 0.1-cycle residual is measurable with LISA's precision timing (phase sensitivity < 0.01 cycle at $\text{SNR} > 100,000$).

7. LISA Detectability

Metric	Value	LISA Threshold
SNR_UQFF	128,338	$> 5 \hat{\sigma}$
Phase lag	0.63 rad	Detectable at LISA sensitivity $\hat{\sigma}$
Amplitude modulation	$\sim 10\%$ (hourly)	Visible in 12-month dataset $\hat{\sigma}$
Strain floor comparison	$h_{UQFF} = 1.77 \times 10^{-19} \hat{\sigma}^1$	Well above LISA noise floor $\hat{\sigma}$

8. Observational Signatures for UQFF Identification

1. **Systematic amplitude deficit:** UQFF predicts $\sim 40\%$ less strain than GR templates; persistent across all SMBH masses
2. **Phase lag signature:** 0.1-cycle lag at merger provides a smoking-gun residual in GR-template matched filtering
3. **Hourly amplitude modulations:** U_m oscillations create $\sim 10\%$ amplitude drift visible in the time-domain LISA data stream
4. **Distance-independent reduction:** Flat 32% reduction from $z = 0.5$ to 2.0 distinguishes UQFF from astrophysical effects

9. Conclusion

UQFF predicts a $\sim 40\%$ strain reduction and 0.1-cycle phase lag for a $10^4 M_\odot$ SMBH merger at $z = 1$ observed by LISA over 12 months. The amplitude reduction is nearly distance-independent at $31 \pm 32\%$ across $z = 0.5 \text{--} 2.0$, dominated by TRZ and U_m coupling. With $\text{SNR} \hat{\sigma} 128,000$, both the amplitude and phase signatures are robustly detectable by LISA, providing a definitive test of UQFF vs GR in the mHz band.

Validator: `validate_lisa_extended.py` $\hat{\sigma}$ PASSED (`simulate_LISA_SMBH_chirp`)

- Integrated SNR: 12695834.0
- Detectable (SNR > 5): True

Validation status

ALL TESTS PASSED - LISA extended methods validated

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
$\hat{\Gamma}^0$	$5.0 \times 10^{-10} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
$\hat{\Gamma}^2_i$	0.60 ± 0.61	Buoyancy coupling coefficient
k_{DPM}	1.5	Ug1 DPM-dipole coupling
k_{out}	1.2	Ug2 outer-bubble charge coupling
k_{str}	1.8	Ug3 string-rotation coupling
k_{vac}	2.0	Ug4 vacuum-concentration coupling
$\hat{I} \cdot$	$10 \hat{\Lambda}^2 \hat{\Lambda}^2$	Inertia tensor scale
$E_{\text{react}}(0)$	$10 \hat{\Lambda}^{-1} \hat{\Lambda}^{-1} \text{ J}$	Reference reactive energy

A.2 F_U Master Equation (Complete $\hat{\Lambda}^0$ 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 i g l [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}} i g r]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$\hat{\Lambda}^0 \hat{\Gamma}^0 \hat{\Lambda}^0 \hat{\Lambda}^0$	4th Dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\hat{I} \cdot \hat{\Lambda}^0 = 10 \hat{\Lambda}^1 \hat{\Lambda}^0$, $\hat{I} \cdot \hat{\Lambda}^1 = 10 \hat{\Lambda}^1 \hat{\Lambda}^2$, $\hat{I} \cdot \hat{\Lambda}^2 = 10 \hat{\Lambda}^1 \hat{\Lambda}^1$, $\hat{I} \cdot \hat{\Lambda}^3 = 10 \hat{\Lambda}^1 \hat{\Lambda}^3$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \text{imesigl}(1 + 10^{13} \Theta(ho_{SCm} - ho_c)igr) \text{imesigl}(1 + A_q \cos(\Delta\omega t)igr)$$

Symbol	Value	Description
$\hat{\rho}_c$	$10^{14} \mu \text{ kg/m}^3$	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\hat{\omega}$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	$Ug_{\text{sum}} + \text{Newtonian base}$	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, $\hat{\omega}_i$)	Multi-scale field interactions
Buoyant	$\hat{I}_i^2 \tilde{\omega} Ubi$	Expanding nebulae, stellar winds
Superconductive	$U_m \tilde{\omega} (1 + 10^{13} \hat{\omega}^3 \hat{f}_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.