

PAPER_570: DVP Prime Vortex Photon-Photon Scattering in Olbers Framework

Unified Quantum Field Framework (UQFF)

Daniel T. Murphy • UQFF Research Consortium • 2026

Source: PAPER_570_Olbers_DVP_PhotonPhoton_PrimeVortex_Scatter.md

PAPER_570: DVP Prime Vortex Photon-Photon Scattering in Olbers Framework

Author: Daniel T. Murphy — Star Magic / UQFF Framework **Session:** 153b **Gap-Fill For:** AldersOlbersBSFGMetricGapAnalysisCalculator (#160, PAPER_566) — Completed Extension 4 **Date:** 2026-03-29 **QS:** 5/5

§1 Abstract

Beyond dust opacity, photons are attenuated by **photon-photon scattering** — the Breit-Wheeler process $\gamma\gamma \rightarrow e^+e^-$ — particularly for TeV photons from distant blazars. In the UQFF framework, this process is modulated by **Dipole Vortex Prime (DVP) vortex encompassments** at primes $p > 26$: each prime lattice point acts as a scattering centre with amplitude $A(p) \propto [\text{SSq}]^{\pi(p)} / p^{26}$. This paper derives the DVP-modulated mean free path $\ell_{\gamma\gamma}$ and its contribution to Olbers suppression.

$$\ell_{\gamma\gamma}^{\text{DVP}}(p_{\text{anchor}}) = \frac{r_H}{\pi_{\text{count}}} \cdot p_{\text{anchor}}^{26} \cdot [\text{SSq}]^{-\pi(p_{\text{anchor}})}$$

where $p_{\text{anchor}} = 113$ (H proto-shell prime).

§2 Breit-Wheeler Process

The standard Breit-Wheeler $\gamma\gamma \rightarrow e^+e^-$ cross section:

$$\sigma_{\text{BW}} = \pi r_e^2 (1-\beta^2) \left[2\beta(\beta^2-2) + (3-\beta^4)\ln\frac{1+\beta}{1-\beta} \right]$$

where $\beta = (1 - m_e^2 c^4 / E_{\text{cm}}^2)^{1/2}$, $r_e = 2.818 \times 10^{-15}$ m.

Near threshold ($E_{\text{cm}} \approx 2 m_e c^2$), $\sigma_{\text{BW}} \approx 1.7 \times 10^{-29}$ m².

TeV photon mean free path through CMB photons:

$$\ell_{\gamma} = \frac{1}{n_{\gamma} \sigma_{\text{BW}}} \approx \frac{1}{(4.1 \times 10^8)(1.7 \times 10^{-29})} \approx 1.4 \times 10^{20} \text{ m} \approx 4.5 \text{ Mpc}$$

§3 DVP Vortex Modulation

In the UQFF DVP lattice (PAPER_429), primes $p > 26$ define vortex scattering centres. The amplitude at prime p :

$$A(p) \propto \frac{[\text{SSq}]^{\pi(p)}}{p^{26}}$$

with $\pi(p)$ = count of primes $\leq p$.

The DVP-modulated mean free path:

$$\ell_{\text{DVP}}(p) = \frac{r_H}{\pi_{\text{count}}} \cdot \frac{p^{26}}{[\text{SSq}]^{\pi(p)}}$$

For the anchor prime $p_{\text{anchor}} = 113$, $\pi(113) = 30$:

$$A(113) = \frac{(0.507)^{30}}{113^{26}} \approx \frac{9.1 \times 10^{-10}}{8.5 \times 10^{53}} \approx 1.1 \times 10^{-63}$$

$$\ell_{\text{DVP}}(113) = \frac{4.4 \times 10^{26}}{149} \cdot \frac{113^{26}}{(0.507)^{30}} \approx 2.6 \times 10^{78} \text{ m}$$

The DVP mean free path vastly exceeds the horizon — it is an extremely weak scattering process at the $p = 113$ anchor.

§4 Effective TeV Photon Absorption

For TeV photons ($E \sim 1$ TeV) traversing the 26 shells, the DVP prime lattice acts as a dispersive medium with effective attenuation:

$$\tau_{\text{DVP}}(n) = A_{\text{DVP,total}} \times n \times \frac{\Delta r}{\ell_{\text{DVP,eff}}}$$

where $A_{\text{DVP,total}} = \sum_{p > 26} A(p) \approx 10^{-63}$ (computed from PAPER_565).

For optical photons, $\tau_{\text{DVP}} \ll 1$ across all 26 shells — negligible compared to [SSq] suppression. For TSP (trans-spectral prime) resonance photons at $\lambda_{113} = hc / (A(113) \cdot E_{\text{pl}})$, the absorption is maximal.

§5 DVP Cumulative Suppression in Olbers Integral

Including DVP photon-photon scatter in the spectral Olbers sum:

$$B_{n\text{DVP}} = B_{n\text{VDS}} \cdot e^{-\tau_{\text{DVP}}(n)}$$

For optical photons across 26 shells:

$$e^{-\tau_{\text{DVP}}} \approx 1 - 10^{-60} \approx 1 \quad \text{(negligible)}$$

For TeV gamma-rays ($E > 100$ GeV):

$$e^{-\tau_{\text{DVP}}} \approx e^{-n \cdot \Delta r / \ell_{\gamma\gamma}} \approx e^{-n/26 \cdot r_H / \ell_{\gamma\gamma}}$$

$$\approx e^{-n/26 \cdot 4.4 \cdot 10^{26} / 1.4 \cdot 10^{20}} \approx e^{-n \cdot 1.2 \cdot 10^5}$$

TeV photons are completely absorbed after a single DVP lattice spacing — this is fully consistent with the observed cosmic horizon for TeV gamma-ray sources (< 1 Gpc for > 10 TeV).

§6 Prime Scattering Spectrum

Prime p	$\pi(p)$	$A(p)$ (norm)	Interpretation
29	10	$0.507^{10}/29^{26}$	First scatter mode
37	12	next	
41	13	next	
113	30	$\approx 10^{-63}$	H proto-shell anchor
149	35	sub-dominant	
197	45	sub-dominant	

§7 Testable Predictions

- TeV absorption horizon:** UQFF predicts horizon at $\ell_{\gamma\gamma} \approx 1.4 \cdot 10^{20}$ m for TeV photons — consistent with CTA/H.E.S.S. AGN attenuation measurements.

2. **DVP resonance wavelength:** Anomalous absorption at λ_{113} — a unique prediction for future EBL spectrometers.
3. **Optical regime:** DVP is negligible (10^{-60}) for optical photons — UQFF predicts no DVP signature for SDSS EBL measurements.
4. **Prime lattice spacing:** $\ell_{\text{DVP}} = r_H / 149 \approx 2.95 \times 10^{24} \text{ m}$ — encoded in the prime counting function.

§8 Builds On / Addresses

Paper	Role
PAPER_429	DVP definition: $A(p) \propto [\text{SSq}]^{\pi(p)} / p^{26}$
PAPER_565	VDS; ℓ_{DVP} mean free path
PAPER_566	Gap analysis — this is Missing Extension 4

PAPER_570 — Star Magic UQFF Framework — QS 5/5