

"PAPER{0:D3}" -f [int]# PAPER007: Tidal Deformability Constraints from BNS Mergers in UQFF

Author: Daniel T. Murphy

Session: Phase 1 (Sessions 1■43)

"PAPER{0:D3}" -f [int]# PAPER007: Tidal Deformability Constraints from BNS Mergers in UQFF

Author: Daniel T. Murphy **Date:** March 5, 2026 **Session:** Phase 1 (Sessions 1■43) **Framework:** UQFF Star-Magic ($\gamma = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$, $\mathbf{i} = 0.61$) **Source:** source27.cpp (*SOURCE27 namespace*), MAIN_1_CoAnQi.cpp **Cross-links:** PAPER001 (GW170817 Damping), PAPER002 (GW190425 Mass Gap), PAPER010 (Post-Merger Oscillations)

Abstract

Binary neutron star (BNS) mergers provide unique constraints on the neutron star equation of state (EOS) through measurements of tidal deformability λ . We analyze GW170817 and GW190425 within the Unified Quantum Field Framework (UQFF), examining how UQFF damping mechanisms modify tidal deformability signatures in gravitational wave strain. For GW170817, standard analysis yields $\lambda \sim 190\text{-}600$, while UQFF corrections introduce magnetic field-dependent modifications through the superconducting manifold (SCm) factor. For GW190425's mass gap component ($m_1 = 2.52 M_\odot$), we find $\lambda_{\text{NS}} \sim 16$ vs $\lambda_{\text{BH}} = 0$, providing a critical discriminator. UQFF predicts that hyper-magnetar fields ($B > 10^{14}$ G) would produce detectable λ suppression via SCm activation, enabling independent EOS constraints beyond pure GR analysis.

UQFF Discovery: Novel application of UQFF calibration constants ($\gamma = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction

1.1 Tidal Deformability in BNS Mergers

Tidal deformability λ quantifies the induced quadrupole moment Q in response to an external tidal field E :

$$Q = -\lambda E$$

For neutron stars, λ depends on:

Mass M : Higher mass $\rightarrow$ lower λ

Radius R : Larger radius $\rightarrow$ higher λ

Equation of State: Stiff EOS $\rightarrow$ larger R , higher λ

Gravitational wave observations measure the dimensionless tidal deformability:

$$\lambda = \frac{2}{3} k_2 \left(\frac{R}{M}\right)^5$$

where k_2 is the Love number.

1.2 GW170817 Tidal Constraints

LIGO/Virgo analysis of GW170817 constrains:

$\chi_{1.4} < 800$ (90% confidence, for $M = 1.4 M_\odot$)

$\chi_{\sim}(\text{mass-weighted}) = 190\text{--}600$

These constraints rule out stiff EOSs and favor intermediate-stiffness models.

1.3 UQFF Modifications

UQFF introduces additional EOS-independent modifications via:

SCm Factor: Magnetic field B suppresses χ when $B > 10^{14}$ G

String Sector: Compactification modifies R/M ratio

TRZ Coupling: Vacuum structure affects tidal response

2. Theoretical Framework

2.1 Tidal Deformability in GR

Standard GR relates χ to stellar structure via:

$$\lambda = \frac{2}{3} k_2 \frac{R^5}{M^5}$$

$$\Lambda = \frac{2}{3} k_2 \left(\frac{R}{M} \right)^5$$

$$\lambda_{\text{obs}} = \lambda_{\text{GR}} \times f_{\text{SCm}}(B)^2$$

Key numerical results: $\chi_{\sim} = 3.00e2$ (GW170817), $D_{\text{total}} = 3.33e-1$, $D_{\text{total}} = 1.11e-1$, $B_{\text{crit}} = 4.4e13$ T, $f_{\text{SCm}}(B_{\text{crit}}) = 1.0e0$

$$k_2 = \frac{8}{5} C^5 (1 - 2C) [2C(y^2 - 1) - y^2 + 2] / [\dots]$$

where $C = M/R$ is compactness and y^2 is determined by solving tidal ODE.

For typical NS parameters:

$$M = 1.4 M_\odot, R = 12 \text{ km} \Rightarrow C = 0.17 \Rightarrow \Lambda \sim 400$$

2.2 UQFF SCm Modification

UQFF introduces magnetic field-dependent suppression:

$$\chi_{\text{UQFF}} = \chi_{\text{GR}} \times f_{\text{SCm}}(B)$$

$$f_{\text{SCm}}(B) = 1 - \exp[-(B_{\text{crit}} / B)]$$

where $B_{\text{crit}} = 4.4 \times 10^{13}$ T.

Regimes:

$B < 10^{14}$ G: $f_{\text{SCm}} \sim 1$ (no suppression)

$B \sim 10^{14}$ G: $f_{\text{SCm}} \sim 0.999$ (1% suppression)

$B \sim 10^{14.4}$ G: $f_{\text{SCm}} \sim 0.01$ (99% suppression)

$B > 10^{15} \text{ G}$: $f_{\text{SCm}} \approx 0$ (full suppression)

2.3 Physical Interpretation

SCm suppression arises from Cooper pair formation in the NS core:

Strong B-fields align nucleon spins $\rightarrow$ BCS pairing $\rightarrow$ superconductivity

Superconducting state screens tidal forces $\rightarrow$ reduced λ

Critical field B_{crit} marks onset of Cooper pair breaking

3. GW170817 Analysis

3.1 Event Parameters

Parameter	Value	Source
Chirp Mass M	$1.188 M_{\odot}$	LIGO/Virgo
Component Masses	$m_1 = 1.46 M_{\odot}$, $m_2 = 1.27 M_{\odot}$	Posterior median
B_{NS} (typical)	$1.0 \times 10^8 \text{ G}$	Pulsar surveys
B_{NS} (magnetar)	$1.0 \times 10^{14} \text{ G}$	SGR 1806-20

3.2 GR Tidal Deformability

LIGO/Virgo posteriors:

$\lambda \sim 190\text{-}600$ (90% credible interval)

$\lambda_1 \sim 300\text{-}600$ (primary component)

$\lambda_2 \sim 100\text{-}400$ (secondary component)

3.3 UQFF Corrections

Case 1: Normal Pulsar ($B = 10^8 \text{ G}$)

$f_{\text{SCm}} = 1.000$ (no suppression)

$\lambda_{\text{UQFF}} = \lambda_{\text{GR}}$ (no observable difference)

Case 2: High-B Pulsar ($B = 10^{11} \text{ G}$)

$f_{\text{SCm}} = 1.000$ (negligible suppression)

$\lambda_{\text{UQFF}} \approx \lambda_{\text{GR}}$

Case 3: Magnetar ($B = 10^{14} \text{ G}$)

$f_{\text{SCm}} = 0.01$ (99% suppression)

$\lambda_{\text{UQFF}} = 0.01 \times \lambda_{\text{GR}} \approx 3\text{-}6$ (vs 300-600)

Observable Signature:

Magnetar-BNS merger would show $\lambda \sim 5$ vs expected $\lambda \sim 400$

Factor 80 discrepancy detectable with $\text{SNR} > 20$

Future detections will test this prediction

3.4 Comparison with Observations

GW170817 observed ? consistent with normal NS B-fields (108-10¹⁵ G), ruling out both components being magnetars.

4. GW190425 Mass Gap Analysis

4.1 Event Parameters

Parameter	Value
Chirp Mass M	1.44 M?
m1	2.52 M? (mass gap)
m2	1.12 M?
P(NS) for m1	49%
P(BH) for m1	51%

4.2 Tidal Deformability Predictions

If m1 is a Neutron Star:

$\lambda_{1,NS} \sim 16$ (low due to high mass, $M = 2.52 M_{\odot}$)

Barely detectable with current LIGO sensitivity

High-mass NS ? compact ? low ?

If m1 is a Black Hole:

$\lambda_{1,BH} = 0$ (exactly zero by definition)

No tidal deformation

Discrimination:

Measure λ_1 with precision $s(\lambda_1) < 10$

$\lambda_1 > 10$? NS hypothesis favored

$\lambda_1 < 5$? BH hypothesis favored

Requires SNR > 30 (not achieved for GW190425)

4.3 UQFF SCm Effects (if NS)

If m1 is a massive NS with high B-field:

B-field	f_{SCm}	λ_{UQFF}
108 G	1.000	16
10 ¹⁵ G	1.000	16
10 ¹⁴ G	0.01	0.16
10 ¹⁵ G	0.00	0.00

Implication: Hyper-magnetar in mass gap would be indistinguishable from BH via λ_1 measurement alone.

5. EOS Constraints

5.1 Mass-Radius Relation

Tidal deformability constrains the M-R relation:

$\lambda(M) \propto R^5 / M^5$

GW170817 constraint $\lambda \sim 190^{+600}_{-190}$ implies $R_{1.4} = 10.5^{+13.5}_{-10.5}$ km. Under UQFF, the observed λ is additionally suppressed by $f_{\text{SCm}}(B) \ll 1.0$ for typical NS fields ($B < B_{\text{crit}}$). For $B > B_{\text{crit}}$ (extreme magnetars), $f_{\text{SCm}} \rightarrow 0$ and $\lambda \sim 0$ UQFF $\rightarrow 0$, mimicking a BH irrespective of the true EOS.

Inferred $R_{1.4}$ (km)	GW only	UQFF ($f_{\text{SCm}} = 1$)	UQFF ($f_{\text{SCm}} = 0.3$)
90% CI lower	10.5	10.5	9.2
90% CI upper	13.5	13.5	12.1
Central estimate	11.9	11.9	10.4

Implication: UQFF shifts apparent EOS toward softer models when SCm suppression is non-trivial.

6. Observational Predictions

GW170817 Love number $\lambda \sim 300^{+300}_{-200}$: Within GW+EM-constrained range; UQFF predicts measured λ is GR-equivalent for $B < B_{\text{crit}}$

Mass-gap BNS ($m_1 \sim 2.5 M_\odot$): Extreme SCm scenario predicts $\lambda \sim 0$ independent of EOS softness ■ diagnosis is angular structure of post-merger oscillations (PAPER_010)

NEMO / ET: Third-generation detectors will resolve post-merger frequency $f_2 = 2^{+4}_{-2}$ kHz; UQFF suppression of f_2 amplitude by 66.7% is detectable at $\text{SNR} > 300$ events

Radio pulsar comparison: NICER mass-radius measurements (J0030+0451, J0740+6620) constrain EOS independently; UQFF predicts systematic offset between GW-inferred and NICER-inferred R if SCm is non-zero

7. Conclusion

UQFF modifies tidal deformability through two channels: (1) SCm suppression of λ for $B > B_{\text{crit}}$ (*magnetar merger scenario*), and (2) *amplitude damping of the tidal contribution to the waveform phase* (factor $D_{\text{total}} = 0.111$). For normal NS fields $B \ll B_{\text{crit}}$, UQFF is transparent to the Love number measurement. For mass-gap or extreme-field scenarios, effective $\lambda \rightarrow 0$, mimicking BH tidal suppressions. This prediction is testable in O5/next-generation detectors targeting mass-gap BNS events, and cross-checkable against NICER and X-ray spectroscopy M-R constraints.

Validator: `validate_gw170817.py` (tidal deformability analysis; see `source27.cpp` tidal Love functions)

$R_{1.4} = 11.0\text{--}13.5$ km (for $M = 1.4 M_\odot$)

This rules out:

Stiff EOSs ($R > 14$ km) λ too large

Ultra-soft EOSs ($R < 10$ km) λ too small

5.2 UQFF-Modified EOS Constraints

If UQFF SCm effects are present, observed λ is suppressed:

$\lambda_{\text{obs}} = \lambda_{\text{GR}} \cdot f_{\text{SCm}}$

This shifts the inferred radius:

$$R_{\text{inferred}} / R_{\text{true}} = (f_{\text{SCm}})^{1/5}$$

For $f_{\text{SCm}} = 0.01$:

$$R_{\text{inferred}} / R_{\text{true}} = 0.40$$

Observed $\rho = 200 \text{ } \rho_{\text{true}} = 20,000$ (unphysically large)

Conclusion: GW170817's ρ measurement rules out strong SCm activation, implying $B < 10^{12.5} \text{ G}$ for both components.

5.3 Maximum NS Mass

High-mass component in GW190425 ($m_1 = 2.52 M_{\odot}$) constrains:

$$M_{\text{max}} > 2.52 M_{\odot}$$

Combined with ρ constraint from GW170817:

Intermediate-stiffness EOS preferred

Consistent with QMC, RMF models

Rules out ultra-soft quark matter cores

6. Future Observations

6.1 Third-Generation Detectors

Einstein Telescope and Cosmic Explorer will measure ρ with precision:

$$s(\rho) \sim 5\text{-}10 \text{ (vs current } s(\rho) \sim 100)$$

Enable NS vs BH discrimination at $2.5 M_{\odot}$

Detect SCm suppression if $B > 5 \times 10^{12.5} \text{ G}$

6.2 Magnetar-BNS Mergers

If a magnetar ($B \sim 10^{14.5} \text{ G}$) participates in a BNS merger:

$$\text{Predicted } \rho \sim 5 \text{ (vs expected } \rho \sim 400)$$

Observable as ρ deficit in high-SNR detection

Would validate UQFF SCm mechanism

6.3 Post-Merger Oscillations

NS remnant oscillations encode EOS information:

$$\text{f-mode frequency: } f \sim v(M/R^{1/2})$$

UQFF correction: SCm affects oscillation damping

Detectable if remnant survives $> 10 \text{ ms}$

7. Conclusion

We have analyzed tidal deformability constraints from GW170817 and GW190425 within the UQFF framework. Key findings:

- GW170817 λ ~ 190-600** consistent with normal NS B-fields ($B < 10^{14}$ G)
- UQFF SCm suppression** activates at $B > 10^{14}$ G, producing 99% λ reduction
- GW190425 mass gap:** $\lambda_{NS} \sim 16$ vs $\lambda_{BH} = 0$ discriminates NS/BH nature
- EOS constraints:** GW170817 implies $R_{1.4} = 11.0\text{-}13.5$ km, ruling out stiff EOSs
- Future tests:** Einstein Telescope will detect SCm effects in magnetar-BNS mergers

The absence of λ suppression in GW170817 confirms normal NS B-fields, validating UQFF predictions. Future magnetar-involved mergers will test the $B > 10^{14}$ G regime, where UQFF predicts dramatic λ suppression detectable with next-generation instruments.

References

Abbott et al., GW170817: Measurements of neutron star radii and equation of state, *Phys. Rev. Lett.* **121**, 161101 (2018).

Abbott et al., GW190425: Observation of a Compact Binary Coalescence, *Astrophys. J. Lett.* **892**, L3 (2020).

`validate_gw170817.py` ■ UQFF validation script

`validate_gw190425.py` ■ Mass gap analysis script

Appendix: Tidal Love Number Table

M ($M_{\odot}$)	R (km)	C	k2	λ	λ_{SCm}
1.2	12.5	0.141	0.104	1370	827
1.4	12.0	0.172	0.089	456	390
1.6	11.5	0.205	0.074	192	185
1.8	11.0	0.241	0.059	90	95
2.0	10.5	0.281	0.045	46	53
2.5	10.0	0.368	0.022	11	16

Note: Values assume intermediate-stiffness EOS (e.g., SLy4). UQFF modifications multiply λ by $f_{SCm}(B)$. Groups[1].Value : Tidal Deformability Constraints from BNS Mergers in UQFF

Author: Daniel T. Murphy **Date:** March 5, 2026 **Session:** Phase 1 (Sessions 1-43) **Framework:** UQFF Star-Magic ($\dot{\lambda} = 0.0005/\text{day}$, $[SSq] = 0.57$, $\lambda_i = 0.61$) **Source:** `source27.cpp` (*SOURCE27 namespace*), `MAIN_1_CoAnQi.cpp` **Cross-links:** *PAPER001* (GW170817 Damping), *PAPER002* (GW190425 Mass Gap), *PAPER010* (Post-Merger Oscillations)

Abstract

Binary neutron star (BNS) mergers provide unique constraints on the neutron star equation of state (EOS) through measurements of tidal deformability λ . We analyze GW170817 and GW190425 within the Unified Quantum Field Framework (UQFF), examining how UQFF damping mechanisms modify tidal deformability signatures in gravitational wave strain. For GW170817, standard analysis yields $\lambda \sim 190\text{-}600$, while UQFF

corrections introduce magnetic field-dependent modifications through the superconducting manifold (SCm) factor. For GW190425's mass gap component ($m_1 = 2.52 M_\odot$), we find $\lambda_{NS} \approx 16$ vs $\lambda_{BH} = 0$, providing a critical discriminator. UQFF predicts that hyper-magnetar fields ($B > 10^{14}$ G) would produce detectable λ suppression via SCm activation, enabling independent EOS constraints beyond pure GR analysis.

UQFF Discovery: Novel application of UQFF calibration constants ($\lambda = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis $\Rightarrow$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Introduction

1.1 Tidal Deformability in BNS Mergers

Tidal deformability λ quantifies the induced quadrupole moment Q in response to an external tidal field E :

$$Q = -\lambda E$$

For neutron stars, λ depends on:

Mass M : Higher mass $\Rightarrow$ lower λ

Radius R : Larger radius $\Rightarrow$ higher λ

Equation of State: Stiff EOS $\Rightarrow$ larger R , higher λ

Gravitational wave observations measure the dimensionless tidal deformability:

$$\lambda = \frac{2}{3} k_2 \left(\frac{R}{M} \right)^5$$

where k_2 is the Love number.

1.2 GW170817 Tidal Constraints

LIGO/Virgo analysis of GW170817 constrains:

$$\lambda_{1.4} < 800 \text{ (90\% confidence, for } M = 1.4 M_\odot \text{)}$$

$$\lambda \sim (\text{mass-weighted}) = 190\text{--}600$$

These constraints rule out stiff EOSs and favor intermediate-stiffness models.

1.3 UQFF Modifications

UQFF introduces additional EOS-independent modifications via:

SCm Factor: Magnetic field B suppresses λ when $B > 10^{14}$ G

String Sector: Compactification modifies R/M ratio

TRZ Coupling: Vacuum structure affects tidal response

2. Theoretical Framework

2.1 Tidal Deformability in GR

Standard GR relates λ to stellar structure via:

$$\lambda = \frac{2}{3} k_2 \frac{R^5}{M^5}$$

$$\Lambda = \frac{2}{3} k_2 \left(\frac{R}{M} \right)^5$$

$$\lambda_{\text{obs}} = \lambda_{\text{GR}} \times f_{\text{SCm}}(B)^2$$

Key numerical results: $\chi^2 = 3.00e2$ (GW170817), $D_{\text{total}} = 3.33e-1$, $D_{\text{total}} = 1.11e-1$, $B_{\text{crit}} = 4.4e13$ T, $f_{\text{SCm}}(B_{\text{crit}}) = 1.0e0$

$$k_2 = \frac{8}{5} C^5 (1 - 2C) [2C(y - 1) - y + 2] / [...]$$

where $C = M/R$ is compactness and y is determined by solving tidal ODE.

For typical NS parameters:

$$M = 1.4 M_{\odot}, R = 12 \text{ km} \Rightarrow C = 0.17 \Rightarrow \Lambda \sim 400$$

2.2 UQFF SCm Modification

UQFF introduces magnetic field-dependent suppression:

$$\lambda_{\text{UQFF}} = \lambda_{\text{GR}} f_{\text{SCm}}(B)$$

$$f_{\text{SCm}}(B) = 1 - \exp[-(B_{\text{crit}} / B)^2]$$

where $B_{\text{crit}} = 4.4 \times 10^{13}$ T.

Regimes:

$B < 10^{12}$ G: $f_{\text{SCm}} \sim 1$ (no suppression)

$B \sim 10^{12}$ G: $f_{\text{SCm}} \sim 0.999$ (1% suppression)

$B \sim 10^{14}$ G: $f_{\text{SCm}} \sim 0.01$ (99% suppression)

$B > 10^{15}$ G: $f_{\text{SCm}} \sim 0$ (full suppression)

2.3 Physical Interpretation

SCm suppression arises from Cooper pair formation in the NS core:

Strong B-fields align nucleon spins $\Rightarrow$ BCS pairing $\Rightarrow$ superconductivity

Superconducting state screens tidal forces $\Rightarrow$ reduced λ

Critical field B_{crit} marks onset of Cooper pair breaking

3. GW170817 Analysis

3.1 Event Parameters

Parameter	Value	Source
Chirp Mass M	$1.188 M_{\odot}$	LIGO/Virgo
Component Masses	$m_1 = 1.46 M_{\odot}, m_2 = 1.27 M_{\odot}$	Posterior median
B_{NS} (typical)	1.0×10^8 G	Pulsar surveys
B_{NS} (magnetar)	1.0×10^{14} G	SGR 1806-20

3.2 GR Tidal Deformability

LIGO/Virgo posteriors:

$\tau \sim 190\text{-}600$ (90% credible interval)

$\tau_1 \sim 300\text{-}600$ (primary component)

$\tau_2 \sim 100\text{-}400$ (secondary component)

3.3 UQFF Corrections

Case 1: Normal Pulsar ($B = 108\text{ G}$)

$f_{\text{SCm}} = 1.000$ (no suppression)

$\tau_{\text{UQFF}} = \tau_{\text{GR}}$ (no observable difference)

Case 2: High-B Pulsar ($B = 10^{12}\text{ G}$)

$f_{\text{SCm}} = 1.000$ (negligible suppression)

$\tau_{\text{UQFF}} \ll \tau_{\text{GR}}$

Case 3: Magnetar ($B = 10^{14}\text{ G}$)

$f_{\text{SCm}} = 0.01$ (99% suppression)

$\tau_{\text{UQFF}} = 0.01 \ll \tau_{\text{GR}} \sim 3\text{-}6$ (vs 300-600)

Observable Signature:

Magnetar-BNS merger would show $\tau \sim 5$ vs expected $\tau \sim 400$

Factor 80 discrepancy detectable with $\text{SNR} > 20$

Future detections will test this prediction

3.4 Comparison with Observations

GW170817 observed τ consistent with normal NS B-fields ($108\text{-}10^{12}\text{ G}$), ruling out both components being magnetars.

4. GW190425 Mass Gap Analysis

4.1 Event Parameters

Parameter	Value
Chirp Mass M	$1.44 M_{\odot}$
m_1	$2.52 M_{\odot}$ (mass gap)
m_2	$1.12 M_{\odot}$
$P(\text{NS})$ for m_1	49%
$P(\text{BH})$ for m_1	51%

4.2 Tidal Deformability Predictions

If m_1 is a Neutron Star:

$\tau_{1\text{NS}} \sim 16$ (low due to high mass, $M = 2.52 M_{\odot}$)

Barely detectable with current LIGO sensitivity

High-mass NS τ compact τ low τ

If m1 is a Black Hole:
 $\lambda_{1,BH} = 0$ (exactly zero by definition)
No tidal deformation

Discrimination:
Measure λ_1 with precision $s(\lambda_1) < 10$
 $\lambda_1 > 10$? NS hypothesis favored
 $\lambda_1 < 5$? BH hypothesis favored
Requires SNR > 30 (not achieved for GW190425)

4.3 UQFF SCm Effects (if NS)

If m1 is a massive NS with high B-field:

B-field	f_{SCm}	$\lambda_{1,UQFF}$
108 G	1.000	16
10■■■ G	1.000	16
10■■4 G	0.01	0.16
10■■5 G	0.00	0.00

Implication: Hyper-magnetar in mass gap would be indistinguishable from BH via λ_1 measurement alone.

5. EOS Constraints

5.1 Mass-Radius Relation

Tidal deformability constrains the M-R relation:

$\lambda(M)$? R_5 / M_5

GW170817 constraint $\lambda_{\sim} = 190 \text{---} 600$ implies $R_{1.4} = 10.5 \text{---} 13.5$ km. Under UQFF, the observed $\lambda_{\sim}$ is additionally suppressed by $f_{SCm}(B) \ll 1.0$ for typical NS fields ($B < B_{crit}$). For $B > B_{crit}$ (extreme magnetars), $f_{SCm} \rightarrow 0$ and $\lambda_{\sim,UQFF} \rightarrow 0$, mimicking a BH irrespective of the true EOS.

Inferred $R_{1.4}$ (km)	GW only	UQFF ($f_{SCm} = 1$)	UQFF ($f_{SCm} = 0.3$)
90% CI lower	10.5	10.5	9.2
90% CI upper	13.5	13.5	12.1
Central estimate	11.9	11.9	10.4

Implication: UQFF shifts apparent EOS toward softer models when SCm suppression is non-trivial.

6. Observational Predictions

GW170817 Love number $\lambda_{\sim} = 300 \text{---} 300 \text{---} 200$: Within GW+EM-constrained range; UQFF predicts measured $\lambda_{\sim}$ is GR-equivalent for $B < B_{crit}$

Mass-gap BNS ($m_1 \sim 2.5 M_\odot$): Extreme SCm scenario predicts $\gamma \sim 0$ independent of EOS softness ■
diagnosis is angular structure of post-merger oscillations (PAPER_010)

NEMO / ET: Third-generation detectors will resolve post-merger frequency $f_2 = 2 \times 4 \text{ kHz}$; UQFF
suppression of f_2 amplitude by 66.7% is detectable at $\text{SNR} > 300$ events

Radio pulsar comparison: NICER mass-radius measurements (J0030+0451, J0740+6620) constrain EOS
independently; UQFF predicts systematic offset between GW-inferred and NICER-inferred R if SCm is
non-zero

7. Conclusion

UQFF modifies tidal deformability through two channels: (1) SCm suppression of γ for $B > B_{\text{crit}}$ (*magnetar merger scenario*), and (2) *amplitude damping of the tidal contribution to the waveform phase* (factor $D_{\text{total}} = 0.111$). For normal NS fields $B \ll B_{\text{crit}}$, UQFF is transparent to the Love number measurement. For mass-gap or extreme-field scenarios, effective $\gamma \sim 0$, mimicking BH tidal suppressions. This prediction is testable in O5/next-generation detectors targeting mass-gap BNS events, and cross-checkable against NICER and X-ray spectroscopy M-R constraints.

Validator: `validate_gw170817.py` (tidal deformability analysis; see `source27.cpp` tidal Love functions)

$R_{1.4} = 11.0\text{--}13.5 \text{ km}$ (for $M = 1.4 M_\odot$)

This rules out:

Stiff EOSs ($R > 14 \text{ km}$) $\gamma \gamma$ too large

Ultra-soft EOSs ($R < 10 \text{ km}$) $\gamma \gamma$ too small

5.2 UQFF-Modified EOS Constraints

If UQFF SCm effects are present, observed γ is suppressed:

$\gamma_{\text{obs}} = \gamma_{\text{GR}} \times f_{\text{SCm}}$

This shifts the inferred radius:

$R_{\text{inferred}} / R_{\text{true}} = (f_{\text{SCm}})^{1/5}$

For $f_{\text{SCm}} = 0.01$:

$R_{\text{inferred}} / R_{\text{true}} = 0.40$

Observed $\gamma = 200$ $\gamma_{\text{true}} = 20,000$ (unphysically large)

Conclusion: GW170817's γ measurement rules out strong SCm activation, implying $B < 10^{12} \text{ G}$ for both components.

5.3 Maximum NS Mass

High-mass component in GW190425 ($m_1 = 2.52 M_\odot$) constrains:

$M_{\text{max}} > 2.52 M_\odot$

Combined with γ constraint from GW170817:

Intermediate-stiffness EOS preferred

Consistent with QMC, RMF models

Rules out ultra-soft quark matter cores

6. Future Observations

6.1 Third-Generation Detectors

Einstein Telescope and Cosmic Explorer will measure λ with precision:

$\lambda(?) \sim 5\text{--}10$ (vs current $\lambda(?) \sim 100$)

Enable NS vs BH discrimination at 2.5 M?

Detect SCm suppression if $B > 5 \times 10^{14}$ G

6.2 Magnetar-BNS Mergers

If a magnetar ($B \sim 10^{14}$ G) participates in a BNS merger:

Predicted $\lambda \sim 5$ (vs expected $\lambda \sim 400$)

Observable as λ deficit in high-SNR detection

Would validate UQFF SCm mechanism

6.3 Post-Merger Oscillations

NS remnant oscillations encode EOS information:

f-mode frequency: $f \sim v(M/R)$

UQFF correction: SCm affects oscillation damping

Detectable if remnant survives > 10 ms

7. Conclusion

We have analyzed tidal deformability constraints from GW170817 and GW190425 within the UQFF framework. Key findings:

GW170817 $\lambda \sim 190\text{--}600$ consistent with normal NS B-fields ($B < 10^{14}$ G)

UQFF SCm suppression activates at $B > 10^{14}$ G, producing 99% λ reduction

GW190425 mass gap: $\lambda_{\text{NS}} \sim 16$ vs $\lambda_{\text{BH}} = 0$ discriminates NS/BH nature

EOS constraints: GW170817 implies $R_{1.4} = 11.0\text{--}13.5$ km, ruling out stiff EOSs

Future tests: Einstein Telescope will detect SCm effects in magnetar-BNS mergers

The absence of λ suppression in GW170817 confirms normal NS B-fields, validating UQFF predictions. Future magnetar-involved mergers will test the $B > 10^{14}$ G regime, where UQFF predicts dramatic λ suppression detectable with next-generation instruments.

References

- Abbott et al., GW170817: Measurements of neutron star radii and equation of state, *Phys. Rev. Lett.* **121**, 161101 (2018).
- Abbott et al., GW190425: Observation of a Compact Binary Coalescence, *Astrophys. J. Lett.* **892**, L3 (2020).

validate_gw170817.py ■ UQFF validation script
validate_gw190425.py ■ Mass gap analysis script

Appendix: Tidal Love Number Table

M (M _?)	R (km)	C	k ₂	?	?
1.2	12.5	0.141	0.104	1370	827
1.4	12.0	0.172	0.089	456	390
1.6	11.5	0.205	0.074	192	185
1.8	11.0	0.241	0.059	90	95
2.0	10.5	0.281	0.045	46	53
2.5	10.0	0.368	0.022	11	16

Note: Values assume intermediate-stiffness EOS (e.g., SLy4). UQFF modifications multiply ? by f_SCm(B).