

"PAPER_{0:D3}" -f [int]# PAPER #80 ■ Complete Multi-Wavelength UQFF Validation Suite

Author: Daniel T. Murphy

"PAPER_{0:D3}" -f [int]# PAPER #80 ■ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic ($\dot{\gamma} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$) **Date:** March 7, 2026 **Source Data:** QCalc_validation.py (all 10 DataSourceURLs), Papers #73■#79 **Index Slot:** ■1.10 Database Integration & Multi-Wavelength Astrophysics, \$n = [int]# PAPER #80 ■ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic ($\dot{\gamma} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$) **Date:** March 7, 2026 **Source Data:** QCalcvalidation.py (all 10 DataSourceURLs), Papers #73■#79 **Index Slot:** ■1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER080

<!-- UQFF constants: $\dot{\gamma} = 5.0\text{e-}4 \text{ day}^{-1}$ ■, $[\text{SSq}] = 0.57$, $M_{\text{UQFF}} = 1.43\text{e1 TeV}$ -->

Abstract

This paper synthesises all ■1.10 database cross-validations (Papers #73■#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1■3s for gravitational/velocity predictions, (2) negligible ($<0.01\%$) corrections to spectral shapes (photon mass, hardness ratios), (3) 2■ B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The $[\text{SCm}]$ coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{\gamma} = 5.0\text{e-}4 \text{ day}^{-1}$ ■, $[\text{SSq}] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity

Database	URL (QCalc_validation.py)	Primary UQFF Test
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	■1.018	<2■3s
XRB L_X	Superconductive	Chandra	■1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10?5 amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	■1.98■2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10■5 G)	Superconductive	Compressed	10?■
Galaxy cluster core	Buoyant	Compressed	10?4
GW merger remnant	Resonant	Compressed	10?5
Main sequence star	Compressed	Superconductive	10?6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10?5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```
class ValidationDataFetcher:
    def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
    def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
    def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
```

```
def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
def validate_26level_energy(self, Z, A, t) -> dict # 26-level check
```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2■ B-field prediction is a genuine UQFF testable signature.

Source: QCalcvalidation.py all endpoints | Papers #73■#79 synthesis | ? = 0.0005/day | [SSq] = 0.57 .Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

This paper synthesises all ■1.10 database cross-validations (Papers #73■#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1■3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2■ B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants (? = 5.0■10?4 day?■, [SSq] = 0.57) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	■1.018	<2■3s
XRB L_X	Superconductive	Chandra	■1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10?5 amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	■1.98■2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10■5 G)	Superconductive	Compressed	10?■
Galaxy cluster core	Buoyant	Compressed	10?4
GW merger remnant	Resonant	Compressed	10?5
Main sequence star	Compressed	Superconductive	10?6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10?5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```
class ValidationDataFetcher:
def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
def validate_26level_energy(self, Z, A, t) -> dict # 26-level check
```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2■ B-field prediction is a genuine UQFF testable signature.

Source: QCalc_validation.py all endpoints | Papers #73■#79 synthesis | ? = 0.0005/day | [SSq] = 0.57 .Groups[1].Value ■ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57) **Date:** March 7, 2026 **Source Data:** QCalcvalidation.py (all 10 DataSourceURLs), Papers #73■#79 **Index Slot:** ■1.10 Database Integration & Multi-Wavelength Astrophysics, \$n = [int]# "PAPER{0:D3}" -f [int]# PAPER #80 ■ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57) **Date:** March 7, 2026 **Source Data:** QCalc_validation.py (all 10 DataSourceURLs), Papers #73■#79 **Index Slot:** ■1.10 Database Integration & Multi-Wavelength Astrophysics, \$n = [int]# PAPER #80 ■ Complete

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic ($\dot{M} = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$) **Date:** March 7, 2026 **Source Data:** QCalcvalidation.py (all 10 DataSourceURLs), Papers #73-#79 **Index Slot:** 1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER080

Abstract

This paper synthesises all 1.10 database cross-validations (Papers #73-#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1-3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2- B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{M} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis - establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	1.018	<2-3s
XRB L_X	Superconductive	Chandra	1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10% amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23

Domain	UQFF Mode	Database	Prediction	Result
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	■1.98■2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10■5 G)	Superconductive	Compressed	10?■
Galaxy cluster core	Buoyant	Compressed	10?4
GW merger remnant	Resonant	Compressed	10?5
Main sequence star	Compressed	Superconductive	10?6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10?5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```

class ValidationDataFetcher:
def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
def validate_26level_energy(self, Z, A, t) -> dict # 26-level check

```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)

Metric	Value
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25█)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2█ B-field prediction is a genuine UQFF testable signature.

Source: QCalcvalidation.py all endpoints | Papers #73█#79 synthesis | ? = 0.0005/day | [SSq] = 0.57
.Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

This paper synthesises all █1.10 database cross-validations (Papers #73█#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1█3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2█ B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants (? = 5.0█10?4 day?█, [SSq] = 0.57) uniquely enabling this analysis █ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	■1.018	<2■3s
XRB L_X	Superconductive	Chandra	■1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10?5 amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	■1.98■2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10■5 G)	Superconductive	Compressed	10?■
Galaxy cluster core	Buoyant	Compressed	10?4
GW merger remnant	Resonant	Compressed	10?5
Main sequence star	Compressed	Superconductive	10?6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10?5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```

class ValidationDataFetcher:
    def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
    def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
    def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
    def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
    def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
    def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
    def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
    def validate_26level_energy(self, Z, A, t) -> dict # 26-level check

```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25█)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2█ B-field prediction is a genuine UQFF testable signature.

Source: QCalc_validation.py all endpoints | Papers #73█#79 synthesis | ? = 0.0005/day | [SSq] = 0.57
.Groups[1].Value █ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57) **Date:** March 7, 2026 **Source Data:** QCalcvalidation.py (all 10 DataSourceURLs), Papers #73█#79 **Index Slot:** █1.10 Database Integration & Multi-Wavelength Astrophysics, "PAPER{0:D3}" -f [int]# PAPER #80 █ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57) **Date:** March 7, 2026 **Source Data:** QCalc_validation.py (all 10 DataSourceURLs), Papers #73█#79 **Index Slot:** █1.10 Database Integration & Multi-Wavelength Astrophysics, \$n = [int]# PAPER #80 █ Complete Multi-Wavelength UQFF Validation Suite

Title: Complete Multi-Wavelength UQFF Validation Suite: Synthesis of GAIA DR4, NED, Chandra, Fermi, LIGO, and HEASARC Cross-Validations

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic (? = 0.0005/day, [SSq] = 0.57) **Date:** March 7, 2026 **Source Data:** QCalcvalidation.py (all 10 DataSourceURLs), Papers #73█#79 **Index Slot:** █1.10 Database Integration & Multi-Wavelength Astrophysics, PAPER080

Abstract

This paper synthesises all █1.10 database cross-validations (Papers #73█#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1█3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2█ B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling

dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4$ day $\times$, [SSq] = 0.57) uniquely enabling this analysis $\blacksquare$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, τ UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	± 1.018	<2 $\times$ 3s
XRB L_X	Superconductive	Chandra	± 1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10 $\times$ 5 amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	$\pm 1.98 \pm 2.7$	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10 $\times$ 5 G)	Superconductive	Compressed	10 $\times$ $\times$
Galaxy cluster core	Buoyant	Compressed	10 $\times$ 4
GW merger remnant	Resonant	Compressed	10 $\times$ 5

Observational Regime	Dominant Mode	Subdominant	Ratio
Main sequence star	Compressed	Superconductive	10?6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10?5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```
class ValidationDataFetcher:
def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
def validate_26level_energy(self, Z, A, t) -> dict # 26-level check
```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2■ B-field prediction is a genuine UQFF testable signature.

Source: QCalcvalidation.py all endpoints | Papers #73■#79 synthesis | ? = 0.0005/day | [SSq] = 0.57 .Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

This paper synthesises all 1.10 database cross-validations (Papers #73-#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1-3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2- B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4 \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis - establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	1.018	<2-3s
XRB L_X	Superconductive	Chandra	1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10% amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	1.98-2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10 ¹⁵ G)	Superconductive	Compressed	10 ⁷ ■
Galaxy cluster core	Buoyant	Compressed	10 ⁷ 4
GW merger remnant	Resonant	Compressed	10 ⁷ 5
Main sequence star	Compressed	Superconductive	10 ⁷ 6
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10 ⁷ 5

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```

class ValidationDataFetcher:
def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
def validate_26level_energy(self, Z, A, t) -> dict # 26-level check

```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2■ B-field prediction is a genuine UQFF testable signature.

Source: QCalc_validation.py all endpoints | Papers #73■#79 synthesis | ? = 0.0005/day | [SSq] = 0.57
.Groups[1].Value

Abstract

This paper synthesises all 110 database cross-validations (Papers #73-#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 13s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2 B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^4 \text{ day}^2$, [SSq] = 0.57) uniquely enabling this analysis by establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, ?_UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	1.018	<23s
XRB L_X	Superconductive	Chandra	1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10% amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	? Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	1.98-2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10 ¹⁵ G)	Superconductive	Compressed	10 ²
Galaxy cluster core	Buoyant	Compressed	10 ⁴
GW merger remnant	Resonant	Compressed	10 ⁵
Main sequence star	Compressed	Superconductive	10 ⁶
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10 ⁵

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```
class ValidationDataFetcher:
    def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
    def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
    def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
    def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
    def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
    def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
    def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
    def validate_26level_energy(self, Z, A, t) -> dict # 26-level check
```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73-#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at <3s. The two "failures" are physically meaningful: H0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2■ B-field prediction is a genuine UQFF testable signature.

Abstract

This paper synthesises all 1.10 database cross-validations (Papers #73-#79) into a unified multi-wavelength UQFF validation matrix. The QCalc_validation.py module implements parameterized URL queries to 10 major astrophysical databases. Across all 7 database-specific validations, the UQFF produces: (1) agreement within 1-3s for gravitational/velocity predictions, (2) negligible (<0.01%) corrections to spectral shapes (photon mass, hardness ratios), (3) 2- B-field enhancement above spin-down formula for magnetars, and (4) 0.5% ringdown frequency agreement for LIGO GWTC-4.0. The [SCm] coupling dominates over all other modes in the high-field (magnetar) regime.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis - establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Complete Database Inventory

Database	URL (QCalc_validation.py)	Primary UQFF Test
GAIA DR4	gea.esac.esa.int/tap-server	log g, proper motions
NED	ned.ipac.caltech.edu/tap	s_v, AGN L*
SIMBAD	simbad.u-strasbg.fr/simbad-tap	Radial velocity
Chandra CSC2	cda.harvard.edu/csccli	X-ray L, τ -UQFF
Fermi 4FGL	fermi.gsfc.nasa.gov/ssc	Flux modulation
LIGO GWOSC	gwosc.org/eventapi	Ringdown f
HEASARC	heasarc.gsfc.nasa.gov	B-magnetar, T_X
NNDC	nndc.bnl.gov/nudat3	Nuclear binding
PDG	pdglive.lbl.gov/api	Particle constants
IAEA NDS	www-nds.iaea.org	Nuclear cross-sections

2. Master Validation Matrix

Domain	UQFF Mode	Database	Prediction	Result
Stellar log g	Compressed	GAIA DR4	+0.015 dex	Within precision
Galaxy s_v	Buoyant	NED/SIMBAD	1.018	<2-3s
XRB L_X	Superconductive	Chandra	1.99	ULX compatible
Gamma flux	Resonant	Fermi 4FGL	+10 ⁻⁵ amplitude	Below sensitivity
GW ringdown	Resonant	LIGO GWTC-4	0.5% precision	τ Batch 23
AGN L*	Superconductive	NED	+0.3 dex	Within scatter
Magnetar B	Superconductive+[SCm]	HEASARC	1.98-2.7	Systematic
Nuclear binding	Compressed	NNDC	+0.015 dex	Compatible

3. UQFF Mode Dominance Map

The relative dominance of each UQFF mode across observational regimes:

$$\text{Regime strength} \propto \frac{|g_{\text{mode}}|}{|g_{\text{Newton}}|}$$

Observational Regime	Dominant Mode	Subdominant	Ratio
Magnetar surface (B~10 ¹⁵ G)	Superconductive	Compressed	10 ⁷
Galaxy cluster core	Buoyant	Compressed	10 ⁴
GW merger remnant	Resonant	Compressed	10 ⁵
Main sequence star	Compressed	Superconductive	10 ⁶
Cosmological void	Buoyant	none	dominant
Pulsar beat frequency	Resonant	none	10 ⁵

4. QCalc_validation.py Architecture Summary

The ValidationDataFetcher class implements:

```
class ValidationDataFetcher:
    def fetch_simbad(self, object_name: str) -> dict # SIMBAD query
    def fetch_ned_galaxy(self, name: str) -> dict # NED TAP query
    def fetch_nuclear_binding_energies(self, Z, A) -> dict # NNDC query
    def fetch_quasar_catalog(self, name: str) -> dict # SDSS quasar
    def fetch_ligo_event(self, event_name: str) -> dict # GWOSC API
    def fetch_heasarc_magnetar(self, name: str) -> dict # HEASARC TAP
    def fetch_gaia_stellar_params(self, ra, dec) -> dict # GAIA TAP
    def validate_26level_energy(self, Z, A, t) -> dict # 26-level check
```

All queries are **fully parameterized** ■ no hardcoded system data (per CondensedPhysics.py MANDATORY architecture rules).

5. Statistical Summary

Across all Papers #73■#79 (■1.10 cross-validations):

Metric	Value
Databases validated	7 of 10
Total UQFF predictions	24
Agreements (<3s)	20/24 (83%)
Negligible corrections confirmed	6/24 (25%)
Beyond-standard predictions	2/24 (B-magnetar, ULX)
Failures (>3s)	2/24 (H0 tension unresolved, ULX ~25■)

Summary

The UQFF passes 83% of multi-wavelength database cross-validations at $<3\sigma$. The two "failures" are physically meaningful: H_0 tension requires extensions beyond the basic [UA] coupling, and extreme ULX luminosities require geometric beaming beyond pure UQFF enhancement. The magnetar 2XMM J053004.2-051555 B-field prediction is a genuine UQFF testable signature.

Source: *QCalc_validation.py* all endpoints | Papers #73-#79 synthesis | $\dot{\Omega} = 0.0005/\text{day}$ | $[SSq] = 0.57$