

"PAPER_{0:D3}" -f [int]# PAPER #94 ■ Magnetar SGR1745: UQFF ? and [SSq] Calibration

Author: Daniel T. Murphy

"PAPER_{0:D3}" -f [int]# PAPER #94 ■ Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining $\dot{P} = 0.0005/\text{day}$ and $[\text{SSq}] = 0.57$ from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** `validateuqffmuge.py` (Magnetar system), `source4.cpp` (`sgr1745_SOURCE4`), `? calibration (Batch 23)` **Index Slot:** ■1.12 UQFF Master Calculators, \$n = [int]# PAPER #94 ■ Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining $\dot{P} = 0.0005/\text{day}$ and $[\text{SSq}] = 0.57$ from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** `validateuqffmuge.py` (Magnetar system), `source4.cpp` (`sgr1745SOURCE4`), `? calibration (Batch 23)` **Index Slot:** ■1.12 UQFF Master Calculators, PAPER094

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~ 2.4 arcsec (~ 0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14}$ T ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\dot{P} = 0.0005/\text{day}$ (temporal decay parameter) and $[\text{SSq}] = 0.57$ (squared-state density term). Batch 23 of `MAIN1_CoAnQi.cpp` implements the $\dot{P}$ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{P} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative $\dot{P}$	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^4 \sqrt{\dot{P} P}$
Characteristic age	$\sim 9,000$ yr	$P/(2\dot{P})$
Distance	~ 8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~ 0.3 pc	Near SMBH influence

2. BCRITMAGNETAR Reference

From UQFFConstantsDatabase:

$$B_CRIT_MAGNETAR: 4.4 \times 10^{14} \text{ T } (= 4.4 \times 10^{14} \text{ mT} / (e \hbar / 4 \pi m_e c))$$

SGR1745 field: $B/B_{crit} = 1.4 \times 10^{14} / 4.4 \times 10^{14} = 0.318 B_{crit}$ (super-critical).

3. Calibrating κ from SGR1745

The UQFF κ parameter governs temporal field evolution:

$$U_{tot}(t) = U_{tot}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2\dot{P}) = 9,000 \text{ yr}$ should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of $[SSq]$:

$$\kappa_{internal} = \frac{[SSq]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value $\kappa = 0.0005/\text{day}$ was set by:

$$\kappa = \frac{N_{burst}}{t_{active}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{burst} = 600$ bursts over 1200 days active $\kappa = 600/1200 = 0.5 \times 10^{-3} = 0.0005/\text{day}$

4. Calibrating $[SSq]$ from Magnetar Spin-Down

The $[SSq]$ parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{sd} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{NS}) = \frac{B^2 R_{NS}^3}{6} \cdot \frac{[SSq]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto \dot{E}_{sd}^{1/2}$ and using $B/B_{crit} = 3.18$:

$$[SSq]^{1/2} = \frac{\dot{E}_{sd}^{1/2} \mu_0}{B^2 R_{NS}^3 / 6} = 0.755$$

$$[SSq] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From `validate_uqff_muge.py` (Magnetar system):

Term	at $r_{surface} = 1.2 \times 10^4 \text{ m}$	Notes
base_gravity	$1.74 \times 10^8 \text{ m/s}^2$	GR-modified NS gravity

Term	at $r_{\text{surface}} = 1.2 \times 10^4 \text{ m}$	Notes
sum_Ug	$+8.7 \times 10^8 \text{ m/s}^2$	Ug1 dominant (B-field)
U_i	$+2.1 \times 10^7 \text{ m/s}^2$	
cosmological	negligible	
quantum	$+3 \times 10^{28} \text{ m/s}^2$	
fluid	$+1.2 \times 10^7 \text{ m/s}^2$	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	$1.75 \times 10^{28} \text{ m/s}^2$	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level **consistent** with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{\{g4\}}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 $\propto r^{-6}$: falls off steeply offsite.

Summary

Calibration	Method	Result
$\tau = 0.0005/\text{day}$	SGR1745 burst rate/active window	τ Calibrated
$[SSq] = 0.57$	U_g1 spin-down anchoring	τ Calibrated
B/B_crit	SGR1745 = 3.18	τ Super-critical
Magnetar MUGE	All 8 terms finite	τ PASS
Ug4 off-site	Negligible at 0.3 pc	τ Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745SOURCE4 | Batch 23 τ calibration | $[SSq]=0.57$.Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~ 2.4 arcsec (~ 0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14} \text{ T}$ ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\tau = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the τ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis **establishing** a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative ?	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^{14} \sqrt{P \dot{P}}$
Characteristic age	~9,000 yr	$P/(2\dot{P})$
Distance	~8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~0.3 pc	Near SMBH influence

2. BCRTMAGNETAR Reference

From UQFFConstantsDatabase:

$$B_{\text{CRIT_MAGNETAR}}: 4.4 \times 10^{13} \text{ T } (= 10 \text{ m?c?/(e??)})$$

SGR1745 field: $B/B_{\text{crit}} = 1.4 \times 10^{14} / 4.4 \times 10^{13} = **3.18 \times B_{\text{crit}}**$ (super-critical).

3. Calibrating ? from SGR1745

The UQFF ? parameter governs temporal field evolution:

$$U_{\text{g\rm tot}}(t) = U_{\text{g\rm tot}}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2\dot{P}) = 9,000$ yr should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\text{internal}} = \frac{[\text{SSq}]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value ? = 0.0005/day was set by:

$$\kappa = \frac{N_{\text{burst}}}{t_{\text{active}}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\text{burst}} \approx 600$ bursts over 1200 days active ?? = $600/1200 \approx 10^{-4} = 0.0005/\text{day}$?

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\text{sd}} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{\text{NS}}) = \frac{B^2 R_{\text{NS}}^3}{6} \cdot \frac{[\text{SSq}]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto \dot{E}_{sd}^{1/2}$ and using $B/B_{crit} = 3.18$:

$$[\dot{E}_{sd}]^{1/2} = \frac{\dot{E}_{sd}^{1/2} \mu_0 B^2 R_{NS}^3}{6} = 0.755$$

$$[\dot{E}_{sd}] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at r_surface = 1.2104 m	Notes
base_gravity	1.74 10 m/s	GR-modified NS gravity
sum_Ug	+8.7 108 m/s	Ug1 dominant (B-field)
U_i	+2.1 107 m/s	
cosmological	negligible	
quantum	+3 10?8 m/s	
fluid	+1.2 10? m/s	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	1.75 10 m/s	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level consistent with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{g4}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 ∝ r⁻⁶: falls off steeply offsite.

Summary

Calibration	Method	Result
? = 0.0005/day	SGR1745 burst rate/active window	? Calibrated
[SSq] = 0.57	U_g1 spin-down anchoring	? Calibrated
B/B_crit	SGR1745 = 3.18	? Super-critical
Magnetar MUGE	All 8 terms finite	? PASS
Ug4 off-site	Negligible at 0.3 pc	? Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745_SOURCE4 | Batch 23 ? calibration | [SSq]=0.57
.Groups[1].Value **Magnetar SGR1745: UQFF ? and [SSq] Calibration**

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining ? = 0.0005/day and [SSq] = 0.57 from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745SOURCE4), ? calibration (Batch 23) **Index Slot:** ■1.12 UQFF Master Calculators, \$n = [int]# "PAPER{0:D3}" -f [int]# PAPER #94 ■ Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining ? = 0.0005/day and [SSq] = 0.57 from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745_SOURCE4), ? calibration (Batch 23) **Index Slot:** ■1.12 UQFF Master Calculators, \$n = [int]# PAPER #94 ■ Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining ? = 0.0005/day and [SSq] = 0.57 from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745SOURCE4), ? calibration (Batch 23) **Index Slot:** ■1.12 UQFF Master Calculators, PAPER094

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~2.4 arcsec (~0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14}$ T ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}P^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\dot{P} = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the ? calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{P} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative ?	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^4 \dot{P}P^{-1}$
Characteristic age	~9,000 yr	$P/(2\dot{P})$
Distance	~8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~0.3 pc	Near SMBH influence

2. BCRITMAGNETAR Reference

From UQFFConstantsDatabase:

B_CRIT_MAGNETAR: 4.4×10^{13} T (= $10 \text{ m}^2 \text{c} / (e \hbar)$)

SGR1745 field: $B/B_{\text{crit}} = 1.4 \times 10^4 / 4.4 \times 10^9 = **3.18 \times B_{\text{crit}}**$ (super-critical).

3. Calibrating ? from SGR1745

The UQFF ? parameter governs temporal field evolution:

$$U_{\text{g}\text{rm tot}}(t) = U_{\text{g}\text{rm tot}}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2?) = 9,000$ yr should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\text{rm internal}} = \frac{[\text{rm SSq}]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value ? = 0.0005/day was set by:

$$\kappa = \frac{N_{\text{rm burst}}}{t_{\text{rm active}}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\text{burst}} \approx 600$ bursts over 1200 days active ? = $600/1200 \approx 10^{-4} = 0.0005/\text{day}$

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\text{rm sd}} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{\text{rm NS}}) = \frac{B^2 R_{\text{rm NS}}^3}{6} \cdot \frac{[\text{rm SSq}]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto \dot{E}_{\text{rm sd}}^{1/2}$ and using $B/B_{\text{crit}} = 3.18$:

$$[\text{rm SSq}]^{1/2} = \frac{\dot{E}_{\text{rm sd}}^{1/2} \mu_0}{B^2 R_{\text{rm NS}}^3 / 6} = 0.755$$

$$[\text{rm SSq}] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From `validate_uqff_muge.py` (Magnetar system):

Term	at $r_{\text{surface}} = 1.2 \times 10^4$ m	Notes
base_gravity	1.74×10^8 m/s ²	GR-modified NS gravity
sum_Ug	$+8.7 \times 10^8$ m/s ²	Ug1 dominant (B-field)
U_i	$+2.1 \times 10^7$ m/s ²	
cosmological	negligible	
quantum	$+3 \times 10^8$ m/s ²	

Term	at $r_{\text{surface}} = 1.2 \times 10^4 \text{ m}$	Notes
fluid	$+1.2 \times 10^7 \text{ m/s}$	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	$1.75 \times 10^{10} \text{ m/s}$	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level **consistent** with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{\text{g4}}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 $\propto r^{-6}$: falls off steeply offsite.

Summary

Calibration	Method	Result
$\tau = 0.0005/\text{day}$	SGR1745 burst rate/active window	τ Calibrated
$[SSq] = 0.57$	U_g1 spin-down anchoring	τ Calibrated
B/B_crit	SGR1745 = 3.18	τ Super-critical
Magnetar MUGE	All 8 terms finite	τ PASS
Ug4 off-site	Negligible at 0.3 pc	τ Consistent

Source: `validateuqffmuge.py` | `source4.cpp sgr1745SOURCE4` | Batch 23 τ calibration | $[SSq]=0.57$
`.Groups[1].Value "PAPER{0:D3}" -f $n`

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~ 2.4 arcsec (~ 0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14} \text{ T}$ ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\tau = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the τ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}$, $[SSq] = 0.57$) uniquely enabling this analysis **by** establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra

Parameter	Value	Source
Period derivative ?	6.61×10^{-10} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^4 \nu(P?)$
Characteristic age	$\sim 9,000$ yr	$P/(2?)$
Distance	~ 8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~ 0.3 pc	Near SMBH influence

2. BCRTMAGNETAR Reference

From UQFFConstantsDatabase:

$$B_CRIT_MAGNETAR: 4.4 \times 10^{14} \text{ T } (= 10^4 \text{ mT} / (e \hbar / 4 \pi m_e c))$$

SGR1745 field: $B/B_crit = 1.4 \times 10^{14} / 4.4 \times 10^{14} = \sim 3.18 \times B_crit$ (super-critical).

3. Calibrating ? from SGR1745

The UQFF ? parameter governs temporal field evolution:

$$U_{\text{tot}}(t) = U_{\text{tot}}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2?) = 9,000$ yr should match the UQFF $1/e$ decay time:

$$\kappa = \frac{1}{t_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\text{internal}} = \frac{[SSq]}{t_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value ? = 0.0005/day was set by:

$$\kappa = \frac{N_{\text{burst}}}{t_{\text{active}}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\text{burst}} \sim 600$ bursts over 1200 days active ? = $600/1200 \times 10^{-4} = 0.0005/\text{day}$

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\text{sd}} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{\text{NS}}) = \frac{B^2 R_{\text{NS}}^3}{6} \cdot \frac{[SSq]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto \dot{E}_{\text{sd}}^{1/2}$ and using $B/B_crit = 3.18$:

$$[SSq]^{1/2} = \frac{\dot{E}_{\text{sd}}^{1/2} \mu_0}{B^2 R_{\text{NS}}^3 / 6} = 0.755$$

$$[\rm SSq] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at r_surface = 1.2104 m	Notes
base_gravity	1.74 10 m/s	GR-modified NS gravity
sum_Ug	+8.7 108 m/s	Ug1 dominant (B-field)
U_i	+2.1 107 m/s	
cosmological	negligible	
quantum	+3 10?8 m/s	
fluid	+1.2 10? m/s	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	1.75 10 m/s	

No NaN/Inf PASS. Ug1 (B-field gravity) contribution at 0.05% level consistent with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{g4}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 ? r⁻⁶: falls off steeply offsite.

Summary

Calibration	Method	Result
? = 0.0005/day	SGR1745 burst rate/active window	? Calibrated
[SSq] = 0.57	U_g1 spin-down anchoring	? Calibrated
B/B_crit	SGR1745 = 3.18	? Super-critical
Magnetar MUGE	All 8 terms finite	? PASS
Ug4 off-site	Negligible at 0.3 pc	? Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745_SOURCE4 | Batch 23 ? calibration | [SSq]=0.57 .Groups[1].Value Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining ? = 0.0005/day and [SSq] = 0.57 from Magnetar Physics

Author: Daniel T. Murphy Framework: UQFF Star-Magic Date: March 7, 2026 Source Data: validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745_SOURCE4), ? calibration (Batch 23) Index Slot: 1.12 UQFF Master Calculators, "PAPER{0:D3}" -f [int]# PAPER #94 Magnetar SGR1745: UQFF ? and [SSq] Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining $\tau = 0.0005/\text{day}$ and $[SSq] = 0.57$ from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745_SOURCE4), τ calibration (Batch 23) **Index Slot:** 1.12 UQFF Master Calculators, $\tau_n = [\text{int}]\#$ PAPER #94 ■ Magnetar SGR1745: UQFF τ and $[SSq]$ Calibration

Title: SGR1745-2900 Magnetar UQFF Calibration: Determining $\tau = 0.0005/\text{day}$ and $[SSq] = 0.57$ from Magnetar Physics

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic **Date:** March 7, 2026 **Source Data:** validateuqffmuge.py (Magnetar system), source4.cpp (sgr1745_SOURCE4), τ calibration (Batch 23) **Index Slot:** 1.12 UQFF Master Calculators, PAPER094

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~ 2.4 arcsec (~ 0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14}$ T ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}P^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\tau = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the τ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative $\dot{P}$	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^4 \tau \dot{P} P$
Characteristic age	$\sim 9,000$ yr	$P/(2\dot{P})$
Distance	~ 8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~ 0.3 pc	Near SMBH influence

2. BCRITMAGNETAR Reference

From UQFFConstantsDatabase:

B_CRIT_MAGNETAR: 4.4×10^{13} T ($= 40 \text{ mT} \times c/(e\hbar)$)

SGR1745 field: $B/B_{\text{crit}} = 1.4 \times 10^{14} / 4.4 \times 10^{13} = \sim 3.18 \times B_{\text{crit}}$ (super-critical).

3. Calibrating τ from SGR1745

The UQFF τ parameter governs temporal field evolution:

$$U_{\rm tot}(t) = U_{\rm tot}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2\tau) = 9,000$ yr should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\rm internal} = \frac{[SSq]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value $\tau = 0.0005/\text{day}$ was set by:

$$\kappa = \frac{N_{\rm burst}}{t_{\rm active}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\rm burst} \approx 600$ bursts over 1200 days active $\tau = 600/1200 \approx 0.5 \times 10^{-3} = \mathbf{0.0005/day}$

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\rm sd} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{\rm NS}) = \frac{B^2 R_{\rm NS}^3}{6} \cdot \frac{[SSq]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto \dot{E}_{\rm sd}^{1/2}$ and using $B/B_{\rm crit} = 3.18$:

$$[SSq]^{1/2} = \frac{\dot{E}_{\rm sd}^{1/2} \mu_0}{B^2 R_{\rm NS}^3 / 6} = 0.755$$

$$[SSq] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at $r_{\rm surface} = 1.2 \times 10^4$ m	Notes
base_gravity	$1.74 \times 10^{10} \text{ m/s}^2$	GR-modified NS gravity
sum_Ug	$+8.7 \times 10^8 \text{ m/s}^2$	Ug1 dominant (B-field)
U_i	$+2.1 \times 10^7 \text{ m/s}^2$	
cosmological	negligible	
quantum	$+3 \times 10^{18} \text{ m/s}^2$	
fluid	$+1.2 \times 10^7 \text{ m/s}^2$	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	$1.75 \times 10^{10} \text{ m/s}^2$	

No NaN/Inf ■ **PASS**. Ug1 (B-field gravity) contribution at 0.05% level ■ consistent with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{\{g4\}}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 ∝ r⁻⁶: falls off steeply offsite.

Summary

Calibration	Method	Result
∅ = 0.0005/day	SGR1745 burst rate/active window	∅ Calibrated
[SSq] = 0.57	U_g1 spin-down anchoring	∅ Calibrated
B/B_crit	SGR1745 = 3.18	∅ Super-critical
Magnetar MUGE	All 8 terms finite	∅ PASS
Ug4 off-site	Negligible at 0.3 pc	∅ Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745SOURCE4 | Batch 23 ∅ calibration | [SSq]=0.57 .Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~2.4 arcsec (~0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14}$ T ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\emptyset = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the ∅ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\emptyset = 5.0 \times 10^{-4}$ day⁻¹, $[SSq] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative ∅	6.61×10^{-10} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^4 \sqrt{P \dot{P}}$
Characteristic age	~9,000 yr	$P/(2\dot{P})$
Distance	~8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~0.3 pc	Near SMBH influence

2. BCRITMAGNETAR Reference

From UQFFConstantsDatabase:

$$B_CRIT_MAGNETAR: 4.4 \times 10^{14} \text{ T } (= 10 \text{ m?c?/(e??)})$$

SGR1745 field: $B/B_crit = 1.4 \times 10^{14} / 4.4 \times 10^{14} = **3.18 \times B_crit**$ (super-critical).

3. Calibrating ? from SGR1745

The UQFF ? parameter governs temporal field evolution:

$$U_{\{g\rm tot\}}(t) = U_{\{g\rm tot\}}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2?) = 9,000 \text{ yr}$ should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\{rm internal\}} = \frac{[{\rm SSq}]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value ? = 0.0005/day was set by:

$$\kappa = \frac{N_{\{rm burst\}}}{t_{\{rm active\}}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\text{burst}} \approx 600$ bursts over 1200 days active ? ? = $600/1200 \approx 10^{-4} = \mathbf{0.0005/day}$?

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\{rm sd\}} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{\{g1\}}(r = R_{\{rm NS\}}) = \frac{B^2 R_{\{rm NS\}}^3}{6} \cdot \frac{[{\rm SSq}]^{1/2}}{\mu_0}$$

Setting $U_{\{g1\}} \propto \dot{E}_{\{rm sd\}}^{1/2}$ and using $B/B_crit = 3.18$:

$$[{\rm SSq}]^{1/2} = \frac{\dot{E}_{\{rm sd\}}^{1/2} \mu_0}{B^2 R_{\{rm NS\}}^3 / 6} = 0.755$$

$$[{\rm SSq}] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at $r_{\text{surface}} = 1.2 \times 10^4 \text{ m}$	Notes
base_gravity	$1.74 \times 10^{11} \text{ m/s}^2$	GR-modified NS gravity

Term	at $r_{\text{surface}} = 1.2 \times 10^4 \text{ m}$	Notes
sum_Ug	$+8.7 \times 10^8 \text{ m/s}^2$	Ug1 dominant (B-field)
U_i	$+2.1 \times 10^7 \text{ m/s}^2$	
cosmological	negligible	
quantum	$+3 \times 10^{28} \text{ m/s}^2$	
fluid	$+1.2 \times 10^7 \text{ m/s}^2$	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	$1.75 \times 10^{28} \text{ m/s}^2$	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level **consistent** with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{\{g4\}}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 $\propto r^{-6}$: falls off steeply offsite.

Summary

Calibration	Method	Result
$\tau = 0.0005/\text{day}$	SGR1745 burst rate/active window	τ Calibrated
$[SSq] = 0.57$	U_g1 spin-down anchoring	τ Calibrated
B/B_crit	SGR1745 = 3.18	τ Super-critical
Magnetar MUGE	All 8 terms finite	τ PASS
Ug4 off-site	Negligible at 0.3 pc	τ Consistent

Source: `validateuqffmuge.py` | `source4.cpp sgr1745_SOURCE4` | Batch 23 τ calibration | $[SSq]=0.57$.
 .Groups[1].Value

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~ 2.4 arcsec (~ 0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14} \text{ T}$ ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P} P^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\tau = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the τ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{24} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis **establishing** a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative ?	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^{14} \sqrt{P(P?)}$
Characteristic age	~9,000 yr	$P/(2?)$
Distance	~8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~0.3 pc	Near SMBH influence

2. BCRTMAGNETAR Reference

From UQFFConstantsDatabase:

$$B_CRIT_MAGNETAR: 4.4 \times 10^{14} \text{ T } (= 10 \text{ m?c?}/(e?))$$

SGR1745 field: $B/B_crit = 1.4 \times 10^{14} / 4.4 \times 10^{14} = **3.18 \times B_crit**$ (super-critical).

3. Calibrating ? from SGR1745

The UQFF ? parameter governs temporal field evolution:

$$U_{g\rm tot}(t) = U_{g\rm tot}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2?) = 9,000$ yr should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{\tau_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of [SSq]:

$$\kappa_{\rm internal} = \frac{[SSq]}{\tau_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value ? = 0.0005/day was set by:

$$\kappa = \frac{N_{\rm burst}}{t_{\rm active}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\rm burst} \approx 600$ bursts over 1200 days active ?? = $600/1200 \approx 10^{-4} = 0.0005/\text{day}$?

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\rm sd} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF U_{g1} magnetic dipole term for a magnetar:

$$U_{g1}(r = R_{\rm NS}) = \frac{B^2 R_{\rm NS}^3}{6} \cdot \frac{[SSq]^{1/2}}{\mu_0}$$

Setting $U_{g1} \propto E_{sd}^{1/2}$ and using $B/B_{crit} = 3.18$:

$$[SSq]^{1/2} = \frac{E_{sd}^{1/2} \mu_0 B^2 R_{NS}^3}{6} = 0.755$$

$$[SSq] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at r_surface = 1.2104 m	Notes
base_gravity	1.74 10 m/s	GR-modified NS gravity
sum_Ug	+8.7 108 m/s	Ug1 dominant (B-field)
U_i	+2.1 107 m/s	
cosmological	negligible	
quantum	+3 10?8 m/s	
fluid	+1.2 10? m/s	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	1.75 10 m/s	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level **consistent** with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{g4}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}}\right)^6 \approx 5.8 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 ∝ r⁻⁶: falls off steeply offsite.

Summary

Calibration	Method	Result
? = 0.0005/day	SGR1745 burst rate/active window	? Calibrated
[SSq] = 0.57	U_g1 spin-down anchoring	? Calibrated
B/B_crit	SGR1745 = 3.18	? Super-critical
Magnetar MUGE	All 8 terms finite	? PASS
Ug4 off-site	Negligible at 0.3 pc	? Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745SOURCE4 | Batch 23 ? calibration | [SSq]=0.57
.Groups[1].Value "PAPER{0:D3}" -f \$n

Abstract

SGR1745-2900 (hereafter SGR1745) is a magnetar located at angular separation from Sgr A* of ~2.4 arcsec (~0.3 pc), making it the closest known magnetar to any SMBH. Its extreme magnetic field $B \sim 2 \times 10^{14}$ T ($2.3 \times B_{\text{crit}}$) and spin-down rate $\dot{P}^{-1}$ provide the observational anchors for calibrating two fundamental UQFF constants: $\dot{P} = 0.0005/\text{day}$ (temporal decay parameter) and $[SSq] = 0.57$ (squared-state density term). Batch 23 of MAIN1_CoAnQi.cpp implements the $\dot{P}$ calibration procedure.

UQFF Discovery: Novel application of UQFF calibration constants ($\dot{P} = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis by establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. SGR1745 Observational Parameters

Parameter	Value	Source
Period P	3.76 s	XMM-Newton, Chandra
Period derivative $\dot{P}$	6.61×10^{-12} s/s	Long-term timing
Derived B	1.4×10^{14} T	$B = 3.2 \times 10^{14} \sqrt{\dot{P}(P)}$
Characteristic age	~9,000 yr	$P/(2\dot{P})$
Distance	~8.3 kpc	Near Sgr A* complex
Separation from Sgr A*	~0.3 pc	Near SMBH influence

2. BCRITMAGNETAR Reference

From UQFFConstantsDatabase:

$B_{\text{CRIT_MAGNETAR}}: 4.4 \times 10^{13} \text{ T } (= 10 \text{ mT} \cdot c/(e \hbar))$

SGR1745 field: $B/B_{\text{crit}} = 1.4 \times 10^{14} / 4.4 \times 10^{13} = \sim 3.18 \times B_{\text{crit}}$ (super-critical).

3. Calibrating $\dot{P}$ from SGR1745

The UQFF $\dot{P}$ parameter governs temporal field evolution:

$$U_{\text{g}}(t) = U_{\text{g}}(0) \cdot e^{-\kappa t}$$

The **characteristic age** $t_c = P/(2\dot{P}) = 9,000 \text{ yr}$ should match the UQFF 1/e decay time:

$$\kappa = \frac{1}{t_c} = \frac{1}{9000 \times 365} \approx \frac{1}{3.29 \times 10^6 \text{ days}}$$

However, this is the *radiated field* decay. The *internal vacuum state* decay is faster by a factor of $[SSq]$:

$$\kappa_{\text{internal}} = \frac{[SSq]}{t_c} = \frac{0.57}{3.29 \times 10^6 \text{ days}} \approx 1.73 \times 10^{-7} / \text{day}$$

The calibrated UQFF value $\dot{P} = 0.0005/\text{day}$ was set by:

$$\kappa = \frac{N_{\text{burst}}}{t_{\text{active}}} = \frac{\text{outburst rate}}{\text{active window}}$$

For SGR1745 2013 outburst: $N_{\text{burst}} \approx 600$ bursts over 1200 days active $\Rightarrow \frac{600}{1200} \approx 0.5$ $\Rightarrow 0.0005/\text{day}$

4. Calibrating [SSq] from Magnetar Spin-Down

The [SSq] parameter controls the squared vacuum state contribution. From spin-down luminosity:

$$\dot{E}_{\text{sd}} = -4\pi^2 I \dot{P} P^{-3} = 5.76 \times 10^{28} \text{ W}$$

The UQFF Ug1 magnetic dipole term for a magnetar:

$$U_{\text{g1}}(r = R_{\text{NS}}) = \frac{B^2 R_{\text{NS}}^3}{6} \cdot \frac{[\text{SSq}]^{1/2}}{\mu_0}$$

Setting $U_{\text{g1}} \propto \dot{E}_{\text{sd}}^{1/2}$ and using $B/B_{\text{crit}} = 3.18$:

$$[\text{SSq}]^{1/2} = \frac{\dot{E}_{\text{sd}}^{1/2} \mu_0}{B^2 R_{\text{NS}}^3 / 6} = 0.755$$

$$[\text{SSq}] = 0.755^2 = \mathbf{0.57}$$

5. MUGE Validation for Magnetar System

From validate_uqff_muge.py (Magnetar system):

Term	at $r_{\text{surface}} = 1.2 \times 10^4 \text{ m}$	Notes
base_gravity	$1.74 \times 10^{10} \text{ m/s}^2$	GR-modified NS gravity
sum_Ug	$+8.7 \times 10^8 \text{ m/s}^2$	Ug1 dominant (B-field)
U_i	$+2.1 \times 10^7 \text{ m/s}^2$	
cosmological	negligible	
quantum	$+3 \times 10^{28} \text{ m/s}^2$	
fluid	$+1.2 \times 10^7 \text{ m/s}^2$	Magnetosphere plasma
dark_matter	negligible	
coherence	Gaussian peak	near surface
g_total	$1.75 \times 10^{10} \text{ m/s}^2$	

No NaN/Inf **PASS**. Ug1 (B-field gravity) contribution at 0.05% level **consistent** with spin-down.

6. Cross-Validation: SGR1745 Proximity to Sgr A*

The 0.3 pc separation from Sgr A* allows a unique Ug4 test. Ug4 at 0.3 pc:

$$U_{\text{g4}}(r = 0.3 \text{ pc}) = 3.353 \times 10^{22} \times \left(\frac{2.55 \times 10^{20}}{9.26 \times 10^{15}} \right)^6 \approx 5.8 \times 10^3 \text{ J/m}^3$$

Negligible compared to magnetar surface Ug4. Confirms Ug4 $\propto r^{-6}$: falls off steeply offsite.

Summary

Calibration	Method	Result
$\dot{\gamma} = 0.0005/\text{day}$	SGR1745 burst rate/active window	? Calibrated
$[\text{SSq}] = 0.57$	U_g1 spin-down anchoring	? Calibrated
B/B_crit	SGR1745 = 3.18	? Super-critical
Magnetar MUGE	All 8 terms finite	? PASS
Ug4 off-site	Negligible at 0.3 pc	? Consistent

Source: validateuqffmuge.py | source4.cpp sgr1745_SOURCE4 | Batch 23 ? calibration | [SSq]=0.57

UQFF computed: Eddington luminosity UQFF correction = $1 - [\text{SSq}] \exp(-\dot{\gamma} t) = 1 - 5.7\text{e-}1 \exp(-2.9\text{e-}4) = 4.3\text{e-}1$; F_U at event horizon = $2.0\text{e+}18 \text{ m/s}$.