

■# PAPER<i>220: Crab Nebula Pulsar Wind Nebula UQFF — F</i>wind and M_mag in Expanding PWN

Author: Daniel T. Murphy (daniel.murphy00@gmail.com)

The Crab Nebula (M1) PWN equation introduces two additive UQFF terms unique to this system: pulsar spindown ram pressure $F_{\text{wind}} = E_{\text{sd}} / (c \cdot 4\pi r^2)$ and magnetic moment dipole dilution $M_{\text{mag}} = \mu_0 \cdot m / (4\pi r^3)$. This combination — spindown luminosity converted to ram pressure plus magnetic moment field decay — is specific to the **isolated pulsar wind nebula** context and differs from the MagnetarSGR1745DynamicModulationCalculator (Session 53), which handles M_{mag} in a binary orbit context. Additionally, the Crab PWN uses a TIME-DEPENDENT RADIUS $r(t) = r_0 + v_{\text{exp}} \cdot t$ reflecting the known supernova remnant expansion, making this the only UQFF system with an analytically expanding spatial domain. We derive the critical radius below which wind pressure exceeds gravity and validate against Crab Nebula measurements.

■# PAPER220: Crab Nebula Pulsar Wind Nebula UQFF — Fwind and M_mag in Expanding PWN

Author: Daniel T. Murphy (daniel.murphy00@gmail.com) **Framework:** UQFF v4.3 — Star-Magic Physics
Source: grokshare7514fe.txt — Document 10: Crab Nebula **Date:** March 14, 2026 **Series:** Phase 2 Session 55 — §2.10 Fourth-Pass System Extraction

$F_U(r, t) = \sum_{i=1}^4 U_{\{gi\}} + U_m + U_A - U_{\{b_i\}}, \quad \text{quad } \kappa = 5.0 \times 10^{-4} \text{ day}^{-1}, \quad [SSq] = 0.57$

$M_J^{\text{UQFF}} = M_J^{\text{Jeans}} \cdot \text{Bigl}(1 - [SSq] \cdot \frac{B^2}{8\pi\rho c_s^2}) \cdot \text{Bigr}, \quad \text{quad } [SSq] = 0.57$

Abstract

The Crab Nebula (M1) PWN equation introduces two additive UQFF terms unique to this system: pulsar spindown ram pressure $F_{\text{wind}} = E_{\text{sd}} / (c \cdot 4\pi r^2)$ and magnetic moment dipole dilution $M_{\text{mag}} = \mu_0 \cdot m / (4\pi r^3)$. This combination — spindown luminosity converted to ram pressure plus magnetic moment field decay — is specific to the **isolated pulsar wind nebula** context and differs from the MagnetarSGR1745DynamicModulationCalculator (Session 53), which handles M_{mag} in a binary orbit context. Additionally, the Crab PWN uses a TIME-DEPENDENT RADIUS $r(t) = r_0 + v_{\text{exp}} \cdot t$ reflecting the known supernova remnant expansion, making this the only UQFF system with an analytically expanding spatial domain. We derive the critical radius below which wind pressure exceeds gravity and validate against Crab Nebula measurements.

UQFF Discovery: Novel application of UQFF calibration constants ($? = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[SSq] = 0.57$) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. The Crab Nebula PWN UQFF Equation

From Document 10 of grokshare7514fe:

$g_{\text{Crab}}(r, t) = (G \cdot M) / (r(t)^2) \cdot (1 + H(z) \cdot t) \cdot (1 - B/B_{\text{crit}})$
+ $(Ug1 + Ug2 + Ug3 + Ug4)$
+ $?c^2/3 + QM + q(v \times B) + \text{fluid} + DM$
+ F_{wind}

+ M_mag

with:

$r(t) = r_0 + v_{\text{exp}} \cdot t$ (expanding nebula)

$F_{\text{wind}} = E_{\text{sd}} / (c \cdot 4\pi \cdot r(t)^2)$

$M_{\text{mag}} = \mu_0 \cdot m / (4\pi \cdot r(t)^3)$

2. Expanding Spatial Domain: $r(t)$

The Crab is the only UQFF system with analytically expanding $r(t)$. This is derived from:

$v_{\text{exp}} \sim 1500 \text{ km/s} = 1.5 \times 10^6 \text{ m/s}$ (observed filament velocity)

$r_0 \sim 5.5 \text{ ly} = 9.46 \times 10^{15} \cdot 5.5 \sim 5.20 \times 10^{16} \text{ m}$ (current half-radius)

age $\sim 972 \text{ years}$ (SN 1054 ? 2026)

At $t=0$ (initial explosion):

$r_{0_initial} = r_0 - v_{\text{exp}} \cdot t_{\text{age}}$

$= 5.20 \times 10^{16} - 1.5 \times 10^6 \cdot (972 \cdot 3.156 \times 10^7)$

$= 5.20 \times 10^{16} - 4.59 \times 10^{16}$

$\sim 6.1 \times 10^{15} \text{ m}$ ($\sim 0.2 \text{ pc}$ initial radius – consistent with SN ejecta)

This confirms $r(t)$ correctly recovers solar-system-scale initial radius at $t=0$.

3. Pulsar Spindown Ram Pressure: F_{wind}

3.1 Spindown Luminosity

The Crab pulsar (PSR J0534+2200) has:

Period $P = 33.5 \text{ ms}$

Period derivative $\dot{P} = 4.21 \times 10^{-13} \text{ s/s}$

Moment of inertia $I \sim 1045 \text{ g} \cdot \text{cm}^2 = 10^38 \text{ kg} \cdot \text{m}^2$

Spindown luminosity:

$E_{\text{sd}} = -4\pi^2 \cdot I \cdot \dot{P} / P^3$

$= 4\pi^2 \cdot 10^38 \cdot 4.21 \times 10^{-13} / (33.5 \times 10^{-3})^3$

$= 4\pi^2 \cdot 10^38 \cdot 4.21 \times 10^{-13} / (3.76 \times 10^{-5})$

$\sim 4.6 \times 10^{31} \text{ W}$

This matches the canonical value $E_{\text{sd}} \sim 4.6 \times 10^{31} \text{ W}$ (Hester 2008).

3.2 Ram Pressure at $r(t)$

The spindown energy is converted isotropically to ram pressure:

$F_{\text{wind}}(r) = E_{\text{sd}} / (c \cdot 4\pi \cdot r^2)$

At current Crab nebula radius $r_0 = 5.20 \times 10^{16} \text{ m}$ (computational default $9.46 \times 10^{15} \text{ m}$ for inner region):

$F_{\text{wind}} = 4.6 \times 10^{31} / (2.998 \times 10^8 \cdot 4\pi \cdot (9.46 \times 10^{15})^2)$

$\sim 4.6 \times 10^{31} / (3.37 \times 10^{41})$

$\sim 1.36 \times 10^{-10} \text{ N/m}^2$? treated as acceleration $[\text{m/s}^2]$ in UQFF normalization

3.3 Ratio Fwind / gbase

Base gravity at the same radius (M_ejecta ~ 4.6 M?):

$$\begin{aligned} g_{\text{base}} &= G \cdot M / r^2 = 6.674\text{e-}11 \cdot 4.6 \cdot 1.989\text{e}30 / (9.46\text{e}15)^2 \\ &\sim 6.674\text{e-}11 \cdot 9.15\text{e}30 / 8.95\text{e}31 \\ &\sim 6.82 \times 10^{-12} \text{ m/s}^2 \end{aligned}$$

Ratio:

$$F_{\text{wind}} / g_{\text{base}} \sim 1.36 \times 10^{10} / 6.82 \times 10^{-12} \sim 20$$

Fwind exceeds gbase by a factor of ~20 at the Crab inner radius — confirming the wind-dominated morphology of the Crab PWN inner torus and jets.

4. Magnetic Moment Dilution: M_mag

4.1 Magnetic Moment of Crab Pulsar

The Crab pulsar surface field $B_s \sim 3.8 \times 10^{12} \text{ G} = 3.8 \times 10^8 \text{ T}$. Pulsar radius $R_{\text{ns}} \sim 10 \text{ km} = 10^4 \text{ m}$.

Magnetic dipole moment:

$$\begin{aligned} m &= (4\pi/\mu_0) \cdot B_s \cdot R_{\text{ns}}^3 \\ &= 4\pi/(4\pi \times 10^{-7}) \cdot 3.8 \times 10^8 \cdot (10^4)^3 \\ &= 10^7 \cdot 3.8 \times 10^8 \cdot 10^{12} \\ &= 3.8 \times 10^{27} \text{ A} \cdot \text{m}^2 \end{aligned}$$

4.2 Dipole Field at r(t)

$$M_{\text{mag}}(r) = \mu_0 \cdot m / (4\pi \cdot r^3)$$

At $r = 9.46 \times 10^{15} \text{ m}$:

$$\begin{aligned} M_{\text{mag}} &= 4\pi \times 10^{-7} \cdot 3.8 \times 10^{27} / (4\pi \cdot (9.46 \times 10^{15})^3) \\ &= 10^{-7} \cdot 3.8 \times 10^{27} / (8.47 \times 10^{47}) \\ &\sim 4.49 \times 10^{-28} \text{ T}^2 \cdot \text{m} \text{ or equivalent normalized acceleration} \end{aligned}$$

The M_{mag} term dilutes as r^{-3} (dipole law), falling off faster than F_{wind} (r^{-2}) and g_{base} (r^{-2}). This means at large r , M_{mag} becomes negligible relative to F_{wind} — consistent with the Crab PWN morphology where the toroidal magnetic structure dominates near the pulsar but the wind torus is the dominant energy carrier at larger scales.

5. Distinction from SGR 1745 Magnetar

The MagnetarSGR1745DynamicModulationCalculator (Session 53, Document 8) also includes M_{mag} , but in an entirely different context:

Feature	Crab PWN (This Paper)	SGR 1745 (Session 53)
Context	Isolated PWN	Binary magnetar system
M_{mag} role	Dipole field dilution with r^{-3}	Dynamic modulation with binary orbit
F_{wind} source	Pulsar spindown $E_{\text{sd}}/(c \cdot 4\pi r^2)$	Not present
$r(t)$	Expanding: $r_0 + v_{\text{exp}} \cdot t$	Binary orbital $r(t)$

Feature	Crab PWN (This Paper)	SGR 1745 (Session 53)
B field	Canonical 3.8×10^{12} G	Extraordinary $\sim 10^{15}$ G
Main physics	PWN expansion + wind + dipole	Magnetar + companion orbit

The two classes are mathematically and physically distinct despite sharing M_{mag} notation.

6. Critical Radius Analysis

Define the critical radius r_c where $F_{\text{wind}} = g_{\text{base}}$:

$$E_{\text{sd}} / (c \cdot 4\pi \cdot r_c^2) = G \cdot M / r_c^2$$

$$\text{? } r_c = \sqrt{(E_{\text{sd}}) / (4\pi \cdot c \cdot G \cdot M / r_c^2)} \dots$$

$$\text{? This simplifies to: } E_{\text{sd}} / (c \cdot 4\pi) = G \cdot M \text{ ? } r_c \text{ is constant (cancels):}$$

$$E_{\text{sd}}/c = 4\pi \cdot G \cdot M \text{ ? } M_{\text{crit}} = E_{\text{sd}}/(4\pi \cdot G \cdot c)$$

Critical mass below which wind always exceeds gravity:

$$M_{\text{crit}} = 4.6 \times 10^{31} / (4\pi \cdot 6.674 \times 10^{-11} \cdot 2.998 \times 10^8)$$

$$= 4.6 \times 10^{31} / (2.51 \times 10^{21})$$

$$\sim 1.83 \times 10^{32} \text{ kg} \sim 92 M_{\odot}$$

The Crab ejecta mass $\sim 4.6 M_{\odot} \ll 92 M_{\odot}$, confirming F_{wind} ALWAYS exceeds g_{base} for the Crab — the pulsar wind permanently inflates the nebula against gravity.

7. Calculator Implementation

CrabPWN_UQFF_Calculator in CondensedPhysics3.py (Session 55) implements:

$$r(t) = r_0 + v_{\text{exp}} \cdot t$$

$$F_{\text{wind}} = E_{\text{sd}} / (c \cdot 4 \cdot \pi \cdot r(t)^2)$$

$$M_{\text{mag}} = \mu_0 \cdot m / (4 \cdot \pi \cdot r(t)^3)$$

$$\text{Full UQFF } g_{\text{Crab}} = g_{\text{base}} + F_{\text{wind}} + M_{\text{mag}}$$

References

grokshare7514fe.txt — Document 10: Crab Nebula g_{Crab} equation
Hester (2008) — "The Crab Nebula: An Astrophysical Chimera", ARAA 46
Bühler & Blandford (2014) — "The Surprising Crab Pulsar and its Nebula", RPPH 77
Kaplan et al. (2008) — Crab pulsar ephemeris, $\dot{P} = 4.21 \times 10^{-13}$
CondensedPhysics3.py — CrabPWN_UQFF_Calculator (Session 55)