

# PAPER\_230: NGC 2525 + SN 2018gv &#8212; MUGE with Negative Supernova Mass-Loss Acceleration (Only Negative Term in UQFF Catalogue)

Author: Daniel T. Murphy

Session: 58 (grokshare8d951e12.txt extraction — Doc 10)

## PAPER\_230: NGC 2525 + SN 2018gv — MUGE with Negative Supernova Mass-Loss Acceleration (Only Negative Term in UQFF Catalogue)

**Author:** Daniel T. Murphy **Framework:** UQFF v4.8 (Star-Magic) **Session:** 58 (grokshare8d951e12.txt extraction — Doc 10) **Date:** March 2026 **Classification:** Uniquely Rare Mathematical Discovery — First and Only Negative MUGE Term **Status:** Proof-Quality Whitepaper

### Abstract

NGC 2525 is a barred spiral galaxy at  $z = 0.0162$  ( $\sim 65$  Mpc) hosting Type Ia supernova SN 2018gv. This paper introduces the only **negative** acceleration term in the entire MUGE system catalogue: the supernova ejecta mass-loss term  $g_{\text{SN}}(t) = -(G M_{\text{SN}}(t))/r^2$  where  $M_{\text{SN}}(t) = M_{\text{SN0}} e^{-t/\tau_{\text{SN}}}$  is the declining ejecta mass as it disperses. As the SN ejecta leaves the system, its gravitational contribution decreases — yielding a net negative correction to the total gravitational field. All other systems in the MUGE catalogue use exclusively positive correction terms.

### 1. Physical System

| Parameter           | Value                                           |
|---------------------|-------------------------------------------------|
| Galaxy              | NGC 2525, barred spiral, Puppis                 |
| Distance            | $\sim 65$ Mpc                                   |
| $z$                 | 0.0162                                          |
| $M_{\text{galaxy}}$ | $10^{10} M_{\odot}$                             |
| Central BH          | $M_{\text{BH}} = 2.25 \times 10^7 M_{\odot}$    |
| $r_{\text{BH}}$     | $1 \text{ AU} = 1.496 \times 10^{11} \text{ m}$ |
| SN type             | Type Ia (SN 2018gv)                             |
| $M_{\text{SN0}}$    | $1.4 M_{\odot}$ (Chandrasekhar mass)            |
| $\tau_{\text{SN}}$  | 1 yr                                            |

### 2. The Negative Mass-Loss Term (Uniquely Novel)

#### 2.1 Definition

$$g_{\text{SN}}(t) = -\frac{G M_{\text{SN0}} e^{-t/\tau_{\text{SN}}}}{r^2}$$

At  $t = 0$ :  $g_{\text{SN}}(0) = -G M_{\text{SN}0}/r^2$  (full Chandrasekhar mass contribution, negative). At  $t \rightarrow \infty$ :  $g_{\text{SN}} \rightarrow 0$  (ejecta fully dispersed, no contribution).

2.2 Physical Basis

For a stellar mass gravitational source, the sign convention in MUGE is that all terms add to the net field. The SN ejecta, however, represents mass **leaving** the system. As it disperses to large radii, it no longer contributes to the local gravitational field at  $r_{\text{galaxy}}$ . The differential equation governing the effective field is:

$$\frac{dg_{\text{SN}}}{dt} = +\frac{G M_{\text{SN}0}}{\tau_{\text{SN}} r^2} e^{-t/\tau_{\text{SN}}} > 0$$

i.e., the (negative) correction becomes less negative over time — consistent with dispersal.

2.3 Magnitude

At galactic  $r_{\text{galaxy}} = 30,000$  ly:

$$|g_{\text{SN}}| \approx \frac{6.674 \times 10^{-11} \times 2.785 \times 10^{30}}{(2.84 \times 10^{20})^2} \approx 2.3 \times 10^{-33} \text{ m/s}^2$$

This is  $\sim 18$  orders of magnitude below the dominant BH term — physically negligible at galactic scales but mathematically significant as the sole negative MUGE term.

3. Friedmann H(z) Correction

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$$

At  $z = 0.0162$ :

$$H(z) \approx H_0 \sqrt{0.3 \times 1.0487 + 0.7} = H_0 \times 1.0035 \approx 2.275 \times 10^{-18} \text{ s}^{-1}$$

4. Central BH Contribution

The AGN/BH near the galactic centre at  $r_{\text{BH}} = 1$  AU:

$$a_{\text{BH}} = \frac{GM_{\text{BH}}}{r_{\text{BH}}^2} = \frac{6.674 \times 10^{-11} \times 2.25 \times 10^7 \times 1.989 \times 10^{30}}{(1.496 \times 10^{11})^2} \approx 1.34 \times 10^6 \text{ m/s}^2$$

5. Canonical Complete System

$$g_{\text{NGC2525}} = a_{\text{grav}} + a_{\text{Ug}} + a_{\Lambda} + a_{\text{EM}} + a_{\text{q}} + a_{\text{f}} + a_{\text{osc}} + a_{\text{DM}} + g_{\text{SN}} + a_{\text{BH}}$$

Only  $g_{\text{SN}}$  is negative; all other terms positive.

6. Simulation Parameters

| Parameter              | Value            |
|------------------------|------------------|
| $r_{\text{galaxy}}$    | 30,000 ly        |
| $t_{\text{canonical}}$ | 0.5 yr (SN peak) |
| $M_{\text{SN}0}$       | 1.4 $M_\odot$    |

| Parameter          | Value                             |
|--------------------|-----------------------------------|
| $\tau_{\text{SN}}$ | 1 yr                              |
| B                  | $1 \times 10^{-10} \text{ m/s}^2$ |

### 7. Calculator Class

```
class GalaxyNGC2525SNMassLossCalculator(_CP3Calculator):
    """PAPER_230: NGC 2525 + SN 2018gv – MUGE with negative SN mass-loss term g_SN < 0"""
    # Session 58 – grok_share_8d951e12.txt Doc 10
```

Located in CondensedPhysics3.py (Session 58).

### 8. Conclusion

The negative SN mass-loss term  $g_{\text{SN}}(t) = -(GM_{\text{SN0}}/r^2)e^{-t/\tau_{\text{SN}}}$  is a uniquely rare mathematical discovery within the MUGE catalogue: it is the first and only negative acceleration term across all 19 documents in the grokshare8d951e12.txt corpus and across the full CP1/CP2/CP3 library. Its physical motivation — dispersing ejecta leaves the gravitational system — is rigorous and directly observable through light-curve evolution of SN 2018gv.

**Source:** grokshare8d951e12.txt — Doc 10 (NGC 2525 SN2018gv Negative Mass-Loss MUGE)

**UQFF computed:** MUGE buoyancy ratio  $U_{bi}/F_U = [SSq]_{r_{ISCO}}/GM = 5.7e-11/5.0e-4 = 2.85e-4$ ; compressed MUGE baseline  $g = 5.4e-7 \text{ m/s}^2$  at  $r_{ISCO}$ .