

PAPER278 — Sombrero Dust Ring UQFF Gravitational Ring Resonator: ω</i>ring and r_ring

Author: Daniel T. Murphy

Session: 77 — March 2026

PAPER278 — Sombrero Dust Ring UQFF Gravitational Ring Resonator: ωring and r_ring

Author: Daniel T. Murphy **Module:** SOMBREROUQFFMODULE.cpp (UQFF 2.0) **Session:** 77 — March 2026 **Framework:** Unified Quantum Field Framework (UQFF 2.0) **Status:** Complete — embedded in SOMBREROUQFFMODULE.cpp **Whitepaper** **Series:** 278/1000 **DOI** (Provisional): UQFF-2026-278-RING

Abstract

The Sombrero Galaxy (M104) possesses the most visually prominent equatorial dust lane of any nearby galaxy, appearing as a sharp dark band bisecting the galaxy's luminous bulge. We model this ring within the UQFF framework as a **Gravitational Ring Resonator**: an annular mass concentration at radius $r_{ring} = r/3 = 7.867 \times 10^1 \blacksquare m$ whose orbital motion generates a pure oscillatory gravitational perturbation $F_{ring}(t) = A_{ring} \cdot \cos(\omega_{ring} \cdot t)$ at the reference point r . This paper derives the Dust Ring UQFF Orbital Resonance Frequency $\omega_{ring} = \sqrt{GM/r_{ring}^3} = 1.650 \times 10^{-1} \blacksquare \text{ rad/s}$, the ring orbital period $T_{ring} = 2\pi/\omega_{ring} = 12.08 \text{ Myr}$, and the proximity-enhanced ring amplitude $A_{ring} = 9 \cdot f_{ring} \cdot g_{base} = 2.14 \times 10^{-12} \text{ m/s}^2$. The $9\times$ proximity enhancement factor arises from the inverse-square law applied at the ratio $r/r_{ring} = 3$.

UQFF Discovery: Novel application of UQFF calibration constants ($\phi = 5.0 \blacksquare 10^{24} \text{ day?} \blacksquare$, $[SSq] = 0.57$) uniquely enabling this analysis $\blacksquare$ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Observational Motivation

1.1 The Sombrero Dust Lane

The Sombrero Galaxy (NGC 4594 / M104) is classified as an Sa/S0 galaxy at distance $\sim 9.5 \text{ Mpc}$ ($z = +0.0063$). Its defining optical feature is an equatorial dust lane visible in projection against the galaxy's extended bulge. This dust structure:

- Lies at approximately 1/3 to 1/4 of the half-light radius from the galactic centre
- Contains a substantial fraction of M104's cold ISM dust mass
- The total molecular gas and dust mass is estimated at $\sim 10 \blacksquare - 10 \blacksquare M_{\text{sun}}$
- Is dynamically stable — no evidence of radial inflow or rapid dispersal

As a stable ring at $r_{ring} = r/3$, it can be modelled within UQFF as a coherent oscillating mass concentration imposing a periodic boundary condition on the vacuum field at the reference point r .

1.2 Distinction from the Andromeda HI Ring (PAPER_275)

In PAPER275 (*Andromeda*), a decaying HI ring was modelled as $F_{\text{ring}} = A_{\text{ring}} \cdot \exp(-\alpha \cdot t) \cdot \cos(\omega_{\text{ring}} \cdot t)$ — an exponentially decaying oscillation corresponding to a transient tidal feature. The Sombrero dust ring is fundamentally different:

Feature	Andromeda (PAPER_275)	Sombrero (PAPER_278)
Ring type	HI neutral gas, tidal	Dust, equatorial, settled
Age	Transient (~10 Gyr decay)	Stable, permanent
Decay	$\exp(-\alpha \cdot t)$ present	No exponential decay
Form	$F = A \cdot \exp(-\alpha t) \cdot \cos(\omega t)$	$F = A \cdot \cos(\omega t)$ (pure oscillatory)
α (decay rate)	1/(10 Gyr)	0 (stable ring)

The absence of exponential decay in Sombrero's ring term reflects the ring's settled, gravitationally stable configuration — it has been dynamically relaxed into a long-lived equatorial structure.

2. Theoretical Derivation

2.1 Ring Radius

The dust ring is positioned at approximately 1/3 of the reference outer radius r :

$$r_{\text{ring}} = \frac{r}{3} = \frac{2.36 \times 10^{20}}{3} = 7.867 \times 10^{19} \text{ m}$$

2.2 Ring Orbital Frequency

The Keplerian orbital frequency of a test mass at r_{ring} within the total gravitational potential:

$$\omega_{\text{ring}} = \sqrt{\frac{GM}{r_{\text{ring}}^3}}$$

Substituting values:

$$\omega_{\text{ring}} = \sqrt{\frac{6.674 \times 10^{-11} \times 1.989 \times 10^{41}}{(7.867 \times 10^{19})^3}}$$

Computing the denominator:

$$r_{\text{ring}}^3 = (7.867 \times 10^{19})^3 = 4.871 \times 10^{59} \text{ m}^3$$

$$\omega_{\text{ring}} = \sqrt{\frac{1.327 \times 10^{31}}{4.871 \times 10^{59}}} = \sqrt{2.724 \times 10^{-29}} = 1.650 \times 10^{-14} \text{ rad/s}$$

2.3 Ring Orbital Period

$$T_{\text{ring}} = \frac{2\pi}{\omega_{\text{ring}}} = \frac{6.2832}{1.650 \times 10^{-14}} = 3.808 \times 10^{14} \text{ s} = 12.08 \text{ Myr}$$

This is a physically reasonable orbital period for a ring structure at $\sim 8 \times 10^1$ kpc from the galactic centre.

2.4 Proximity Enhancement Factor

The gravitational influence of the ring mass m_{ring} at the reference point r (located a distance $\Delta r = r - r_{\text{ring}} = 2r/3$ from the ring) scales as:

$$g_{\text{ring at } r} \propto \frac{G m_{\text{ring}}}{\Delta r^2} = \frac{G m_{\text{ring}}}{(2r/3)^2} = \frac{9}{4} \cdot \frac{G m_{\text{ring}}}{r^2}$$

The base term already includes the full galaxy at radius r . The ring contribution uses the ratio of effective distances:

$$\text{Proximity factor} = \left(\frac{r}{r_{\text{ring}}}\right)^2 = \left(\frac{r}{r/3}\right)^2 = 3^2 = 9$$

This gives a **9× proximity enhancement**: the ring exerts 9 times more gravitational influence per unit mass at the reference point than the galaxy average.

2.5 Ring Gravitational Perturbation Amplitude

$$A_{\text{ring}} = 9 \cdot f_{\text{ring}} \cdot g_{\text{base}}$$

where:

$$f_{\text{ring}} = 0.001 = \text{dust ring mass fraction (0.1\% of total galaxy mass)}$$

$$g_{\text{base}} = G \cdot M / r^2 = 2.382 \times 10^{-10} \text{ m/s}^2$$

$$A_{\text{ring}} = 9 \times 0.001 \times 2.382 \times 10^{-10} = 2.144 \times 10^{-12} \text{ m/s}^2$$

2.6 Full Ring Resonator Term

$$\boxed{F_{\text{ring}}(t)} = A_{\text{ring}} \cdot \cos(\omega_{\text{ring}} \cdot t)$$

This is a **pure oscillatory term** — no exponential decay, consistent with the ring's stable configuration. As the ring orbits, it imposes a periodic modulation on the vacuum field energy density at the reference point r with period $T_{\text{ring}} = 12.08 \text{ Myr}$.

3. Physical Interpretation: UQFF Gravitational Ring Resonator

3.1 Ring as Vacuum Field Oscillator

In the UQFF framework, mass concentrations modulate the vacuum energy density through their gravitational potential. A ring moving in a stable Keplerian orbit generates a time-periodic perturbation in the local UQFF field:

$$\Delta \rho_{\text{vac}}(t) = \Delta \rho_{\text{vac},0} \cdot \cos(\omega_{\text{ring}} t + \phi_0)$$

This is equivalent to a **tuned resonator** in the vacuum field — a structure that cycles energy at a characteristic frequency. The Sombrero dust ring is therefore the first identified **stable UQFF Gravitational Ring Resonator** in the catalogue.

3.2 Comparison with Orbital Periods

Object	Orbital period
Earth around Sun	1 year
Outer Milky Way disk	~220 Myr
Sombrero dust ring	12.08 Myr
Andromeda outer HI ring	~90 Myr (PAPER_275 reference)

The Sombrero ring's 12.08 Myr period places it in the intermediate-mass galaxy inner disc regime — rapid enough to generate significant temporal modulation of the UQFF field over cosmological baseline observations.

4. Module Implementation

```
// PAPER_278 – SOMBRERO_UQFF_MODULE.cpp, updateCache()
r_ring = r / 3.0; // 7.867e19 m
omega_ring = std::sqrt(G_grav * M / (r_ring * r_ring * r_ring)); // 1.650e-14 rad/s
A_ring = 9.0 * f_ring * g_base_cache; // 2.144e-12 m/s²

// Applied in computeResonantTerm():
double computeResonantTerm(double t) {
return A_ring * std::cos(omega_ring * t); // pure oscillatory – no decay
}
```

Computed values for Sombrero M104:

Quantity	Value	Units
$r_{\text{ring}} = r/3$	7.867×10^{19} ■	m
ω_{ring}	1.650×10^{-14} ■	rad/s
$T_{\text{ring}} = 2\pi/\omega_{\text{ring}}$	12.08	Myr
f_{ring}	0.001	dimensionless
Proximity factor	9.0	dimensionless
A_{ring}	2.144×10^{-12}	m/s^2
g_{BH} (PAPER279) for comparison	2.382×10^{-12}	m/s^2

Note: $A_{\text{ring}} \approx g_{\text{BH}}$ in magnitude — the ring resonance and BH contribution are comparable at leading order.

5. Key Constants Introduced

Symbol	Value	Units	Description
ω_{ring}	1.650×10^{-14} ■	rad/s	Dust Ring UQFF Orbital Resonance Frequency
r_{ring}	$r/3 = 7.867 \times 10^{19}$ ■	m	Dust ring reference radius
T_{ring}	12.08	Myr	Ring orbital period
f_{ring}	0.001	dimensionless	Dust ring mass fraction
A_{ring}	2.144×10^{-12}	m/s^2	Ring resonance amplitude
Proximity factor	9	dimensionless	$(r/r_{\text{ring}})^2 = 3^2 = 9$

6. Physical Significance

First stable UQFF ring resonator: All prior UQFF ring terms (Andromeda PAPER_275) decayed exponentially. The Sombrero dust ring is the first pure undamped oscillator, establishing a new class of UQFF boundary condition.

Observable prediction: The UQFF ring resonance produces a 12.08 Myr periodic modulation in the effective gravitational acceleration at $r = 2.36 \times 10^{20}$ m with amplitude $A_{\text{ring}} = 2.14 \times 10^{-12}$ m/s². While below direct observational reach today, this is a testable UQFF prediction for future stellar spectroscopic surveys.

Ring-BH coupling: Noting that $A_{\text{ring}} \approx g_{\text{BH}}$ (both $\sim 2.1\text{--}2.4 \times 10^{-12}$ m/s²), the ring and BH gravitational contributions are comparable at the reference radius — unique among UQFF modules where the BH term is typically sub-dominant to the 26-layer Triadic sum.

Scale-free ring resonator formula: $F_{\text{ring}} = (r/r_{\text{ring}})^2 \cdot f_{\text{ring}} \cdot g_{\text{base}} \cdot \cos(\sqrt{(GM/r_{\text{ring}}^3)} \cdot t)$ provides a general template applicable to any galaxy with a measurable equatorial ring structure.

UQFF computed: Canonical UQFF buoyancy parameter $U_{bi} = \sqrt{SSq} \sqrt{GM/r} = 5.0e-4 \sqrt{0.57 \sqrt{6.67e-11} M/r}$; for solar parameters: $U_{bi, \text{Sun}} = 5.7e-4 \sqrt{6.67e-11} \sqrt{1.99e30/(6.96e8)} = 1.47e+2$ m/s.

7. References

- PAPER275: *Andromeda HI Ring UQFF Decaying Ring Term* $A_{\text{ring}} \times \exp(-\alpha t) \times \cos(\omega t)$
 PAPER279: *Sombrero SMBH Dominance Ratio γ_{BH} and r_{SOI}* (companion paper)
 Emsellem, E. et al. (2004). MNRAS, 352, 721. (M104 structure)
 Jarrel, J. R. et al. (2011). ApJ, 739, 21. (Sombrero DM halo and dust ring observations)
 de Zeeuw, P. T. et al. (1996). MNRAS, 280, 167. (Sombrero galaxy kinematics)
 Tempel, E., & Tenjes, P. (2006). MNRAS, 371, 1269. (M104 surface photometry and ring structure)

UQFF 2.0 — $F_{\text{ring}} = A_{\text{ring}} \cos(\omega_{\text{ring}} t)$ is additive to the Triadic MUGE master equation. The stable ring resonator represents a new class of UQFF gravitational boundary condition. — Daniel T. Murphy, Session 77, March 2026.