

PAPER_311: NGC 6302 Bipolar Planetary Nebula — UQFF Wind Shock Gravitational Dominance

Author: Daniel T. Murphy

Session: 89 | Date: March 17, 2026

PAPER_311: NGC 6302 Bipolar Planetary Nebula — UQFF Wind Shock Gravitational Dominance

Subtitle: FIRST UQFF Bipolar PN Wind Shock Analysis — $\eta_{wind} = 7.127 \times 10^{-11}$; $KE/gravwell = 3.564 \times 10^{-11}$

Author: Daniel T. Murphy **Session:** 89 | **Date:** March 17, 2026 **Module:** NGC6302_UQFF_MODULE.cpp (31st C++ UQFF module) **WOLFRAM_TERM:** NGC6302_WIND_SHOCK **UQFF First:** FIRST UQFF explicit bipolar planetary nebula wind shock gravitational dominance analysis

<!-- UQFF constants: $\kappa = 5.0e-4 \text{ day}^{-1}$, [SSq] = 0.57, $M_{UQFF} = 1.43e1 \text{ TeV}$ -->

Abstract

This paper presents UQFF derivations and numerical results for: PAPER_311: NGC 6302 Bipolar Planetary Nebula — UQFF Wind Shock Gravitational Dominance. Calibration constants: $\kappa = 0.0005/\text{day}$, [SSq] = 0.57. Results validated against observational data and prior UQFF whitepaper series.

1. System Parameters

Parameter	Value	Notes
M	$3.978 \times 10^{30} \text{ kg}$	Total PN ejected mass = $2.0 M_{\text{sun}}$
r	$9.46 \times 10^{15} \text{ m}$	Half-lobe radius $\approx 1 \text{ ly}$
v_wind	$1.0 \times 10^6 \text{ m/s}$	Central star fast wind (100 km/s)
t_eject	$6.312 \times 10^{10} \text{ s}$	Bipolar lobe age (2000 yr)
ρ_{fluid}	$1.0 \times 10^{-20} \text{ kg/m}^3$	Ionized lobe gas
z	$9.5 \times 10^4 \text{ pc}$	Distance $\approx 1.2 \text{ kpc}$

2. Unique Physics

2.1 UQFF Gravitational Base

$$g_{\text{base}} = \frac{G M}{r^2} = \frac{6.6743 \times 10^{-11} \times 3.978 \times 10^{30}}{(9.46 \times 10^{15})^2}$$

$$g_{\text{base}} = \frac{2.655 \times 10^{20}}{8.949 \times 10^{31}} = \mathbf{2.967 \times 10^{-12} \text{ m/s}^2}$$

2.2 Wind Shock Acceleration (Properly Normalized)

The stellar wind ram pressure per unit length gives the characteristic wind acceleration:

$$a_{\text{wind}}(t) = \frac{v_{\text{wind}}^2}{r} \left(1 + \frac{t}{t_{\text{eject}}} \right)$$

This formulation is dimensionally consistent: $[\text{m}^2/\text{s}^2] / [\text{m}] = [\text{m}/\text{s}^2]$, representing the kinematic gradient of the wind momentum deposition over the PN lobe scale length.

At $t = 0$:

$$a_{\text{wind}}(0) = \frac{(10^5)^2 \{9.46 \times 10^{15}\}}{10^{10}} = \frac{10^{10} \{9.46 \times 10^{15}\}}{10^{10}} = \mathbf{1.057 \times 10^{-6} \text{ m/s}^2}$$

At $t = t_{\text{eject}}$ (2000 yr):

$$a_{\text{wind}}(t_{\text{eject}}) = 2 \times \frac{(10^5)^2 \{9.46 \times 10^{15}\}}{10^{10}} = \mathbf{2.114 \times 10^{-6} \text{ m/s}^2}$$

2.3 Wind-to-Gravity Dominance Ratio

$$\eta_{\text{wind}} \equiv \frac{a_{\text{wind}}(t_{\text{eject}})}{g_{\text{base}}} = \frac{2.114 \times 10^{-6}}{2.967 \times 10^{-12}} = \mathbf{7.127 \times 10^5}$$

The stellar wind shock acceleration exceeds the gravitational binding by 7.127×10^5 at $t = t_{\text{eject}}$. This explains the observed bipolar expansion kinematics: the wind force is $\sim 712,700\times$ stronger than gravity at the lobe radius, making gravitational confinement impossible and guaranteeing outward bipolar expansion.

2.4 Wind Kinetic Energy vs Gravitational Well Depth

$$\frac{\text{KE}_{\text{wind}}}{\Phi_{\text{grav}}} = \frac{v_{\text{wind}}^2 \{G M / r\}}{10^{10} \{6.6743 \times 10^{-11}\} \times 3.978 \times 10^{30} / 9.46 \times 10^{15}} = \frac{10^{10} \{2.806 \times 10^4\}}{10^{10}} = \mathbf{3.564 \times 10^5}$$

The specific kinetic energy of the stellar wind exceeds the gravitational well depth by a factor of 3.564×10^5 , confirming wind outflow is thermodynamically guaranteed regardless of the gravitational mass.

3. UQFF Pipeline Term

In the full UQFF 2.0 pipeline, the wind shock term enters as:

$$g_{\text{NGC6302}}^{\text{(wind)}} = \frac{v_{\text{wind}}^2}{r} \left(1 + \frac{t}{t_{\text{eject}}}\right)$$

This additive term is dominant over g_{base} by $\eta_{\text{wind}} \sim 10^5$, confirming that **wind dynamics, not gravity, set the kinematic structure of bipolar PNe**.

4. Astrophysical Context

NGC 6302 (the "Bug Nebula") hosts one of the hottest known white dwarf central stars ($T_{\text{eff}} \approx 200,000 \text{ K}$). The fast stellar wind (100 km/s for the slow component; up to 600 km/s for the fast component observed by HST) carves the bipolar morphology through interaction with the equatorial dust torus. The UQFF analysis quantitatively confirms that the wind kinematic energy exceeds gravitational binding by ~ 6 orders of magnitude at the 1 ly lobe scale.

5. Key Results

Quantity	Value	Unit
g_{base}	2.967×10^{-12}	m/s^2

Quantity	Value	Unit
a_wind(t=0)	1.057×10^{-1}	m/s^2
awind(reject)	2.114×10^{-1}	m/s^2
η_{wind}	7.127×10^{-1}	dimensionless
KE/grav_well	3.564×10^{-1}	dimensionless

6. Classification

UQFF First: FIRST UQFF explicit bipolar planetary nebula wind shock dominance

Scale: Stellar (PN lobe scale, ~ 1 ly)

Physics category: Wind shock / stellar wind kinematics / gravitational dominance hierarchy

Cross-references: PAPER312 (*UV radiation*), PAPER313 (torus magnetic confinement)