

PAPER<i>359 ■ G359 Galactic Center Filament: Magnetic Erosion E(t) and Negative F</i>U<i>Bi</i>i

Author: Daniel T. Murphy

Session: 97

PAPER359 ■ G359 Galactic Center Filament: Magnetic Erosion E(t) and Negative FUBii

Whitepaper Series: Star-Magic UQFF Phase 2 **Session:** 97 **Source:** gokshare31b5c807a4.txt (Supplemental Gap Analysis Block) **Classification:** FIRST UQFF treatment of a Galactic Center radio filament with negative E(t) erosion and F_mag **Author:** Daniel T. Murphy

Abstract

The G359 filament complex is a system of non-thermal radio filaments in the Galactic Center region, magnetically anchored by $B_0 = 10^{25} \text{ T}$ ordered fields threading molecular clouds. UQFF introduces a negative $E(t)$ vacuum energy erosion term for the filament environment, where $E(t) < 0$ corresponds to vacuum depletion by the ordered magnetic field. The magnetic buoyancy force per unit volume $F_{\text{mag}} = B_0^2 / (2\mu_0) V$ is computed alongside the full $F_{\text{UBi}_i} \approx -8.32 \times 10^{217} \text{ N}$.

2. Core Physics

2.1 Negative Vacuum Energy Erosion Term

For the G359 filament, the UQFF $E(t)$ vacuum energy term enters with negative sign:

$$E(t)_{\text{filament}} = -E_0 \cdot f_{\text{mag}}(B_0) \cdot t$$

This represents depletion of vacuum energy by the sustained ordered magnetic field B_0 , reducing the effective UQFF vacuum buoyancy over the filament lifetime.

2.2 Magnetic Buoyancy Force

$$F_{\text{mag}} = \frac{B_0^2}{2\mu_0} \cdot V_{\text{filament}}$$

For $B_0 = 10^{25} \text{ T}$:

$$\frac{B_0^2}{2\mu_0} = \frac{(10^{-5})^2}{2 \times 4\pi \times 10^{-7}} = \frac{10^{-10}}{8\pi \times 10^{-7}} \approx 3.98 \times 10^{-5} \text{ J/m}^3 = 3.98 \times 10^{-5} \text{ Pa}$$

For filament volume $V \sim 1048 \text{ m}^3 (100 \text{ pc}^3 \sim 1 \text{ pc}^3 \sim 1 \text{ pc})$:

$$F_{\text{mag}} \approx 3.98 \times 10^{-5} \times 10^{48} = 3.98 \times 10^{43} \text{ N}$$

2.3 Modified FUBii with Negative E(t)

$$F_{\text{UBi}_i}^{\text{filament}} = F_{\text{UBi}_i}^{\text{standard}} \cdot (1 + E(t)) \approx -8.32 \times 10^{217} \cdot (1 - E_0 t)$$

For small erosion $|E_0| \ll 1$, this produces a time-dependent weakening consistent with filament aging observations.

3. Key Values

Quantity	Formula	Value
B_0	MeerKAT measurement	10 ²⁵ T
F_mag	$B^2 V / (2 \mu_0)$	3.98 × 10 ⁴ N (filament volume)
FUBi_i	UQFF full	-8.32 × 10 ¹⁷ N
E(t) sign	Filament erosion	Negative
Distance	Galactic Center	~8.2 kpc

4. Physical Significance

The Galactic Center radio filaments have resisted unified explanation for 30+ years. UQFF proposes that their near-perpendicular-to-plane orientation reflects the alignment of ordered magnetic fields B_0 with the UQFF vacuum preferred direction. The negative E(t) erosion term is a unique first in UQFF: all earlier E(t) terms were positive (bubble expansion), but filament environments represent vacuum depletion, not expansion. This sign flip provides a new taxonomic marker for UQFF systems: positive E(t) = expanding (jets, bubbles, winds); negative E(t) = eroding (filaments, fossil lobes, dissipating relics).

5. Deduplication Note

****vs. PAPER_361 (Bubble Nebula):**** PAPER_361 uses POSITIVE E(t); PAPER_359 uses NEGATIVE E(t). This is the key distinction.

vs. SOURCE5 (Galactic Center): SOURCE5 computed the Galactic Center 5-frequency resonances; PAPER359 adds the magnetic erosion and Fmag forms.

6. Classification

Physics Territory: FIRST UQFF negative E(t) erosion in a magnetic filament system **Scale:** Galactic Center filament (100 pc) **CP Implementation:** G359FilamentGalacticCenterFUBiCalculator (CondensedPhysics4.py, Session 97)

Standard Model Comparison: Observed astrophysical data from arXiv-published surveys, SIMBAD/NED catalogs, and standard GR calculations provide the quantitative baseline; UQFF deviations are within current observational uncertainty and predict measurable signatures at future facilities.

UQFF computed: Eddington luminosity UQFF correction = $1 - [SSq] \exp(-\tau) = 1 - 5.7e-1 \exp(-2.9e-4) = 4.3e-1$; F_U at event horizon = 2.0e+18 m/s.