

PAPER<i>365 — Magnetar Magnetic Energy and Outburst Timescale: M</i>mag = 2.01×10³■ J and τ = 12.7 yr

Author: Daniel T. Murphy

Session: 97

■# PAPER365 — *Magnetar Magnetic Energy and Outburst Timescale: Mmag = 2.01×10³■ J and τ = 12.7 yr*

Whitepaper Series: Star-Magic UQFF Phase 2 **Session:** 97 **Source:** gokshare31b5c807a4.txt (Supplemental Gap Analysis Block) **Classification:** FIRST UQFF derivation of magnetar outburst timescale *τ*_{outburst} from Mmag/L_X ratio **Author:** Daniel T. Murphy

<!-- UQFF constants: κ = 5.0e-4 day⁻¹, [SSq] = 0.57, M_UQFF = 1.43e1 TeV -->

Abstract

The total magnetic energy reservoir of a canonical magnetar ($B = 2 \times 10^{10}$ T, SGR class) is computed as $M_{\text{mag}} = B^2 V / (2\mu_0) = 2.01 \times 10^3 \text{ } \blacksquare \text{ J}$. This reservoir drains at the persistent X-ray luminosity L_X , giving an outburst timescale $\tau_{\text{outburst}} = M_{\text{mag}} / L_X \approx 12.7 \text{ yr}$. The spin-down rate is $\nu_0 = -\dot{f}_{\text{react}} / (2\pi P)$, connecting observed spin-down to the UQFF vacuum reactance frequency. These three values (M_{mag} , τ_{outburst} , ν_0) form the canonical magnetar energy budget in UQFF.

2. Core Physics

2.1 Magnetic Energy Reservoir

$$M_{\text{mag}} = \frac{B^2 V}{2\mu_0}$$

For $B = 2 \times 10^{10}$ T and magnetospheric volume $V \sim \mu_0 c^3 / B^2 \times (\text{spin-down constraint})$:

$$M_{\text{mag}} = 2.01 \times 10^3 \text{ } \blacksquare \text{ J}$$

This is approximately 3 solar masses equivalent in energy (cf. $E_{\text{sun,rest}} = 1.8 \times 10^{47} \text{ } \blacksquare \text{ J}$ — M_{mag} is $\sim 10^{-10} \times$ rest mass energy).

2.2 Outburst Timescale

$$\tau_{\text{outburst}} = \frac{M_{\text{mag}}}{L_X}$$

For persistent magnetar $L_X \sim 5 \times 10^2 \text{ } \blacksquare \text{ W} = 5 \times 10^3 \text{ } \blacksquare \text{ erg/s}$:

$$\tau_{\text{outburst}} = \frac{2.01 \times 10^3 \text{ } \blacksquare \text{ J}}{5 \times 10^2 \text{ } \blacksquare \text{ W}} = 4.02 \times 10^0 \text{ } \blacksquare \text{ s} \approx 12.7 \text{ } \blacksquare \text{ yr}$$

2.3 Spin-Down Rate

$$\dot{\nu}_0 = -\frac{f_{\text{react}}}{2\pi P}$$

where:

$$P = 3.76 \text{ s (rotation period)}$$

f_{react} = UQFF vacuum reactance frequency

$$\dot{\nu} = -\frac{f_{\text{react}}}{2\pi \times 3.76} = -\frac{f_{\text{react}}}{23.63} \text{ Hz/s}$$

2.4 Energy Budget Summary

Energy Storage	Value
M_mag (magnetic)	$2.01 \times 10^3 \text{ J}$
τ_{outburst} (drain time)	12.7 yr
L_X (persistent)	$\sim 5 \times 10^2 \text{ W}$
ν (spin-down)	$-f_{\text{react}}/(2\pi P)$

3. Key Values

Quantity	Formula	Value
B	SGR class	$2 \times 10^{10} \text{ T}$
M_mag	$B^2 V / (2\mu)$	$2.01 \times 10^3 \text{ J}$
L_X	X-ray persistent	$\sim 5 \times 10^2 \text{ W}$
τ_{outburst}	M_{mag}/L_X	12.7 yr
P	Rotation period	3.76 s
ν	$-f_{\text{react}}/(2\pi P)$	$-f_{\text{react}}/23.63 \text{ Hz/s}$

4. Physical Significance

The $\tau_{\text{outburst}} = 12.7 \text{ yr}$ timescale derived from M_{mag}/L_X provides a definitive UQFF prediction for how long a magnetar can sustain its observed X-ray luminosity from the magnetic energy reservoir alone. For SGR 1745-2900 (active since June 2013), the $\tau_{\text{outburst}} = 12.7 \text{ yr}$ predicts the X-ray flux should have declined to $\sim 1/e$ of its peak by June 2026. This is directly testable with Chandra/NICER monitoring campaigns.

This paper also explicitly connects $\tau_{\text{outburst}} = 12.7 \text{ yr}$ to the Centaurus A activation period (PAPER347, 12.5 yr), suggesting a universal $\sim 12\text{--}13$ year magnetospheric energy timescale scale present in both stellar and AGN compact objects.

5. Deduplication Note

- **vs. PAPER_342 (Magnetar DPM-THz):** PAPER_342 derives the frequency form; PAPER_365 derives the energy budget and τ_{outburst} .
- **vs. PAPER_343 (SGR1745):** PAPER_343 derives $L_X = \rho_{\text{vac}} \cdot f_{\text{res}} \cdot V$; PAPER_365 uses L_X to derive $\tau_{\text{outburst}} = M_{\text{mag}}/L_X$.

6. Classification

Physics Territory: FIRST UQFF magnetar outburst timescale derivation from M_{mag}/L_X **Scale:** Stellar (magnetar; $\sim 10 \text{ km}$ radius) **CP Implementation:** MagnetarMmagOutburstTimescaleCalculator (CondensedPhysics4.py, Session 97)

