

PAPER_874

U_g3=U_i+U_m Electron Tagging and THz Circulation

CP4 Class: #458 Ug3ElectronTaggingTHzCirculationCalc
Session 200C v5.61 | Source: describe mass without using weight.txt

Abstract

U_g3 = U_i + U_m in motion. U_i individually tags each electron via the THz hole pipeline from nucleus (Point A) to electron (Point B), counteracting the strong-force to position electrons at projected quantum energy shell balance. Electron DPM uses limited U_i for coherent circulation in U_g2 imaginary orbit shell. U_g2 monitors and adjusts shell position.

Key Equations

$$\begin{aligned} F_{Ug3} &= (F_{Ui} + F_{Um} + F_{DPM_e}) * G_{geo} / r_{shell}^2 \\ F_{Ui} &= k_i * fUA'_{nuc} * nu_{THz} * R_{EB} \quad (\text{repulsive tagging}) \\ F_{Um} &= k_m * fSCm_{nuc} * nu_{res} \quad (\text{SCm-driven motion}) \\ F_{DPM_e} &= k_e * (fUA'_e * fSCm_e) * nu_{THz} \quad (\text{electron circulation}) \\ G_{geo} &= \sin(\theta) * \cos(\phi) * f(nu_{THz}) \\ E_{shell} &= c * nu_{res} * \hbar * fSCm * \sin(\theta) \quad (U_{g2} \text{ monitor}) \end{aligned}$$

Star-Magic UQFF Codebase - Session 200C - Copyright Daniel T. Murphy