

PAPER_465 — NGC 1316 “Hubble Spies Cosmic Dust Bunnies”: MUGE UQFF Merger History, AGN Jets, Star Cluster Disruption

Whitepaper Series: Star-Magic UQFF Phase 2 — Merger-Active Elliptical Galaxy **Session:** 120 (C++ module encoded) / Whitepapers created Session 122 **Source:** grok_share_dc707f5d3.txt (Doc 47 — NGC1316UQFFModule, “MUGE Hubble spies cosmic dust bunnies”) **Classification:** FIRST MUGE UQFF for NGC 1316; FIRST merger-tidal F_{env} term with spiral disruption; FIRST UQFF dust-fluid coupling for post-merger elliptical **Author:** Daniel T. Murphy **CP4 Class:** Pending (dc707f5d3 batch) **C++ Module:** NGC1316UQFFModule.h / NGC1316UQFFModule.cpp

Abstract

This paper presents the complete NGC 1316 (Fornax A) gravitational evolution model under the Master Universal Gravity Equation (MUGE) + UQFF integration. NGC 1316 is a prominent post-merger elliptical galaxy in the Fornax Cluster with prominent radio lobes (Fornax A), AGN jets, and extensive dust-lane structure formed from cannibalizing a spiral companion. The UQFF model incorporates merger history via an evolving $M_{\text{spiral}}(t)$ mass term, tidal cluster disruption ($M_{\text{cluster}} = 1 \times 10^4 M_{\odot}$), dust-lane fluid coupling ($\rho_{\text{dust}} = 1 \times 10^{-21} \text{ kg/m}^3$), AGN jet energy via Ug4 reactive decay, and dark matter halo. Result: $g_{\text{NGC1316}} = 2 \times 10^{-3} \text{ m/s}^2$ (DM/fluid dominant; repulsive terms advance framework).

2. Core Physics — PAPER_465

2.1 System Parameters

Parameter	Value	Notes
M_{total}	$5 \times 10^{11} M_{\odot}$ ($\sim 9.95 \times 10^{41} \text{ kg}$)	Post-merger mass
M_{spiral}	$1 \times 10^4 M_{\odot}$	Cannibalized spiral companion
M_{BH}	$1 \times 10^6 M_{\odot}$	Central AGN black hole
M_{cluster}	$1 \times 10^4 M_{\odot}$	Star cluster undergoing disruption
r	46 kpc ($\sim 1.42 \times 10^{21} \text{ m}$)	Effective radius
d_{spiral}	50 kpc	Original separation from spiral
ρ_{dust}	$1 \times 10^{-21} \text{ kg/m}^3$	Dust-lane fluid density
B	$1 \times 10^{-4} \text{ T}$	Enhanced AGN magnetic field
z	0.005	Fornax Cluster redshift
M_{DM}	$\sim 0.85 \times M_{\text{total}}$	Dark matter fraction

2.2 Merger-Modified Gravitational Equation

$$g_{\text{NGC1316}}(r, t) = \frac{GM_{\text{sf}}(t)}{r(t)^2} (1 + H_z t) \left(1 - \frac{B}{B_{\text{crit}}} \right) (1 + F_{\text{env}}(t))$$

$$+ \sum U_{gi} + \frac{\Lambda c^2}{3} + U_i + g_{\text{quantum}} + g_{\text{fluid}} + g_{\text{DM}}$$

2.3 Merger-Driven $F_{\text{env}}(t)$

The tidal + cluster disruption environmental force:

$$F_{\text{tidal}} = \frac{GM_{\text{spiral}}}{d_{\text{spiral}}^2}$$

$$F_{\text{cluster}} = \frac{GM_{\text{cluster}}}{r_{\text{cluster}}^2}$$

$$F_{\text{env}}(t) = F_{\text{tidal}} + F_{\text{cluster}}$$

The cannibalized spiral contributes a persistent tidal debris field, while ongoing star cluster disruption deposits angular momentum into the dust lanes.

2.4 Merger Mass Evolution

$$M_{\text{merge}}(t) = M_{\text{spiral}}(1 - e^{-t/\tau_{\text{merge}}})$$

This term tracks the gradual gravitational capture and absorption of the spiral companion debris into the NGC 1316 potential well.

2.5 Ug Sub-terms

- **Ug1** (AGN magnetic dipole): Enhanced by $B = 1 \times 10^{-10}$ T from radio lobe activity — $10 \times$ typical galaxy field
- **Ug2** (superconductor boundary): $U_{g2} = B_{\text{super}}^2 / (2\mu_0)$
- **Ug3** (external spiral gravitational pull): $U'_{g3} = GM_{\text{spiral}} / d_{\text{spiral}}^2$
- **Ug4** (AGN reactive energy): $U_{g4} = k_4 \cdot E_{\text{react}}(t)$, $E_{\text{react}} = 10^{46} e^{-0.0005t}$ — models jet energy deposition over cosmic timescales

2.6 Dust-Lane Fluid Term

$$g_{\text{fluid}} = \rho_{\text{dust}} \cdot V \cdot g_{\text{base}}$$

The dust-lane density ($\rho_{\text{dust}} = 1 \times 10^{-21}$ kg/m³) is significantly lower than typical ISM gas, reflecting the post-merger dispersed dust structure visible in Hubble ACS images.

3. Equation Summary

$$g_{\text{NGC1316}}(r, t) = \frac{GM_{\text{sf}}(t)}{r(t)^2} (1 + H_z t) (1 - B/B_{\text{crit}}) (1 + F_{\text{tidal}} + F_{\text{cluster}}) + \sum U_{gi} + \frac{\Lambda c^2}{3} + U_i + g_{\text{quantum}} + g_{\text{dust}} + g_{\text{DM}}$$

Computed Result: $g_{\text{NGC1316}} \approx 2 \times 10^{37}$ m/s² — DM/fluid dominant; repulsive Ug2 and Λ terms provide merger equilibrium; AGN Ug4 advances the reactive decay framework.

4. Physical Interpretation

- **Post-merger AGN enhancement:** $B = 1 \times 10^{-4}$ T (vs. $B = 1 \times 10^{-5}$ T for normal spirals) directly amplifies Ug1 magnetic dipole term — modeling the Fornax A radio lobe power.
 - **Dust-lane dynamics:** `_dust` term in the fluid equation captures the post-merger dust structure as a pseudo-fluid modifying the gravitational field profile.
 - **Cluster disruption:** `F_cluster` in `F_env` represents ongoing star cluster tidal stripping — a UQFF innovation modeling galactic cannibalism.
-

5. C++ Module Reference

Module: `NGC1316UQFFModule` (root-level, Session 120 from `grok_share_dc707f5d3.txt`)
Key method: `computeG(double t)` — returns total `g_NGC1316` in m/s^2 **Unique feature:** `computeMmerge(double t)` — merger mass evolution term **Integration point:** `MAIN_1_CoAnQi.cpp` multi-system validation

QS=5 — Full UQFF integration: merger `F_env`, AGN Ug4, dust-fluid coupling, post-merger elliptical parameters. *Copyright — Daniel T. Murphy. Encoded Oct 10, 2025.*