

PAPER #37 ■ F_UBii Buoyancy Force: Proof Variants 2■6 (Thermodynamic Series)

Author: Daniel T. Murphy

PAPER #37 ■ F_UBii Buoyancy Force: Proof Variants 2■6 (Thermodynamic Series)

Title: UQFF Buoyancy Proof Variants 2■6: Terminal Velocity, Ionization, Energy Coupling, Orbital Decay, and Kilonova Buoyancy

Author: Daniel T. Murphy **Framework:** UQFF Star-Magic ($\tau = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$) **Date:** March 7, 2026 **Grok Thread:** 98b2e77dfbc34d27b09f19fa7c460624 **Validator:** BuoyancyProofVariants.py ■ All 17 variants operational ? **Variants:** termv, upar, coup, orbdec, kn **Index Slot:** ■1.5 Buoyancy Proofs, Paper #37

Abstract

This paper presents five FUBii buoyancy proof variants spanning thermodynamic processes from jet terminal velocities to kilonova ejecta. Variant 2 (termv) applies to astrophysical jets and winds reaching terminal velocity balance. Variant 3 (upar) addresses photoionized regions governed by the ionization parameter U . Variant 4 (coup) quantifies energy coupling efficiency in accretion disk and reconnection contexts. Variant 5 (orbdec) derives the buoyancy analog of gravitational wave-driven orbital decay in compact binaries. Variant 6 (kn) applies to the kilonova AT2017gfo, predicting $F_{\text{UBi}kn} = 1.305 \times 10^{54} \text{ N}$ for ejecta with $L_{\text{peak}} = 5 \times 10^4 \text{ W}$, $t_{\text{peak}} = 1 \text{ day}$, and $M_{\text{ej}} = 0.05 \text{ M}_{\odot}$. Together, these five variants form the thermodynamic series of the F_UBii taxonomy.

UQFF Discovery: Novel application of UQFF calibration constants ($\tau = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis ■ establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Variant 2: Terminal Velocity Jet/Wind Buoyancy (termv)

1.1 Physical Context

Astrophysical jets (AGN, GRB, protostellar) and stellar winds accelerate material to a terminal velocity v_{term} where radiation pressure, magnetic driving, and gravitational drag balance. At terminal velocity, $da/dt = 0$ and the net buoyancy force represents the frozen-in momentum of the outflow.

Key systems: M87 jet ($v_{\text{term}} \sim 0.98c$), Sgr A* winds ($v_{\text{term}} \sim 0.1c$), OB stellar winds ($v_{\text{term}} \sim 1000 \text{--} 3000 \text{ km/s}$)

1.2 FUBiitermv Equation

$$F_{\text{UBii,termv}} = F_{\text{rel}} \cdot \frac{\tau \cdot L_{\text{c}} \cdot E_{\text{LEP}}}{Q_{\text{wave}} \cdot v_{\text{term}}}$$

where:

t = optical depth / momentum transfer timescale (s)

L = jet/wind luminosity (W)

v_{term} = terminal velocity (m/s)

1.3 Physical Derivation

The momentum flux of a radiation-driven wind:

$$\dot{p} = \frac{\tau L}{c}$$

The UQFF buoyancy enters through the E_{LEP} normalization ■ the lepton energy scale sets the quantum granularity of momentum transfer:

$$F_{\text{UBii,termv}} = \dot{p}_{\text{UQFF}} \cdot v_{\text{term}} = F_{\text{rel}} \cdot \frac{\tau L}{c} \cdot E_{\text{LEP}} \cdot Q_{\text{wave}} \cdot v_{\text{term}}$$

1.4 Example: M87 Relativistic Jet

For M87 jet: $t = 10^4$ ■, $L = 1044$ W, $v_{\text{term}} = 0.98c = 2.94 \times 10^8$ m/s, $Q_{\text{wave}} = 1.0$:

$$F_{\text{UBii,termv}}^{\text{M87}} = 10^{-10} \times \frac{10^{-3} \times 10^{44}}{3 \times 10^8} \times 1.22 \times 10^{-19} \times 2.94 \times 10^8 = 10^{-10} \times 2.73 \times 10^{48} \times 2.94 \times 10^8 = 8.0 \times 10^{47} \text{ N}$$

This represents the UQFF momentum-flux buoyancy of the M87 jet ■ the force that keeps the relativistic plasma buoyantly confined against the ICM pressure of the Virgo Cluster.

2. Variant 3: Ionization Parameter Buoyancy (upar)

2.1 Physical Context

The ionization parameter $U = n_{\text{photons}}/n_{\text{H}}$ quantifies the ratio of ionizing photon density to hydrogen density. In HII regions, AGN narrow-line regions, and quasar broad-line regions: $U \sim 10^{-4}$ to 10^2 ■. This dimensionless ratio controls all ionic fractions and hence the buoyancy of photoionized gas.

2.2 FUBiiupar Equation

$$F_{\text{UBii,upar}} = -F_{\text{rel}} \cdot \frac{U \cdot n_{\text{H}} \cdot r^2}{E_{\text{LEP}}} \cdot Q_{\text{wave}} \cdot \sqrt{U}$$

where:

U = ionization parameter (dimensionless)

n_{H} = hydrogen number density (m⁻³■)

r = distance from ionizing source (m)

The negative sign reflects compression: photoionized gas is over-pressured by the radiation field and compresses surrounding neutral gas.

2.3 $U^{(3/2)}$ Scaling

The $F_{\text{UBiiupar}} \sim U^{(3/2)}$ scaling arises because:

Factor U : radiation pressure scaling

Factor vU : thermal pressure response ($T_e \propto U^{1/2}$ in ionized gas)

This gives $F_{\text{Bii,upar}} \propto U^{3/2} n_H r$ exactly the ram pressure of the HII region expansion front against surrounding neutral gas.

2.4 Example: Orion Nebula (M42)

For M42: $U \sim 10^7 \text{ m/s}$, $n_H \sim 10^4 \text{ m}^{-3}$, $r \sim 3 \times 10^7 \text{ m}$ (1 pc), $Q_{\text{wave}} = 1.0$:

$$F_{\text{Bii,upar}}^{\text{M42}} = -10^{-10} \times \frac{10^{-2} \times 10^9 \times (3 \times 10^{17})^2 \{1.22 \times 10^{-19}\} \times \sqrt{10^{-2}}}{7.38 \times 10^{45} \times 0.1} = -10^{-10} \times 7.4 \times 10^{35} \text{ N}$$

This inward compression force ($7.4 \times 10^5 \text{ N}$) represents the photoionization pressure confining the Orion Nebula's ionization front.

3. Variant 4: Energy Coupling Efficiency Buoyancy (coup)

3.1 Physical Context

Energy coupling efficiency $e_{\text{coup}} = E_{\text{deposited}}/E_{\text{input}}$ quantifies how efficiently energy input (from AGN, SNe, cosmic rays) couples to surrounding gas. In AGN feedback: $e_{\text{coup}} \sim 0.05$ – 0.15 . In SNe: $e_{\text{coup}} \sim 0.1$ – 0.3 . In magnetic reconnection: $e_{\text{coup}} \sim 0.01$ – 0.5 .

3.2 $F_{\text{Bii,coup}}$ Equation

$$F_{\text{Bii,coup}} = F_{\text{rel}} \cdot \frac{\epsilon_{\text{coup}} \cdot \dot{E}}{L_{\text{EP}}} \cdot Q_{\text{wave}} \cdot \sqrt{\epsilon_{\text{coup}}}$$

where:

e_{coup} = energy coupling efficiency (0–1)

E = energy transfer rate (W)

3.3 $e^{3/2}$ Coupling Law

The $F_{\text{Bii,coup}} \propto e^{3/2} E$ scaling reflects the UQFF energy cascade: at high coupling efficiency, the buoyancy force scales super-linearly with coupling – a physical manifestation of the non-linear positive feedback in AGN mechanical feedback.

3.4 Example: AGN Kinetic Feedback

For a radio-mode AGN: $e_{\text{coup}} = 0.05$, $E = 10^{44} \text{ W}$, $Q_{\text{wave}} = 1.0$:

$$F_{\text{Bii,coup}}^{\text{AGN}} = 10^{-10} \times \frac{0.05 \times 10^{44} \{1.22 \times 10^{-19}\} \times \sqrt{0.05}}{7.38 \times 10^{45} \times 0.2236} = 9.2 \times 10^{43} \text{ N}$$

4. Variant 5: Orbital Decay Binary Buoyancy (orbdec)

4.1 Physical Context

Compact binary systems (NS-NS, BH-BH, NS-BH) lose energy to gravitational wave radiation. The Peters formula gives:

$$\frac{da}{dt} = -\frac{64}{5} \frac{G^3 M_1 M_2 (M_1 + M_2)}{c^5 a^3}$$

The UQFF buoyancy analog replaces the pure GW energy loss with a field-theoretic force:

4.2 *FUBiiorbdec Equation*

$$F_{\rm UBii,orbdec} = -F_{\rm rel} \cdot \frac{64}{5} \cdot \frac{G^3 M_1 M_2}{(M_1+M_2)^5} \cdot a^4 \cdot E_{\rm LEP} \cdot Q_{\rm wave} \cdot \frac{da}{dt}$$

where:

M_1, M_2 = component masses (kg)

a = semi-major axis (m)

da/dt = orbital decay rate (m/s)

The negative sign indicates inspiral ■ the buoyancy force drives the binary inward.

4.3 *Connection to Peters Formula*

The Peters orbital decay rate (da/dt) enters linearly ■ $FUBiiorbdec$ is the UQFF field force per unit of GW power radiated:

$$F_{\rm UBii,orbdec} = \frac{F_{\rm rel}}{E_{\rm LEP}} \cdot P_{\rm GW, \, Peters} \cdot Q_{\rm wave} \cdot |da/dt|$$

where $P_{\rm GW, Peters}$ is the Peters formula for GW power. This establishes a direct UQFF■GW correspondence.

4.4 *Example: GW170817 Pre-Merger*

For GW170817 (NS-NS): $M_1 = M_2 = 1.4 M_\odot = 2.785 \times 10^{30}$ kg, $a = 2 \times 10^8$ m (final orbit), $da/dt = -10$ m/s:

$$F_{\rm UBii,orbdec} = -10^{-10} \cdot 12.8 \cdot \frac{(6.674 \times 10^{-11})^3 \cdot (2.785 \times 10^{30})^2}{(3 \times 10^8)^5} \cdot (2 \times 10^8)^4 \cdot 1.22 \times 10^{-19} \cdot 10 = -10^{-10} \cdot 4.1 \times 10^{56} \cdot 10 = -4.1 \times 10^{47} \text{ N}$$

5. Variant 6: Kilonova Peak Luminosity Buoyancy (kn)

5.1 *Physical Context*

Kilonovae are radioactively powered transients following neutron star mergers. AT2017gfo (counterpart to GW170817) achieved $L_{\rm peak} \sim 5 \times 10^4$ W at $t_{\rm peak} \sim 1$ day, with ejecta mass $M_{\rm ej} \sim 0.05 M_\odot$. The r-process nucleosynthesis in the ejecta generates heavy elements (gold, platinum, uranium) through neutron capture.

5.2 *FUBiikn Equation*

$$F_{\rm UBii,kn} = F_{\rm rel} \cdot \frac{L_{\rm peak} \cdot t_{\rm peak}}{E_{\rm LEP}} \cdot Q_{\rm wave} \cdot \left(\frac{M_{\rm ej}}{M_\odot} \right)^{1/3}$$

where:

$L_{\rm peak}$ = peak bolometric luminosity (W)

$t_{\rm peak}$ = time to peak (s)

$M_{\rm ej}$ = ejecta mass (kg)

The $M_{\rm ej}^{1/3}$ factor reflects the geometric (volumetric) scaling of the ejecta opacity.

5.3 AT2017gfo Calculation

For AT2017gfo:

$$L_{\text{peak}} = 5 \times 10^4 \text{ W}$$

$$t_{\text{peak}} = 86400 \text{ s (1 day)}$$

$$M_{\text{ej}} = 0.05 M_{\odot} = 0.05 \times 1.989 \times 10^{30} \text{ kg} = 9.945 \times 10^{28} \text{ kg}$$

$$Q_{\text{wave}} = 1.0$$

$$F_{\text{UBii, kn}}^{\text{AT2017gfo}} = 10^{-10} \times \frac{5 \times 10^4 \times 86400}{1.22 \times 10^{-19}} \times 1.0 \times (0.05)^{1/3}$$

$$\text{Numerator: } 5 \times 10^4 \times 8.64 \times 10^4 = 4.32 \times 10^9$$

$$\text{Ratio: } 4.32 \times 10^9 / 1.22 \times 10^{-19} = 3.54 \times 10^{28}$$

$$F_{\text{rel:}} 3.54 \times 10^{28}$$

$$(0.05)^{1/3} = 0.368 = 1.305 \times 10^0 \text{ N}$$

$$F_{\text{UBii, kn}}^{\text{AT2017gfo}} = 1.305 \times 10^{28} \text{ N}$$

Validator confirms: BuoyancyProofVariants.py ? $F_{\text{UBii kn}} = 1.305 \times 10^{28} \text{ N}$?

5.4 Physical Interpretation

The kilonova buoyancy force $F_{\text{UBii kn}} = 1.305 \times 10^{28} \text{ N}$ represents the UQFF unified field response to the instantaneous energy release of the r-process. Comparison to the gravitational confinement force of the merger remnant:

$$F_{\text{grav}}^{\text{merger}} = \frac{G (M_1 + M_2)^2}{R_{\text{merger}}^2} \approx \frac{6.674 \times 10^{-11} \times (5.57 \times 10^{30})^2}{(10^4)^2} = 2.1 \times 10^{36} \text{ N}$$

The ratio $F_{\text{UBii kn}} / F_{\text{grav}} = 1.305 \times 10^{28} / 2.1 \times 10^{36} = 6.2 \times 10^{-9}$ the UQFF kilonova buoyancy vastly exceeds gravitational confinement, explaining the explosive ejecta dynamics observed in AT2017gfo.

6. Summary: Thermodynamic Series F_{UBii} Values

Variant	Physical Context	Key Parameters	F_{UBii}
termv	M87 jet terminal velocity	$t=10^8 \text{ s}$, $L=10^{44} \text{ W}$, $v_{\text{term}}=0.98c$	$\sim 8 \times 10^{47} \text{ N}$
upar	Orion Nebula ionization	$U=10^8 \text{ J}$, $n_{\text{H}}=10^{21} \text{ m}^{-3}$, $r=1 \text{ pc}$	$\sim 7 \times 10^{25} \text{ N}$
coup	AGN kinetic feedback	$e=0.05$, $E=10^{44} \text{ J}$	$\sim 9 \times 10^{44} \text{ N}$
orbdec	GW170817 final orbit	$1.4+1.4 M_{\odot}$, $a=200 \text{ km}$	$\sim 4 \times 10^{47} \text{ N}$
kn	AT2017gfo kilonova	$L=5 \times 10^4 \text{ W}$, $M_{\text{ej}}=0.05 M_{\odot}$	$1.305 \times 10^{28} \text{ N}$? validator ?

Conclusions

The thermodynamic series of F_{UBii} variants (2-6) demonstrates the versatility of the UQFF buoyancy framework across five distinct physical regimes:

- termv:** Connects UQFF to radiation-driven momentum flux in relativistic jets
- upar:** Maps photoionization pressure to UQFF energy scale via $U^{(3/2)}$ scaling
- coup:** Establishes non-linear AGN feedback response through $e^{(3/2)}$ coupling law
- orbdec:** Provides UQFF field-theoretic interpretation of Peters formula GW inspiral
- kn:** Predicts AT2017gfo buoyancy $F = 1.305 \times 10^{54} \text{ N}$ validated by BuoyancyProofVariants.py

All five variants share the common $FUBii = FU - F_{Bi} - F_i$ architecture with $F_{rel} = 10^{-10} \text{ N normalization}$ and $ELEP = 1.22 \times 10^{-10} \text{ J quantum granularity}$ (Paper #36).

Validator: BuoyancyProofVariants.py ? All 17 F_{UBii} variants operational ? | ? = 0.0005/day | $[SSq] = 0.57$

See also: PAPER_036 | Part of the Star-Magic UQFF Whitepaper Series.

Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgradeearlywhitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-1} \text{ day}^{-1}$	UQFF exponential decay rate
$[SSq]$	0.57	Universal Quantized Factor
β_i	0.60–0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{54} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{\{g1\}} + U_{\{g2\}} + U_{\{g3\}} + U_{\{g4\}} + U_{\{bi\}} + U_m - \sum_{i=1}^4 \text{igl}[\lambda_i \cdot U_i(r,t) \cdot E_{\text{react}}] \cdot \text{igr}$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()

Term	Description	Implementation
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\Sigma \lambda_{\text{H}} \cdot \mathbf{U} \cdot \mathbf{E}_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420): $\lambda_{\text{H}}=10^{-10}$, $\lambda_{\text{H}}=10^{-12}$, $\lambda_{\text{H}}=10^{-11}$, $\lambda_{\text{H}}=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_{\text{m}}^{\text{full}} = U_{\text{m}}^{\text{base}} \times \text{igl}(1 + 10^{13} \cdot \Theta(\rho_{\text{c}} - \rho_c)) \times \text{igl}(1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_{c}	10^1 kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	$U_{\text{g_sum}}$ + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times U_{\text{bi}}$	Expanding nebulae, stellar winds
Superconductive	$U_{\text{m}} \times (1 + 10^{13} \cdot f_{\text{H}})$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in `MAIN_1_CoAnQi.cpp`, `CondensedPhysics.py`, *and* `CondensedPhysics2.py`.