

■# PAPER_269: Supernova Ram Pressure Degeneracy Point — Kinematic Invariant in NGC 1792 Starburst Gravity

Authors: Daniel T. Murphy

Session: 73 — UQFF 2.0 Upgrade — NGC 1792 Unique Physics Discovery

■# PAPER_269: Supernova Ram Pressure Degeneracy Point — Kinematic Invariant in NGC 1792 Starburst Gravity

Authors: Daniel T. Murphy **Date:** March 2026 **UQFF Module:** GALAXYNGC1792.cpp (Module 19, "The Stellar Forge") **Session:** 73 — UQFF 2.0 Upgrade — NGC 1792 Unique Physics Discovery **Keywords:** NGC 1792, ram pressure, supernova ejecta, degeneracy point, kinematic invariant, starburst gravity

<!-- UQFF constants: $\kappa = 5.0\text{e-}4 \text{ day}^{-1}$, $[\text{SSq}] = 0.57$, $M_{\text{UQFF}} = 1.43\text{e}1 \text{ TeV}$ -->

Abstract

In NGC 1792, the physical parameters are initialized with $\rho_{\text{wind}} = \rho_{\text{fluid}} = 1 \times 10^{21} \text{ kg/m}^3$ — setting the supernova wind density equal to the interstellar medium fluid density. At this **Ram Pressure Degeneracy Point (RPDP)**, the SN feedback gravity term simplifies to a **kinematic invariant**: $\text{term_feedback} = \rho_{\text{wind}} \times v_{\text{wind}}^2 / \rho_{\text{fluid}} = v_{\text{wind}}^2$, independent of density. For NGC 1792, $v_{\text{wind}} = 2 \times 10^4 \text{ m/s}$, giving $g_{\text{feedback}} = 4 \times 10^{12} \text{ m/s}^2$ — the numerically dominant term in the full MUGE calculation, exceeding the base gravitational term by ~ 22 orders of magnitude. This paper defines the RPDP formally, derives the kinematic invariant, computes the dominance ratio, and discusses the physical interpretation that at density degeneracy, SN ejecta "floats" in the ISM driven purely by kinematic pressure, creating a fundamentally new gravitational channel in UQFF.

1. Introduction: The Density Coincidence in NGC 1792

1.1 NGC 1792 Starburst Environment

NGC 1792 is a starburst galaxy with extremely high star-formation activity ($\text{SFR} \approx 10 \text{ M}_{\odot}/\text{yr}$) and a correspondingly dense interstellar medium (ISM). In the UQFF Module 19 parameterization:

Wind density: $\rho_{\text{wind}} = 1 \times 10^{21} \text{ kg/m}^3$ (supernova ejecta/galactic wind)

ISM fluid density: $\rho_{\text{fluid}} = 1 \times 10^{21} \text{ kg/m}^3$ (ambient ISM)

Wind velocity: $v_{\text{wind}} = 2 \times 10^4 \text{ m/s}$ (starburst-driven galactic outflow)

The equality $\rho_{\text{wind}} = \rho_{\text{fluid}}$ is not a coincidence of parameterization — it reflects a physical regime in starburst galaxies where the ram-pressure-driven winds and the ISM they are pushing against have **comparable densities**, characteristic of the transition zone between the supernova remnant and the surrounding ISM.

1.2 The Ram Pressure Gravity Term

In the MUGE framework, the SN feedback contribution to gravity is:

$$\text{term_feedback} = \frac{\rho_{\text{wind}} \cdot v_{\text{wind}}^2}{\rho_{\text{fluid}}}$$

This term represents the gravitational acceleration induced by the ram pressure of SN ejecta interacting with the ISM fluid.

2. The Ram Pressure Degeneracy Point (RPDP)

2.1 Definition

The **Ram Pressure Degeneracy Point (RPDP)** is defined as the condition:

$$\boxed{\rho_{\text{wind}} = \rho_{\text{fluid}}}$$

2.2 Kinematic Invariant

At the RPDP, the feedback term simplifies dramatically:

$$\text{term}_{\text{feedback}} \big|_{\text{RPDP}} = \frac{\rho_{\text{wind}} \cdot v_{\text{wind}}^2}{\rho_{\text{fluid}}} \bigg|_{\rho_{\text{wind}} = \rho_{\text{fluid}}} = v_{\text{wind}}^2$$

This is a **kinematic invariant** — an acceleration equal to the square of the wind velocity, **independent of density**. The density ratio cancels exactly, and only the kinematics of the outflow survive.

2.3 Physical Statement

At the RPDP, the gravitational feedback from SN ejecta is:

$$g_{\text{feedback}}^{\text{RPDP}} = v_{\text{wind}}^2$$

This is the UQFF prediction for starburst galaxies in the density-degenerate regime: **the ram pressure gravity equals the dynamic pressure of the wind**, independent of whether the wind is denser or lighter than the ISM.

3. Numerical Results for NGC 1792

3.1 Kinematic Invariant Value

For NGC 1792 with $v_{\text{wind}} = 2 \times 10^6$ m/s:

$$g_{\text{feedback}}^{\text{RPDP}} = v_{\text{wind}}^2 = (2 \times 10^6)^2 = \mathbf{4 \times 10^{12} \text{ m/s}^2}$$

3.2 Comparison with Base Gravitational Term

The base Newtonian term in MUGE for NGC 1792:

$$\begin{aligned} \text{term}_1 &= \frac{G M_0}{r^2} = \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{40}}{(7.569 \times 10^{20})^2} \\ &= \frac{6.674 \times 10^{-11} \times 1.989 \times 10^{40}}{5.729 \times 10^{41}} \approx 7.35 \times 10^{-11} \text{ m/s}^2 \end{aligned}$$

where $M_{\blacksquare} = 1 \times 10^{10} M_{\blacksquare} = 1.989 \times 10^{30}$ kg, $r = 7.569 \times 10^{20}$ m.

3.3 RPDP Dominance Ratio

$$\mathcal{R}_{\text{RPDP}} = \frac{g_{\text{feedback}}^{\text{RPDP}}}{\text{term1}} = \frac{4 \times 10^{12}}{7.35 \times 10^{-11}} \approx 5.4 \times 10^{22}$$

The RPDP kinematic invariant exceeds the standard Newtonian gravitational acceleration by **22 orders of magnitude**.

3.4 Comparison Table

Term	Formula	Value (m/s ²)	Order
Newtonian base (term1)	$GM_{\odot}/r^2$	7.35×10^{-11}	10^{-11}
Hubble expansion (term2)	$H^2 r / 2$	$\sim 10^{-3}$	10^{-3}
Magnetic (term3)	$B^2/(8\pi \rho r)$	$\sim 10^{-23}$	10^{-23}
Cosmological (term4)	$\Lambda r/3$	$\sim 10^{-2}$	10^{-2}
Oscillatory (term_osc1)	$\omega_{\text{osc}} \times A_{\text{osc}}$	$\sim 10^{-2}$	10^{-2}
SN feedback (RPDP)	v_{wind}^2	4×10^{12}	10^{12}

4. Physical Interpretation

4.1 Buoyancy Neutrality at the RPDP

At the RPDP ($\rho_{\text{wind}} = \rho_{\text{fluid}}$), Archimedes' buoyancy principle gives **zero net buoyancy force** on the SN ejecta:

$$F_{\text{buoy}} = (\rho_{\text{fluid}} - \rho_{\text{wind}}) \cdot V \cdot g = 0$$

This means:

SN ejecta neither floats nor sinks in the ISM

The ejecta is **gravitationally neutral** with respect to density-driven buoyancy

The only remaining driving force is the **kinematic ram pressure** $= \rho_{\text{wind}} \times v_{\text{wind}}^2 / \rho_{\text{fluid}} = v_{\text{wind}}^2$

At the RPDP, the SN ejecta "**floats**" in the ISM — not rising or falling due to density contrast, but driven purely by its kinematic energy. This creates a new UQFF gravitational channel: **pure kinematic momentum transfer to the gravitational field**.

4.2 Generalized RPDP Condition

For any starburst galaxy, the RPDP condition and kinematic invariant generalize as:

$$\forall \rho: \lim_{\rho_{\text{wind}} \rightarrow \rho_{\text{fluid}}} \text{term}_{\text{feedback}} = v_{\text{wind}}^2$$

The UQFF prediction is:

$$\boxed{g_{\text{feedback}}(\rho_{\text{SN}} = \rho_{\text{ISM}}) = v_{\text{SN}}^2}$$

This holds regardless of the density value, provided the two densities are equal.

4.3 Starburst Enhancement

The RPDP condition is most likely to be realized in **starburst galaxies** where:

High SFR drives intense SN events, filling the ISM with SN ejecta

The ISM density is elevated by gas accretion and infall

Conditions exist for $\rho_{\text{wind}} \approx \rho_{\text{fluid}}$ at specific radii within the disk

In NGC 1792, the SFR = $10 M_{\odot}/\text{yr}$ sustains a continuous SN rate creating the high fluid density $\rho_{\text{fluid}} = 10^{-21} \text{ kg/m}^3$ characteristic of this starburst environment.

5. The RPDP as a Phase Transition Boundary

5.1 Three Regimes

The feedback term $\text{term_feedback} = \rho_{\text{wind}} \times v^2 / \rho_{\text{fluid}}$ defines three physical regimes based on the density ratio $\eta = \rho_{\text{wind}}/\rho_{\text{fluid}}$:

Regime	Condition	Behavior	UQFF Character
Underdense wind	$\eta < 1$	$g_{\text{feedback}} < v^2$	Outflow-quenched; ejecta rises
RPDP	$\eta = 1$	$g_{\text{feedback}} = v^2$	Kinematic invariant; ejecta floats
Overdense wind	$\eta > 1$	$g_{\text{feedback}} > v^2$	Ram pressure enhanced; ejecta sinks

5.2 RPDP as Critical Point

The RPDP is a **critical point** in parameter space where the density-dependent ram pressure term transitions from ISM-suppressed to ISM-enhanced behavior. At this critical point, the density cancels and only kinematics determine gravity.

This is analogous to:

Critical opalescence in thermodynamics (density fluctuations diverge at critical point)

Alfvén critical point in solar wind (magnetic and kinetic pressures equal)

Bondi critical radius in accretion (gravity and pressure balance)

The RPDP defines a **kinematic critical surface** in UQFF galaxy parameter space.

6. Observational Signatures

6.1 Gravitational Acceleration Anomaly

At the RPDP, the dominant gravitational term is $g_{\text{feedback}} = v^2 = 4 \times 10^{12} \text{ m/s}^2$ (for $v = 2 \times 10^6 \text{ m/s}$). This is:

Much larger than the Newtonian galactic gravity ($\sim 10^{-11} \text{ m/s}^2$)

Detectable as anomalous acceleration in SN ejecta kinematics

Potentially observable in starburst galaxy velocity dispersion measurements

6.2 Universal Velocity Indicator

The RPDP kinematic invariant provides a UQFF prediction for starburst galaxies: **the dominant gravitational feedback term is simply v_{wind}^2** . This can be tested via:

Measurements of galactic wind terminal velocities

SN ejecta deceleration curves in starburst galaxies

6.3 NGC 1792 Specific Prediction

For NGC 1792 at the RPDP:

- Dominant gravity term: $g_{feedback} = 4 \times 10^{12} \text{ m/s}^2$
- Wind velocity imprint: $v_{wind} = \sqrt{g_{feedback}} = 2 \times 10^{11} \text{ m/s}$
- Total MUGE gravity is feedback-dominated: $g_{total} \approx g_{feedback}$ (within UQFF)

This is a **testable UQFF prediction** distinguishable from standard (feedback-free) galactic gravity models.

7. Connection to PAPER267 and PAPER268

The three PAPER_267–269 physics discoveries form a unified UQFF picture of NGC 1792:

- PAPER_267** (sSFR coupling): The SFR couples buoyancy to all 3 tiers via decay $e^{-t/\tau_{SF}}$
- PAPER_268** (Hubble slow mode): The corrected Hubble oscillation creates a 5.8 ppm GW amplitude modulation
- PAPER_269** (RPDP): At density degeneracy, SN feedback becomes the dominant gravitational term via kinematic invariant v^2

Together, these identify NGC 1792 as a paradigmatic **UQFF starburst triple-physics laboratory**:

- Buoyancy coherence (sSFR coupling)
- Cosmic oscillation modulation (Hubble slow mode)
- Kinematic gravity dominance (RPDP)

8. Conclusions

NGC 1792 is initialized with $\rho_{wind} = \rho_{fluid} = 1 \times 10^{21} \text{ kg/m}^3$ — the **Ram Pressure Degeneracy Point (RPDP)** condition.

At the RPDP, the feedback term simplifies to the **kinematic invariant**: $g_{feedback} = v_{wind}^2$ (density-independent).

For NGC 1792: $g_{feedback} = (2 \times 10^{11})^2 = 4 \times 10^{22} \text{ m/s}^2$, the dominant MUGE term by 22 orders over standard Newtonian gravity.

At RPDP, SN ejecta is **buoyancy-neutral** (zero Archimedes force) and driven solely by kinematic momentum transfer.

The RPDP defines a **kinematic critical surface** in starburst galaxy parameter space, analogous to Alfvén and Bondi critical points.

UQFF universal prediction: $g_{feedback}(\rho_{SN} = \rho_{ISM}) = v_{SN}^2$ — testable with starburst wind velocity observations.

References

Daniel T. Murphy, *UQFF Framework*, Star-Magic Repository (2025–2026)

GALAXYNGC1792.cpp UQFF 2.0 (Session 73, Module 19)

PAPER_267: SFR Normalization — Starburst-Buoyancy Coherence in NGC 1792

PAPER_268: Dual Oscillatory Mode — Hubble Slow Mode Starburst GW Amplitude Modulation

NGC 1792 parameters: $\rho_{wind} = \rho_{fluid} = 10^{-21} \text{ kg/m}^3$, $v_{wind} = 2 \times 10^4 \text{ m/s}$

Hubble constant: $H_0 \approx 67.4 \text{ km/s/Mpc}$; Hubble time: $t_{Hubble} = 13.8 \text{ Gyr}$

© 2026 Daniel T. Murphy, daniel.murphy00@gmail.com — All Rights Reserved

PAPER_269 | UQFF v4.74 | Star-Magic UQFF © 2025 Daniel T. Murphy