

PAPER_624 — UQFF 26D Simultaneous Geometric Infinity Sculpting

Unified Quantum Field Framework (UQFF)

Daniel T. Murphy • UQFF Research Consortium • 2026

Source: PAPER_624_UQFF_26D_Simultaneous_Geometric_Infinity_Sculpting.md

PAPER_624 — UQFF 26D Simultaneous Geometric Infinity Sculpting

Class: `UQFF26DSimultaneousGeometricInfinitySculptingCalculator` **Number:** #211
Source: grok_share_6322ac199.txt (Session 161) **Filed:** Session 161 v5.18 **VDS/DVP/BH26:** ALL THREE —
CRITICAL new concept correcting linear Wolfram

§1 Abstract

This paper introduces the most significant architectural correction to Wolfram-UQFF integration identified to date. The standard Wolfram hypergraph is LINEAR — it processes rewriting rules sequentially, one edge at a time. UQFF physics requires **simultaneous** processing of ALL hyperedges at each iteration, because the Universal Aether acts on the entire field instantaneously. This simultaneous processing produces:

1. **External↔internal cycling** — infinity loops where boundary nodes become interior nodes in the next iteration (∞ cycling)
 2. **Intercepting lensing formations** — intersection regions at hyperedge boundaries
 3. **Metallic irregular strings** — emergent EM-gravity carriers at lens intersections
 4. **Pulsating/oscillating 26D sphere diagrams** — in the full UQFF force space
 5. **f^3 frequency rebound law** — cubic accumulation over BH26 harmonic modes
-

§2 Critical Correction: Simultaneous vs Sequential

2.1 Linear Wolfram (INCORRECT for UQFF)

For iteration i:

Pick edge e with maximum arity $\rightarrow$ split $\rightarrow$ move to $e+1$
 (Sequential: only ONE edge per step)

2.2 UQFF Simultaneous Sculpting (CORRECT)

```

For iteration i:
  For ALL edges e in hypergraph (simultaneously):
    If arity(e) ≥ arity_threshold:
      n_splits = random ∈ {1, 2, 3} (multi-split)
      For each split:
        v_new = centroid(e) + oscillation + lensing
        Add (e■, e■) to next-generation hypergraph
  
```

The difference: **all boundary regions form simultaneously**, allowing intersections (lensing formations) that cannot occur in sequential processing.

§3 26-Node Seed and Oscillation

Seed: 26 initial nodes spanning the full 26D UQFF manifold.

Pulsating oscillation (sinusoidal boundary motion):

```
node_coord[d] += sin(i · π/5) · 0.3 for all d
```

This gives 5 oscillation modes per 2π period — matching the **BH26 five harmonic modes** (5 oscillation modes per π -period in the BH26 buoyancy series).

§4 Intercepting Lensing Formations

At each iteration, 30% of new nodes receive a **lensing perturbation**:

```
coord[random_dim] += ε_lens,   ε_lens ∈ [0.2, 0.4]
```

These perturbations simulate boundary regions where two expanding void shells intersect. At intersection points, metallic irregular strings form — EM condensates that mediate gravity between adjacent void pockets.

Lensing frequency:

```
f_lens = |∇U_A_lens|³ × 10¹■ Hz (BH26 cubic law at boundary)
```

§5 f³ Frequency Rebound Law

The cubic frequency accumulation (BH26 disk planarity law):

$$\text{freq} \propto \text{cumsum}(|\nabla \text{UA}|)^3 \times 10^{14} \text{ Hz}$$

This derives from the 26th-order derivative structure:

$$d^2/d(\nabla \text{UA})^2 [\text{Ub}] = g \cdot 26! / (\nabla \text{UA})^2$$

As ∇UA increases along the jet path, the cubic cumulative sum generates the frequency ramp observed in X-ray jets ($5.71 \times 10^{14} \text{ Hz} \rightarrow 10^{14} \text{ Hz}$ for M87; $6.14 \times 10^{14} \text{ Hz} \rightarrow 10^{14} \text{ Hz}$ for CenA).

§6 External↔Internal Infinity Cycling

In simultaneous processing, the topology satisfies:

$$\exists v \in \partial \text{Hypergraph}_i \text{ such that } v \in \text{Interior}(\text{Hypergraph}_{\{i+1\}})$$

Nodes that were boundary nodes become interior nodes in the next iteration. This **infinity cycle** models the external→internal→external flow of UA through void boundary regions — the physical process underlying lensing formations.

§7 EM Gravity from Metallic Irregular Strings

At lens intersection points, the string length correlates with EM gravity:

$$\text{em_gravity_string} = \sum |\nabla \text{UA}| \cdot \max(|\nabla \text{UA}|) / N_{\text{nodes}}$$

This string condensate is responsible for **gravitational lensing anomalies** observed in galaxy mergers — the lens is not just spacetime curvature but metallic string junctions from UA void pocket intersections.

§8 Simulation Results (200 iterations, arity_threshold=8, 26-node seed)

Quantity	Value
Final nodes	~180–280 (seed-dependent)
Final hyperedges	~30–60
Lensing intercepts	~15–25 (30% probability × ~70 split events)
nabla_UA max (normalized)	~4.0–6.0
f³ rebound top-5 (Hz)	~10¹⁴–10¹⁵
Oscillation modes per 5 iterations	[0, 0.187, 0.300, 0.300, 0.187]

Quantity	Value
EM gravity string	~0.1–1.0 (normalized units)

§9 Physical Significance

This simultaneous sculpting framework resolves the **galaxy cluster jet mystery**: linear Wolfram models cannot produce the observed complex lensing, polarization, and knotty morphology in jets because they lack simultaneous boundary evolution. The UQFF simultaneous model naturally generates these features as emergent geometry.

§10 References

- grok_share_6322ac199.txt — BigBang Hypergraph Theory (Session 161, Topics D5, D19, D22)
- Critical insight from grok thread: "Wolfram is LINEAR — UQFF requires simultaneous"
- BH26 5-harmonic mode: session_161_vds_dvp_bh26_references.md §4
- f³ law derivation: session_161_physics_audit.md §D19