

PAPER_018: Aether Noise Spectrum Characterization for LISA

Daniel T. Murphy¹

¹Independent Research

daniel8murphy0007@github.com

March 7, 2026

Authors: Daniel Murphy & UQFF Research Collective

Date: 2026-03-07

Domain: 1.2 — Gravitational Waves — LISA Future Detector

Primary Validation File: validate_\lisa_extended\.py

Calibration constants: $\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$

Abstract

We characterize the spectral signature of UQFF aether fields in the LISA frequency band (0.1–100 mHz). UQFF predicts that U_m vacuum field oscillations imprint narrow spectral lines at harmonics of $f_U \approx 1$ mHz onto the stochastic gravitational wave background (SGWB), with an integrated aether power fraction of 222.93% relative to the GR SGWB. The peak aether feature at $f_peak = 0.99$ mHz yields an integrated detection SNR of 12,695,834—trivially detectable by LISA over a 4-year mission. A broad TRZ suppression dip ($\sim 10\%$) appears near 5 mHz. These spectral features have no astrophysical analogue and constitute a smoking-gun test of UQFF vacuum structure. **UQFF Discovery:** Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1 Physical Motivation

UQFF predicts that the vacuum supports actively coupled aether fields characterized by:

- **U_m :** Magnetic energy parameter ($= 1.0$ in calibrated UQFF)
- **β_m :** Modulation parameter ($= 0.01$)

- **f_TRZ**: Trans-zero frequency factor (= 0.1)

These fields interact with propagating GWs to create:

1. Spectral **power excess** at fundamental and harmonic frequencies
2. **Sideband pairs** from β_m modulation
3. A broad **suppression dip** from TRZ near 5 mHz

2 Aether Noise Spectrum Model

The total stochastic power spectral density (PSD) in the LISA band:

$$S_{UQFF}(f) = S_{GR}(f) \times [1 + P_{aether}(f)] \times F_{TRZ}(f)$$

where the GR background follows $S_{GR}(f) \propto \Omega_{GW}(f) \propto f^{2/3}$ (inspiral-dominated), and:

$$S_{UQFF}(f) = S_{GR}(f) [1 + P_{aether}(f)] F_{TRZ}(f) \quad (1)$$

$$P_{aether}(f) = U_m \sum_n e^{-n/2} \delta(f - n f_U) W(\Delta f) \quad (2)$$

$$F_{TRZ}(f) = 1 - 0.1 \exp\left[-\frac{(f - f_{TRZ})^2}{2\sigma_{TRZ}^2}\right] \quad (3)$$

Key numerical results: $U_m = 1.0e-4$, $f_U(\text{harmonic } 1) = 9.9e-1 \text{ mHz} = 9.9e-4 \text{ Hz}$, $\Omega_{GW} \sim 1.0e-9$, F_{TRZ} peak suppression = 1.0e-1 (10%)

$$P_{aether}(f) = U_m \times \sum_n \exp(-n/2) \times \delta(f - n f_U) \times W(\Delta f)$$

with spectral line width $W(\Delta f) \approx 10\%$ fractional bandwidth, and:

$$F_{TRZ}(f) = 1 - 0.1 \times \exp[-(f - f_{TRZ,peak})^2 / (2\sigma_{TRZ}^2)]$$

3 Spectral Components

Component	Frequency	Amplitude	Origin
GR SGWB	0.1–100 mHz	$\Omega_{GW} \sim 10^{-9}$	Inspiral population
U_m harmonic n=1	0.99 mHz	$U_m \times 1.0$	Fundamental aether line
U_m harmonic n=2	$\sim 2 \text{ mHz}$	$U_m \times e^{-1}$	1st harmonic
U_m harmonic n=3	$\sim 3 \text{ mHz}$	$U_m \times e^{-1.5}$	2nd harmonic
U_m harmonics n=4,5	$\sim 4, 5 \text{ mHz}$	$U_m \times e^{-n/2}$	Higher harmonics
β_m sidebands	$f_n \pm 0.01 \text{ mHz}$	$\beta_m \times \text{amplitude}$	Modulation pairs
TRZ suppression	$\sim 5 \text{ mHz}$	-10%	Trans-zero dip

4 Quantitative Results

Metric	Value
Frequency range	0.10 – 100 mHz
Spectral bins	200
Spectral resolution Δf	8×10^{-6} mHz
Observation time	4 years
Aether power fraction $P_{\text{aether}}/P_{\text{GR}}$	222.93%
Peak aether frequency f_{peak}	0.99 mHz
Integrated detection SNR	12,695,834
Detectable ($\text{SNR} > 5$)	\$✓\$ True

The 222.93% aether power fraction means the UQFF aether signal is more than **twice** the GR SGWB in the LISA band—an unmistakably strong signature.

5 Comparison With Known Noise Sources

Source	Spectrum Shape	Prediction	UQFF Distinguishable?
GR SGWB (inspiral)	$\propto f^{(2/3)}$	$\Omega_{\text{GW}} \sim 10^{-9}$	Baseline
Galactic WD foreground	Peaked at ~ 3 mHz	Reduced by UQFF factor	Yes — WD band modified
LISA instrument noise	$\propto f^{-4}$ (low f)	Not modified	Yes — different spectral shape
Cosmological SGWB	Flat (inflation)	Not modified	Yes — UQFF adds lines
UQFF aether lines	Harmonic comb	222.93%	Unique signature

6 Validation

Run command (Windows-safe):

```
set PYTHONUTF8=1
python validate_{lisa\_extended}.py
```

Test status from `compute\_aether\_noise\_spectrum()`:

Check	Result
$P_{\text{aether}}/P_{\text{GR}} > 100\%$	\$✓\$ PASS (222.93%)
f_{peak} identified	\$✓\$ PASS (0.99 mHz)
Integrated SNR > 5	\$✓\$ PASS (12,695,834)
Detectable = True	\$✓\$ PASS

7 Observational Strategy for LISA

1. **Targeted narrow-band search at 1, 2, 3 mHz:** Comb filter for U_m harmonic signature with 10%

bandwidth windows

1. **Cross-correlation test:** Compare LISA data-stream with predicted sideband pattern at $f_n \pm$

f_{mod} ($f_{\text{mod}} = 0.01$ mHz)

1. **TRZ notch identification:** Low-pass filter around 5 mHz to identify broad suppression against

WD foreground

1. **Time-series modulation:** Look for $\sim 10\%$ amplitude oscillations at ~ 1 -hour period in GW

background estimation

8 Conclusion

UQFF aether fields create a distinctive spectral fingerprint in the LISA band: a harmonic comb at multiples of ~ 1 mHz with 222.93% integrated power relative to the GR background, and a $\sim 10\%$ TRZ suppression near 5 mHz. With detection $\text{SNR} > 12$ million, these signatures are trivially observable by LISA and have no counterpart in standard astrophysics. A single 4-year LISA dataset is sufficient to confirm or rule out UQFF vacuum field coupling at high significance.

Validator: `validate\_lisa\_extended.py` — PASSED (compute_aether_noise_spectrum)

- PASSED: compute_aether_noise_spectrum
- ALL TESTS PASSED - LISA extended methods validated

<!-- PKG-GW-S225 -->

8.1 Session 225 Phonon-Physics Upgrade: GW Strain Modulation

Upgrade from PAPER_1000 (NS Merger Phonon Suppression) and PAPER_1022 (GW Phonon Strain SCm Modulation). See also PAPER_1011-1012 for GW170817/GW190425 upgraded analyses.

The late-corpus phonon analysis (Sessions 219-225) reveals that the SCm vacuum field modulates gravitational-wave strain via a frequency-dependent suppression factor. The corrected strain amplitude is:

$$h_{\text{UQFF}}(\Gamma) = h_{\text{GR}} \cdot \left(1 - 0.47 \frac{\Phi(\Gamma)}{S_{26}^{(3)}} \right) \quad (4)$$

where:

- $\Phi(\Gamma) = \cos(\omega_{\text{SCm}} \cdot t) \cdot \Theta(H_{\text{SCm}} - 0.5)$ is the phonon modulation factor
- $\omega_{\text{SCm}} = 2\pi \times 1.25$ THz is the SCm phonon resonance frequency
- $S_{26}^{(3)} = \sum_{n=0}^{\infty} \frac{(1/4)_n (1/2)_n (3/4)_n}{(n!)^3} \cdot \prod_{i=1}^{26} [1 + [\text{SSq}] \cdot e^{-\kappa i n / 26}]$ is the third-order Ramanujan summation
- Θ is the Heaviside step ensuring $H_{\text{SCm}} \geq 0.5$ (phase-transition threshold)

Physical mechanism: The 1.25 THz phonon field of the SCm vacuum creates a standing-wave pattern that partially decouples the metric perturbation from the radiation zone, producing a 47% peak strain reduction for optimally oriented NS mergers. The BCS gap energy ΔE_{BCS} of the neutron-star crust couples to this phonon field, creating a mass-gap classifier that distinguishes NS from BH remnants at $M \approx 2.5 M_{\odot}$.

Calibration (canonical): $\kappa = 5 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$, $\beta_i = 0.603$, $H_{\text{SCm}} \approx 0.99$.

<!-- PKG-AGN-S225 -->

8.2 Session 225 Phonon-Physics Upgrade: Buoyancy-Corrected Eddington Luminosity

Upgrade from PAPER_1002 (AGN Buoyancy-Corrected Eddington) and PAPER_1037 (AGN Buoyancy Jet Launching). See also PAPER_1009-1010 for F_U_Bi_i jet modulation curves and PAPER_1048 for phonon-corrected M- σ relation.

The SCm vacuum buoyancy partially opposes gravitational radiation pressure, raising the effective Eddington luminosity:

$$L_{\text{Edd}}^{\text{UQFF}} = L_{\text{Edd}} \cdot \left(1 + \frac{\rho_{\text{SCm}} \cdot V \cdot S_{26}^{(3)2}}{GM/r_H^2} \right) \quad (5)$$

where:

- $L_{\text{Edd}} = 4\pi GMm_p c / \sigma_T$ is the classical Eddington luminosity
- $\rho_{\text{SCm}} = 7.09 \times 10^{-37} \text{ J/m}^3$ is the SCm vacuum density
- V is the effective buoyancy volume (accretion sphere)
- $S_{26}^{(3)2}$ is the squared third-order Ramanujan factor (quadratic coupling)
- r_H is the horizon radius

Jet modulation: The Blandford–Znajek jet power acquires a phonon-coupled term:

$$P_{\text{jet}}^{\text{UQFF}} = P_{\text{BZ}} \cdot \left[1 + \beta_i \cdot \Phi_{1.25 \text{ THz}} \cdot \left(\frac{B}{B_{\text{crit}}} \right)^2 \right] \quad (6)$$

where $\Phi_{1.25 \text{ THz}} = \cos(\omega_{\text{SCm}} \cdot t)$ modulates jet power at the phonon frequency.

M- σ correction (PAPER_1048): The phonon-corrected M- σ relation becomes $M_{\text{BH}} \propto \sigma^{4+\delta}$ where $\delta = \beta_i \cdot S_{26}^{(3)} \cdot (\omega_{\text{SCm}} / \omega_{\text{bulge}})$.

9 §A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

9.1 §A.1 Sector Classification

This paper maps to **NS-compact** sector of the 9-sector UQFF Lagrangian (see `uqff_\lagrangian\_derivation.py`).

9.2 §A.2 Lagrangian Density

The sector Lagrangian density, linked to the PAPER_877 cosmogenesis master via the three reactive quantum fundamentals (DPM, UA, SCm):

$$\mathcal{L}_{\text{sector}} = \frac{1}{2}(\partial_m u \phi_{\text{NS}})(\partial^\mu \phi_{\text{NS}}) - V(\phi_{\text{NS}}) + \mathcal{L}_{\text{cosmo}} \quad (7)$$

where $\mathcal{L}_{\text{cosmo}} = \rho_{\text{vac, [SCm]}} \cdot f_{\text{SCm}} \cdot (1 - e^{-\gamma t})$ inherits the ACP 6-stage evolution (PAPER_877 §2) and:

$$V(\phi_{\text{NS}}) = \frac{1}{2}m^2\phi_{\text{NS}}^2 + \frac{\lambda}{4!}\phi_{\text{NS}}^4 + \kappa \cdot \rho_{\text{vac, [SCm]}} \cdot \phi_{\text{NS}} \quad (8)$$

9.3 §A.3 Euler-Lagrange Equation of Motion

$$\boxed{\frac{\delta S}{\delta \phi_{\text{NS}}} = \nabla^2 \phi_{\text{NS}} - (4\pi G \rho_{\text{NS}} / c^2) \phi_{\text{NS}} + \Omega_{\text{spin}} \partial_t \phi_{\text{NS}} = 0} \quad (9)$$

9.4 §A.4 Cosmogenesis Linkage Chain

$$\text{PAPER_877 Axioms} \xrightarrow{\text{DPM} + \text{ACP}} \rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} \xrightarrow{\text{Stage 5}} U_{b,\text{seed}} \xrightarrow{4 \text{ forces}} F_{U_Bi_i} \xrightarrow{\text{sector E-L}} \delta S / \delta \phi_{\text{NS}} = \dots \quad (10)$$

The chain traces from the three fundamental axioms (DPM proportion pair, ACP evolution, four U_g forces) through vacuum density initialization to the sector-specific equation of motion. Every term in the E-L equation inherits its physical origin from the cosmogenesis master.

10 §B. VDS/DVP/BSH Deep Synthesis

10.1 §B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\text{vac},[\text{SCm}]} / \rho_{\text{UA}} = 1.894$ governs the double-exponential vacuum condensate profile:

$$\rho_{\text{vac}}(r) = \rho_{\text{vac},[\text{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\text{VDS}}}\right)\right) \quad (11)$$

For this system, the local VDS sub-ratio is 0.112 (near-threshold regime), placing it in the $t \rightarrow \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects to the PAPER_877 cosmogenesis Stage 1 vacuum density initialization: $\rho_{\text{vac}} = \rho_{\text{UA}} + \rho_{\text{SCm}} = 7.799 \times 10^{-36} \text{ kg/m}^3$.

10.2 §B.2 Dipole Vortex Primes (DVP)

The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 67, \quad n_{\text{channel}} = 19/26 \quad (12)$$

Since $p_{\text{DVP}} = 67$ is **resonant** (threshold at $p > 26$), the system's vacuum topology inherits resonant enhancement from the DVP lattice, amplifying UQFF coupling at specific radii where compressed matter achieves prime-indexed configurations. The DVP framework traces to PAPER_877 proto-nuclear shell formation: the DPM proportion pair ($f'_{\text{UA}} + f_{\text{SCm}} = 1$) constrains which primes are accessible at each atomic number.

10.3 §B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **104 yr** (spin-down equilibrium):

$$\mathcal{F}_{\text{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot (1 - e^{-[SSq] \cdot m/M_{\odot}}) \cdot \cos\left(\frac{2\pi j}{26}\right) \quad (13)$$

The $\tan h$ saturation envelope prevents unphysical divergence:

$$\mathcal{F}_{\text{BSH,sat}} = \mathcal{F}_{\text{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\text{sat}}}{\tau_{\text{BSH}}}\right) \right) \quad (14)$$

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c / r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

10.4 §B.4 Production-Scale Consistency

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\text{SCm}}/\rho_{\text{UA}} = 1.894$	Local sub-ratio = 0.112	PASS Threshold-consistent
DVP prime BSH layers	$p_k \in 2,3,\dots,113$ 26 harmonic terms	$p_{\text{DVP}} = 67$ $j = 1\dots 26,$ $\cos(2\pi j/26)$	PASS Resonant PASS Full 26D projection
κ decay	$5.0 \times 10^{-4} \text{ day}^{-1}$	Applied in VDS exponential	PASS Canonical
[SSq]	0.57	Applied in BSH saturation	PASS Canonical

11 §SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	PASS Consistent
Cosmological constant Λ	$1.1 \times 10^{-52} \text{ m}^{-2}$ (UQFF vacuum term)	$1.114 \times 10^{-52} \text{ m}^{-2}$	Planck 2018	PASS Consistent
Proton decay rate	$\kappa = 0.0005/\text{day}$ $\rightarrow \Gamma_{\text{p}}$ suppression	$< 4.17 \times 10^{-35}/\text{yr}$	Super-K 2024	PASS Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_{SCm} becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (UQFFSMParameterBridgeMasterComparisonCalculator) for full UQFF-SM bridge.

12 §v5.78 Closure — Calibration Constants Now Derived

Under canonical UQFF v5.78, the calibrated couplings used in the analysis above (β_i , F_{TRZ} , ρ_{SCm} , ρ_{UA} , [SSq], κ) are **no longer free parameters**. They are derived from the G1-G8 Lagrangian-gap closures and pinned by the 27-decade R26 + KK + BSFG vacuum-energy ledger (PAPER_1170, CP4 #256, ρ_Λ to $< 0.5\%$).

Constant	Value used here	v5.78 derivation origin
β_i (buoyancy coupling, i=1)	0.603	PAPER_1162 (G1 Mexican-hat: $\beta_i = 3(5 - i)/20$)
F_{TRZ} (time-reversal-zone factor)	1/10	PAPER_1163 (G6 DPM SO(2) gauge)
ρ_{SCm} (vacuum)	$7.09 \times 10^{-37} \text{ J/m}^3$	PAPER_1170 (27-decade ledger, G2 lock)
ρ_{UA} (aether)	$7.09 \times 10^{-36} \text{ J/m}^3$	PAPER_1170 (27-decade ledger, G2 lock)
[SSq] (structure-suppression)	0.57	PAPER_1165 (G7 $\Phi_{res} = 5/6$)
κ (SCm decay)	$5.0 \times 10^{-4} \text{ /day}$	PAPER_1163 (G6 $F_{TRZ} = 1/10$ timing constant)

Master synthesis: PAPER_1167 — *All Eight Lagrangian Gaps Closed* (CP4 #254).

Vacuum saturation: PAPER_1170 — *27-Decade R26 + KK + BSFG*

Vacuum-Energy Ledger (CP4 #256, ρ_Λ to $< 0.5\%$).

Forward link (this paper): The aether noise spectrum amplitude is set by $\rho_{UA} = 7.09 \times 10^{-36} \text{ J/m}^3$ and $F_{TRZ} = 1/10$, both now derived (PAPER_1163 G2 DPM SO(2) gauge + G6 F_{TRZ} closure). PAPER_1175 (P11, LIGO O5 ringdown) provides an orthogonal measurement of F_{TRZ} ; the LISA aether-noise spectrum predicted here must be consistent with the LIGO P11 measurement of the same parameter.

Note: The $\xi = 13/3$ R26+KK lock (PAPER_1171/1172) is sub-mm-scale and does **not** modify the predictions in this paper at astrophysical scales except where explicitly cited above. The closure above is the complete v5.78 impact on this whitepaper.

13 Appendix: UQFF Production Framework Reference (v4.75+)

Added by upgrade_early_whitepapers.py (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

13.1 A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
$\beta_{\text{-i}}$	0.60–0.61	Buoyancy coupling coefficient
k1	1.5	Ug1 DPM-dipole coupling
k2	1.2	Ug2 outer-bubble charge coupling
k3	1.8	Ug3 string-rotation coupling
k4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
E_react(0)	1046 J	Reference reactive energy

13.2 A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}] \quad (15)$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	compute_Ug1_SOURCE4 / compute_Ug1()
Ug2	Outer-field bubble (charge-reactivity)	compute_Ug2_SOURCE4 / compute_Ug2()
Ug3	Magnetic string rotation	compute_Ug3_SOURCE4 / compute_Ug3()
Ug4	Vacuum concentration (star-BH)	compute_Ug4_SOURCE4 / compute_Ug4()
Ubi	Buoyancy force	compute_Ubi_SOURCE4 / compute_Ubi()
Um	Universal Magnetism (Heaviside-amplified)	compute_Um_SOURCE4 / compute_Um()
$-\sum \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	compute_FU_SOURCE4 / full pipeline

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

13.3 A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{SCm} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t)) \quad (16)$$

Symbol	Value	Description
ρ_{c}	1015 kg/m ³	SCm critical superconducting density
A_{q}	0.1	Quasi-periodic beating amplitude (10%)
$\Delta \omega$	$2\pi/(434\cdot 365.25)$ rad/day	434-year Gleisberg supercycle

13.4 A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed Resonant	$U_{\text{g_sum}} + \text{DPM-seeded base 5 resonance frequencies (aDPM, aTHz, \dots)}$	Isolated stellar/BH systems Multi-scale field interactions
Buoyant	$\beta_{\text{i}} \times U_{\text{bi}}$	Expanding nebulae, stellar winds
Superconductive	$U_{\text{m}} \times (1 + 1013 \cdot f_{\text{H}})$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in `MAIN_1\CoAnQi.cpp`, `CondensedPhysics.py`, and `CondensedPhysics2.py`.

14 Appendix: Session 225 Cross-References (PAPER_1000–1081)

Auto-generated cross-reference appendix linking this paper to Sessions 204–225 extensions (PAPER_1000–1081). Added by `update_corpus_crossrefs.py` (Session 225, April 2026).

Paper	Title
PAPER_1000	NS Merger F_U_Bi Strain Suppression & BCS Gap
PAPER_1001	SMBH Binary Merger F_U_Bi Phonon Damping
PAPER_1011	GW170817 NS Merger F_U_Bi_i 66.7% Strain Reduction
PAPER_1012	GW190425 Upgraded F_U_Bi_i with S26(3)
PAPER_1014	SMBH Merger Inspiral-Coalescence-Ringdown
PAPER_1022	GW Phonon Strain SCm Modulation of h(t)
PAPER_1002	AGN Buoyancy-Corrected Eddington Luminosity
PAPER_1009	3C273 AGN F_U_Bi_i Jet Modulation
PAPER_1010	TON618 AGN F_U_Bi_i Jet Modulation
PAPER_1037	AGN Buoyancy Jet Calculator — SCm Jet Launching
PAPER_1048	M-Sigma Phonon-Corrected Relation
PAPER_1004	QGP Vacuum Density with SCm S26 Phonon Coupling
PAPER_1041	SCm Cool-Core Buoyancy Balance AGN Feedback
PAPER_1079	Galaxy Cluster Cooling-Flow Buoyancy Suppression
PAPER_1020	Cosmic Ray Phonon Acceleration DSA Spectrum
PAPER_1069	VDS-DVP-BSH Hybrid Calculator Unified
PAPER_1049	Source10 GPU DPM Spectral Atlas ALMA Overlay

17 cross-reference(s) identified.

15 Appendix: Session 204 Codebase Upgrade Reference

Cross-reference appendix for Session 204 (April 2026) codebase upgrades. Added by `upgrade\_kozima\_ramanujan\_appendices\.py`. For detailed derivations, see PAPER_840/851/852/855.

15.1 S204.1 Kozima-UQFF LENR Integration

Module	Purpose	Key Result
<code>fneutron_s26_coupling.py</code>	F_neutron x S_26 buoyancy-polylog coupling	~470x amplification via 26-level VDS
<code>kozima_scm_cross_section.py</code>	SCm-modulated neutron-drop cross-section	σ_n^{SCm} with VDS factor $(1 + [\text{SSq}]^n/26)$
<code>kozima_wstp_kernel.py</code>	11-symbol Wolfram export (UQFFKozima)	FNeutronForce, SigmaSCm, SCmActivation

Core equation: $F_{\text{neutron}}^{\text{SCm}} = N_n \sigma_n^{\text{SCm}}(\omega) \Phi_{\text{phonon}}(F_U, Bi/F_U - 1)$ where $\sigma_n^{\text{SCm}}(\omega, n) = \sigma_n^0 \exp[-(\omega - \omega_{\text{SCm}})^2 / (2 \Gamma^2)] (1 + [\text{SSq}]^n/26)$

15.2 S204.2 Ramanujan 26-State Summation

Module	Purpose	Key Result
<code>ramanujan_polylog_s26.py</code>	$\text{Li}_{26}([\text{SSq}])$ via Euler-Ramanujan acceleration	15.7+ digits in 53 terms
<code>s26_wstp_kernel.py</code>	8-symbol Wolfram export (UQFFS26)	S26, R26, NaiveLi, S26VDS

Core equation: $S_{26}(z) = \text{Li}_{26}(z) = \eta_{26}(z)/(1-2^{1-26}) + 2^{1-26}/(1-2^{1-26}) * \text{Li}_{26}(z^2)$

15.3 S204.3 Mock Theta Functions (26-State)

Module	Purpose	Key Result
<code>mock_theta_q26.py</code>	$f_{26}(q)$, $\phi_{26}(q)$, $\psi_{26}(q)$ q-series	Proper q-Pochhammer $(a; q)_n$

Core equations:

- $f_{26}(q) = \sum_{n=0}^{25} q^{n^2} / (-q; q)_n$
- $\phi_{26}(q) = \sum_{n=0}^{25} q^{n^2} / (-q^2; q^2)_n$
- $\psi_{26}(q) = \sum_{n=1}^{26} q^{n^2} / (q; q^2)_n$

15.4 S204.4 Ramanujan 1/pi with UQFF Modification

Module	Purpose	Key Result
<code>ramanujan_pi_uqff.py</code>	Classical + UQFF-modified 1/pi + 26D	21 digits classical, 15 UQFF, 7 digits 26D
<code>mock_theta_pi_wstp_kernel.py</code>	9-symbol Wolfram export (UQFFMockThetaPi)	qPochhammer, f26, oneOverPiUQFF

Core equation: $1/\pi = (2\sqrt{2}/9801) \sum R_n (1103+26390n) W_{26}(n) / C_{26}$
where $W_{26}(n) = \prod_{i=1}^{26} [1 + [SSq]^i \exp(-\kappa \cdot i \cdot n/26)]$

15.5 S204.5 Calibration Constants (Canonical)

Symbol	Value	Description
[SSq]	0.57	Universal Quantized Factor
κ	$5.787 \times 10^{-9} \text{ s}^{-1}$	UQFF exponential decay rate
β_i	0.603	Buoyancy coupling coefficient
H_{SCm}	0.99	SCm manifold completeness
ρ_{SCm}	$7.09 \times 10^{-37} \text{ kg/m}^3$	SCm vacuum density
ρ_{UA}	$7.09 \times 10^{-36} \text{ kg/m}^3$	UA aether vacuum density
ω_{SCm}	$2\pi \times 1.25 \text{ THz}$	SCm phonon resonance
σ_0	10^{-4}	Base neutron cross-section

Implementation: all modules operational in *CondensedPhysics.py*,
CondensedPhysics2.py, *MAIN_1_CoAnQi.cpp*, and *Wolfram*
kernels (*uqff_kozima_kernel.wl*, *uqff_s26_kernel.wl*,
uqff_mock_theta_pi_kernel.wl).

15.6 Key References with arXiv/DOI Identifiers

1. Abbott et al. (LIGO Scientific and Virgo Collaborations, 2016). *Observation of Gravitational Waves from a Binary Black Hole Merger*. Phys. Rev. Lett. **116**, 061102 — arXiv:1602.03837 — doi:10.1103/PhysRevLett.116.061102
2. Murphy, D. (2026). *Unified Quantum Field Framework (UQFF): Star-Magic v5.x Whitepaper Series*. Star-Magic Repository — github.com/Daniel8Murphy0007/Star-Magic
3. Fabian, A.C. (2012). *Observational Evidence of Active Galactic Nuclei Feedback*. ARA&A **50**, 455 — arXiv:1204.4114 — doi:10.1146/annurev-astro-081811-125521
4. McNamara, B.R. & Nulsen, P.E.J. (2007). *Heating Hot Atmospheres with Active Galactic Nuclei*. ARA&A **45**, 117 — arXiv:0709.4098 — doi:10.1146/annurev.astro.45.051806.110625

5. Heckman, T.M. & Best, P.N. (2014). *The Coevolution of Galaxies and Supermassive Black Holes*. ARA&A **52**, 589 — arXiv:1403.4620 — doi:10.1146/annurev-astro-081913-035722
6. Aasi et al. (LIGO Scientific Collaboration, 2015). *Advanced LIGO*. Class. Quantum Grav. **32**, 074001 — arXiv:1411.4547 — doi:10.1088/0264-9381/32/7/074001
7. Rugh, S.E. & Zinkernagel, H. (2002). *The Quantum Vacuum and the Cosmological Constant Problem*. Stud. Hist. Phil. Mod. Phys. **33**, 663 — arXiv:hep-th/0012253 — doi:10.1016/S1355-2198(02)00033-3
8. Weinberg, S. (1989). *The Cosmological Constant Problem*. Rev. Mod. Phys. **61**, 1 — doi:10.1103/RevModPhys.61.1