

On the stability of discrete-time Markov jump linear systems

V. Daniel Camarena Pérez

National University of Engineering


vcamarenap@uni.pe

October 5, 2018

Contents

- 1 Introduction
- 2 Discrete-time Markov jump linear systems
- 3 The almost sure stability and the Lyapunov exponent
- 4 Extensions

The solar energy plant



A solar energy plant (Costa et al. 2006).

The solar energy plant

Atmospheric agents:

- Sunny day
- Cloudy day

The solar energy plant

Atmospheric agents:

- Sunny day
- Cloudy day

Three types of system design:


- ① A single control law with perturbation.
- ② Two control laws, one for each operation mode, independent.
- ③ Two control laws, one for each operation mode, dependent.

Discrete-time systems

Let the system of equations in differences

$$\begin{cases} x(k+1) = Ax(k) \\ x(0) = x_0 \end{cases} . \quad (1)$$

Discrete-time systems



Types of stability (Luenberger 1979).

Theorem

They are equivalent:

- ① $\bar{x} = 0$ is the only global asymptotically stable equilibrium point for System (1).
- ② $\rho(A) < 1$.

Theorem

They are equivalent:

- ① $\bar{x} = 0$ is the only global asymptotically stable equilibrium point for System (1).
- ② $\rho(A) < 1$.
- ③ For any $S > 0$, there exists a unique $V > 0$ such that $V - A^*VA = S$.
- ④ For some $V > 0$, we have $V - A^*VA > 0$.

Theorem

They are equivalent:


- ① $\bar{x} = 0$ is the only global asymptotically stable equilibrium point for System (1).
- ② $\rho(A) < 1$.
- ③ For any $S > 0$, there exists a unique $V > 0$ such that $V - A^*VA = S$.
- ④ For some $V > 0$, we have $V - A^*VA > 0$.
- ⑤ For some $0 < \alpha < 1 \leq \beta$, we have for all x_0 , $\|x(k)\|^2 \leq \beta \alpha^k \|x_0\|^2$, $k = 0, 1, \dots$
- ⑥ For all x_0 , we have $\sum_{k=0}^{\infty} \|x(k)\|^2 < \infty$.

Theorem

They are equivalent:

- ① $\bar{x} = 0$ is the only global asymptotically stable equilibrium point for System (1).
- ② $\rho(A) < 1$.
- ③ For any $S > 0$, there exists a unique $V > 0$ such that $V - A^*VA = S$.
- ④ For some $V > 0$, we have $V - A^*VA > 0$.
- ⑤ For some $0 < \alpha < 1 \leq \beta$, we have for all x_0 , $\|x(k)\|^2 \leq \beta \alpha^k \|x_0\|^2$, $k = 0, 1, \dots$
- ⑥ For all x_0 , we have $\sum_{k=0}^{\infty} \|x(k)\|^2 < \infty$.
- ⑦ $\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|A^k\| < 0$.

Markov chains



Random walk on \mathbb{Z} .

Markov chains

Transition probability $P = [P(i, j)]$, initial distribution ν and

$$\begin{cases} \theta(0) \sim \nu \\ \theta(k+1) | \theta(k) = i, \theta(k-1) = i_{k-1}, \dots, \theta(0) = i_0 \sim P(i, \cdot) \end{cases}$$

define a Markov chain $\{\theta(k)\}_{k \geq 0}$.

The stochastic model for the solar energy plant


- $x(k) \in \mathbb{C}^d$: state variable
- $\{A_1, \dots, A_S\}$: operation mode
- $\theta(k) \in \mathbb{S} = \{1, \dots, S\}$: transition dynamics, Markov.
- x_0, θ_0 : initial conditions, independent.

The stochastic model for the solar energy plant

- $x(k) \in \mathbb{C}^d$: state variable
- $\{A_1, \dots, A_S\}$: operation mode
- $\theta(k) \in \mathbb{S} = \{1, \dots, S\}$: transition dynamics, Markov.
- x_0, θ_0 : initial conditions, independent.

$$\begin{cases} x(k+1) = A_{\theta(k)}x(k) \\ x(0) = x_0, \theta(0) = \theta_0 \end{cases} . \quad (2)$$

The stability



Temporal evolution of the system, $d = 3$ (Costa et al. 2006).

The stability

We say that System (2) is mean square stable (**MSS**) if for any initial condition x_0, θ_0 we have

$$\mathbb{E}[x(k)x(k)^*] \rightarrow 0$$

as $k \rightarrow \infty$.

The stability

Theorem (Costa & Fragoso 1993)

They are equivalent:

- 1 *The System (2) is MSS.*
- 2 $\rho(\mathcal{A}) < 1$.

The stability

Theorem (Costa & Fragoso 1993)

They are equivalent:

- 1 *The System (2) is MSS.*
- 2 $\rho(\mathcal{A}) < 1$.
- 3 *For any $U \in \mathbb{H}^{d+}$, $U > 0$, there exists a unique $V \in \mathbb{H}^{d+}$, $V > 0$, such that $V - \mathcal{T}(V) = U$.*
- 4 *For some $V \in \mathbb{H}^{d+}$, $V > 0$, we have $V - \mathcal{T}(V) > 0$.*

The stability

Theorem (Costa & Fragoso 1993)

They are equivalent:

- ① *The System (2) is MSS.*
- ② $\rho(\mathcal{A}) < 1$.
- ③ *For any $U \in \mathbb{H}^{d+}$, $U > 0$, there exists a unique $V \in \mathbb{H}^{d+}$, $V > 0$, such that $V - \mathcal{T}(V) = U$.*
- ④ *For some $V \in \mathbb{H}^{d+}$, $V > 0$, we have $V - \mathcal{T}(V) > 0$.*
- ⑤ *For some $0 < \alpha < 1 \leq \beta$, we have for all x_0 and all θ_0 , $\mathbb{E} \left[\|x(k)\|^2 \right] \leq \beta \alpha^k \mathbb{E} \left[\|x_0\|^2 \right]$, $k = 0, 1, \dots$*
- ⑥ *For all x_0 and all θ_0 , we have $\sum_{k=0}^{\infty} \mathbb{E} \left[\|x(k)\|^2 \right] < \infty$.*

The stability

Consider that $\{\theta(k)\}_{k \geq 0}$ is an Markov chain with $\mathbb{S} = \{1, 2\}$ and transition probability

$$P = \begin{pmatrix} 0.5 & 0.5 \\ 0.5 & 0.5 \end{pmatrix}.$$

The possible operations mode

$$A_1 = \begin{pmatrix} 0 & 2 \\ 0 & 0.5 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0.5 & 0 \\ 2 & 0 \end{pmatrix}.$$

This System (2) is not **MSS**.

The stability

Consider that $\{\theta(k)\}_{k \geq 0}$ is an Markov chain with $\mathbb{S} = \{1, 2\}$ and transition probability

$$P = \begin{pmatrix} 0.5 & 0.5 \\ 0.5 & 0.5 \end{pmatrix}.$$

The possible operations mode

$$A_1 = \begin{pmatrix} 0 & 2 \\ 0 & 0.5 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0.5 & 0 \\ 2 & 0 \end{pmatrix}.$$

This System (2) is not **MSS**. Note that $\rho(A_1) = \rho(A_2) = 0.5 < 1$, i.e.,

Stability for each mode operation \nRightarrow MSS.

The stability

We say that System (2) is stochastic stable (SS) if for any initial condition x_0, θ_0 we have

$$\sum_{k=0}^{\infty} \mathbb{E} \left[\|x(k)\|^2 \right] < \infty.$$

The stability


We say that System (2) is stochastic stable (SS) if for any initial condition x_0, θ_0 we have

$$\sum_{k=0}^{\infty} \mathbb{E} \left[\|x(k)\|^2 \right] < \infty.$$

Theorem (Costa & Fragoso 1993)

If System (2) is MSS (SS) then we have for all x_0 and all θ_0 , $x(k) \rightarrow 0$ with probability one (ASS).

The almost sure stability



Almost sure stability, $d = 2$ (Costa et al. 2006).

The almost sure stability

Ergodic properties: If P irreducible then there exists stationary distribution

$$\pi P = \pi.$$

The almost sure stability

Ergodic properties: If P irreducible then there exists stationary distribution

$$\pi P = \pi.$$

The top Lyapunov exponent of System (2) is

$$\lambda = \limsup_k \frac{1}{k} \ln \|A_{\theta(k-1)} \dots A_{\theta(0)}\|.$$

The almost sure stability

Ergodic properties: If P irreducible then there exists stationary distribution

$$\pi P = \pi.$$

The top Lyapunov exponent of System (2) is

$$\lambda = \limsup_k \frac{1}{k} \ln \|A_{\theta(k-1)} \cdots A_{\theta(0)}\|.$$

Theorem (Fang et al. 1995)

If $\mathbb{E}[\lambda] < 0$ then System (2) is ASS.

The almost sure stability

Consider that $\{\theta(k)\}_{k \geq 0}$ is an irreducible Markov chain with $\mathbb{S} = \{1, 2\}$, stationary distribution $\pi = (0.5, 0.5)$ and transition probability

$$P = \begin{pmatrix} 0.5 & 0.5 \\ 0.5 & 0.5 \end{pmatrix}.$$

The possible operations mode

$$A_1 = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}.$$

This System (2) is **ASS**.

The almost sure stability

Consider that $\{\theta(k)\}_{k \geq 0}$ is an irreducible Markov chain with $\mathbb{S} = \{1, 2\}$, stationary distribution $\pi = (0.5, 0.5)$ and transition probability

$$P = \begin{pmatrix} 0.5 & 0.5 \\ 0.5 & 0.5 \end{pmatrix}.$$

The possible operations mode

$$A_1 = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}.$$

This System (2) is **ASS**. Note that $\rho(A_1) = \rho(A_2) = 2 \geq 1$, i.e.,

ASS \nRightarrow stability for each mode operation.

When the Markov chain has state space countable or Borel

- $x(k) \in \mathbb{C}^d$: state variable.
- $\{A_i : i \in \mathbb{S}\}$: operation mode.
- $\theta(k) \in \mathbb{S}$: transition dynamics, Markov.
- x_0, θ_0 : initial conditions, independent.

When the Markov chain has state space countable or Borel

- $x(k) \in \mathbb{C}^d$: state variable.
- $\{A_i : i \in \mathbb{S}\}$: operation mode.
- $\theta(k) \in \mathbb{S}$: transition dynamics, Markov.
- x_0, θ_0 : initial conditions, independent.

$$\begin{cases} x(k+1) = A_{\theta(k)}x(k) \\ x(0) = x_0, \theta(0) = \theta_0 \end{cases} . \quad (3)$$

References

- Costa, O. L. & Fragoso, M. D. (1993), 'Stability results for discrete-time linear systems with markovian jumping parameters', *Journal of mathematical analysis and applications* **179**(1), 154–178.
- Costa, O. L. V., Fragoso, M. D. & Marques, R. P. (2006), *Discrete-time Markov jump linear systems*, Springer Science & Business Media.
- Fang, Y. (1994), Stability analysis of linear control systems with uncertain parameters, PhD thesis, Case Western Reserve University.
- Fang, Y., Loparo, K. & Feng, X. (1995), 'Stability of discrete time jump linear systems', *Journal of Mathematical Systems, Estimation and Control* **5**(3), 275–321.
- Luenberger, D. G. (1979), *Introduction to dynamic systems: theory, models, and applications*, Vol. 1, Wiley New York.

iThanks!