

*"Something's just not right—our air is clean, our water is pure, we all get plenty of exercise, everything we eat is organic and free-range, and yet nobody lives past thirty."*




Learning general rules  
from experience

# Fundamental of Inductive Learning:

The Empirical Risk Minimization(ERM) rule

# Is Inductive Learning Feasible?

- Let us consider the following two examples:



$$f: \{0,1\}^3 \rightarrow \{0,1\}$$

We know  $f$  only partially in its domain

|       |   |
|-------|---|
| 0 0 0 | 0 |
| 0 0 1 | 1 |
| 0 1 0 | 1 |
| 0 1 1 | 0 |
| 1 0 0 | 1 |
| 1 0 1 | ? |
| 1 1 0 | ? |
| 1 1 1 | ? |

How is  $f$  in the last 3 elements?

# It is Not...

|       | $g$ | $f_1$ | $f_2$ | $f_3$ | $f_4$ | $f_5$ | $f_6$ | $f_7$ | $f_8$ |
|-------|-----|-------|-------|-------|-------|-------|-------|-------|-------|
| 0 0 0 | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     |
| 0 0 1 | 1   | 1     | 1     | 1     | 1     | 1     | 1     | 1     | 1     |
| 0 1 0 | 1   | 1     | 1     | 1     | 1     | 1     | 1     | 1     | 1     |
| 0 1 1 | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     |
| 1 0 0 | 1   | 1     | 1     | 1     | 1     | 1     | 1     | 1     | 1     |
| 1 0 1 | ?   | 0     | 0     | 0     | 0     | 1     | 1     | 1     | 1     |
| 1 1 0 | ?   | 0     | 0     | 1     | 1     | 0     | 1     | 0     | 1     |
| 1 1 1 | ?   | 0     | 1     | 0     | 1     | 0     | 0     | 1     | 1     |

- We can't learn this function !
- Try to verify that the eight solutions are equivalent, this is, all provide the same error.
  - Fix one of them as true solution and count how many of the others provide one, two or three errors on the unknown values.

# What then ?

- **Inductive Learning** is a hopeless approach:

In a strict sense learning out of the sample is not possible!!

( see Inductivist Turkey (Bertrand Russell) 😊 )

Is there any hope to know anything about  $f$  outside the data set **without making assumptions** about  $f$ ?

Yes, if we are willing to give up “for sure”.

Try to learn something less exigent than the proper **unknown** function,  
i.e. some useful property about the **unknown** function

# Let's try to exploit randomness....

- **NEW Hypothesis:** items inside  $\mathcal{D}$  are i.i.d samples from a probability distribution  $\mathcal{P}$
- **Consequences:**
  - $\mathcal{D}$  is the output of a random variable (vector)
  - It is not realistic to expect that every sample  $\mathcal{D}$  represent equally well the distribution  $\mathcal{P}$
  - The function  $g$  depends on  $\mathcal{D}$ , hence its election is also a random process .
- **Where is the novelty ?**
  - Probability theory shows that there are probabilistic dependencies between a random variable and a sample of it (under conditions).
    - Example : Confidence interval for the sample mean  $P(|\bar{x} - \mu| < \epsilon) > 1 - \delta, \quad \delta(\epsilon) \ll 1$

# But is probability sufficient?

- MAIN QUESTION:


There is a learning algorithm  $A$  and a sample size  $m$  such that for each distribution  $P$  if  $A$  receives  $m$  i.i.d. samples from  $P$ , is there exists a high probability that it will generate a predictor  $g$  with a low error?

- **No-Free-Lunch (NFL) Theorem (Informal):** “For every algorithm there exist a  $\mathcal{P}$  on which it fails, even though that  $\mathcal{P}$  can be successfully learned by another learner. Moreover, all algorithms are equivalent in average on all possible target functions  $f$ ”
- Conclusion: In order to succeed each learner  $(\mathcal{A}, \mathcal{H})$  must be applied on the class of distributions  $\mathcal{P}$  that it can learn.
- This highlights the need for **exploiting problem-specific knowledge** to achieve better than random performance (**inductive bias**)
  - Geometric constraint
  - Finite class  $\mathcal{H}$
  - Finite VC dimension
  - .... many others

Can we infer something outside  
the data using only  $\mathcal{D}$ ?:

The PAC answer

# Population Mean from Sample Mean



## The BIN Model

- Bin with red and green marbles.
- Pick a sample of  $N$  marbles *independently*.
- $\mu$ : probability to pick a red marble.  
 $\nu$ : fraction of red marbles in the sample.

Sample  $\longrightarrow$  the data set  $\longrightarrow \nu$   
BIN  $\longrightarrow$  outside the data  $\longrightarrow \mu$

Can we guarantee anything about  $\mu$  (**outside the data**) after observing  $\nu$  (**the data**)?

ANSWER: No. It is possible for the sample to be all green marbles and the bin to be mostly red.

Then, why do we trust polling (e.g. to predict the outcome of a presidential election).

ANSWER: The bad case is possible, but **not probable**.



# Hoeffding's Inequality

Hoeffding/Chernoff proved that, most of the time, **for a fixed  $\mu$** ,  $\nu$  cannot be too far from  $\mu$

$$\mathbb{P}(\mathcal{D}: |\mu - \nu| > \epsilon) \leq 2e^{-2\epsilon^2 N} \quad \text{for any } \epsilon > 0$$

$$\mathbb{P}(\mathcal{D}: |\mu - \nu| \leq \epsilon) \geq 1 - 2e^{-2\epsilon^2 N} \quad \text{for any } \epsilon > 0$$

**Question:** What does the value of  $\nu$  tell us on  $\mu$ :  $\mu \approx \nu \Leftrightarrow \nu \approx \mu$

Example:  $N = 1,000$ ; draw a sample and observe  $\nu$ .

$$99\% \text{ of the time} \quad \mu - 0.05 \leq \nu \leq \mu + 0.05 \quad (\epsilon = 0.05)$$

$$99.9999996\% \text{ of the time} \quad \mu - 0.10 \leq \nu \leq \mu + 0.10 \quad (\epsilon = 0.10)$$

What does this mean? If I repeatedly pick a sample of size 1,000, observe  $\nu$  and claim that

$$\mu \in [\nu - 0.05, \nu + 0.05], \quad (\text{the error bar is } \pm 0.05)$$

I will be right 99% of the time. On any particular sample you may be wrong, but not often.


We learned something. From  $\nu$ , we reached outside the data to  $\mu$ .

# Hoeffding's Inequality: Remarkable facts

- The key ingredient: **samples must be i.i.d.**
  - If the sample is constructed in some arbitrary fashion, then indeed we cannot say anything.
  - Even with independence,  $v$  can take on arbitrary values; but some values are more likely than others.
  - This is what allows us to learn something – it is likely that  $v \approx \mu$ .
- The bound  $2e^{-2\epsilon^2 N}$  does not depend on  $\mu$  or the size of the bin
  - The bin can be infinite.
  - It's great that it does not depend on  $\mu$  because  $\mu$  is unknown.
- The key player in the bound  $2e^{-2\epsilon^2 N}$  is  $N$ .
  - If  $N \rightarrow \infty$ ,  $\mu \approx v$  with very very very . . . high probability, but not for sure.
  - Can you live with  $10^{-100}$  probability of error?

$$\mathbb{P}(\mathcal{D}: |\mu - v| > \epsilon) \leq 2e^{-2\epsilon^2 N}$$

# Learning setup




In learning, the unknown is an entire function  $f$ ; in the bin it was a single number  $\mu$ .



# The Learning Error Function

The function  $h$  defines an unknown but fixed probability error  $E(h)$

Target Function  $f$



Fixed hypothesis  $h$



green:  $h(\mathbf{x}) = f(\mathbf{x})$

red:  $h(\mathbf{x}) \neq f(\mathbf{x})$

$$E(h) = \mathbb{P}_{\mathbf{x}}[h(\mathbf{x}) \neq f(\mathbf{x})]$$




(“size” of the red region)

UNKNOWN

# Relating the Bin to Learning

Let's consider all possible sample points

Now a Bin Model is defined by  $h$  and  $f$



green “marble”:  $h(\mathbf{x}) = f(\mathbf{x})$

red “marble”:  $h(\mathbf{x}) \neq f(\mathbf{x})$

BIN:  $\mathcal{X}$


$$E_{\text{out}}(h) = \mathbb{P}_{\mathbf{x}}[h(\mathbf{x}) \neq f(\mathbf{x})]$$



out-of-sample

UNKNOWN

# Relating the Bin to Learning - the Data




On the same sample, the target function  $f$  and the hypothesis  $h$  provides us with different labels

We have points in different zones of the error function.



If the sample is draw independently according to  $\mathbb{P}$ , each point will be red with probability  $\mu$  and green with probability  $1 - \mu$

# Relating the Bin to Learning - the Data

Target Function  $f$



Fixed a hypothesis  $h$



**KNOWN!**

green data:  $h(\mathbf{x}_n) = f(\mathbf{x}_n)$


red data:  $h(\mathbf{x}_n) \neq f(\mathbf{x}_n)$

$E_{\text{in}}(h)$  = fraction of red data

in-sample

misclassified

# Bin Model and Learning



Unknown  $f$  and  $P(\mathbf{x})$ , fixed  $h$

## Learning

input space  $\mathcal{X}$

green  $\mathbf{x}$  for which  $h(\mathbf{x}) = f(\mathbf{x})$


red  $\mathbf{x}$  for which  $h(\mathbf{x}) \neq f(\mathbf{x})$

$P(\mathbf{x})$

data set  $\mathcal{D}$

**Out-of-sample Error:**  $E_{\text{out}}(h) = \mathbb{P}_{\mathbf{x}}[h(\mathbf{x}) \neq f(\mathbf{x})]$

**In-sample Error:**  $E_{\text{in}}(h) = \frac{1}{N} \sum_{n=1}^N \mathbb{I}[h(\mathbf{x}) \neq f(\mathbf{x})]$



## Bin Model

Bin

green marble

red marble

randomly picking a marble

sample of  $N$  marbles

$\mu$  = probability of picking a red marble

$\nu$  = fraction of red marbles in the sample



# Hoeffding inequality in Learning

- $\mathcal{H}=\{h\}$ , **only one function**, and  $f(x)$  is the unknown true function.
- Let's  $\mathbb{I}[f(x) = h(x)]$  and  $\mathbb{I}[f(x) \neq h(x)]$  represent binary variables in the population and  $\mu = \Pr(\mathbb{I}[f(x) \neq h(x)])$
- For any training sample  $\mathcal{D}$  of size  $N$ ,  $v = \text{Fraction}(\mathbb{I}[f(x) \neq h(x)])$  on  $\mathcal{D}$
- Now  $\mu$  and  $v$  represent the population and sample error respectively.
- Let's denote by  $E_{out}(h) = \mu$  and  $E_{in}(h) = v$  the  $h$ 's global and sample error respectively
- The Hoeffding inequality can be rewritten as:

$$P(\mathcal{D}: |E_{out}(h) - E_{in}(h)| > \epsilon) \leq 2e^{-2\epsilon^2 N} \text{ for any } \epsilon > 0$$

This is called a “**Probably Aproximately Correct (PAC)**” result

- **IMPORTANT**: Note that  $\mathcal{H}$  is set before knowing the data sample.

# Hoeffding says that $E_{\text{in}}(h) \approx E_{\text{out}}(h)$

$$\mathbb{P}(\mathcal{D}: |\mu - \nu| > \epsilon) \leq 2e^{-2\epsilon^2 N}$$



$$\mathbb{P}(\mathcal{D}: |E_{\text{out}}(h) - E_{\text{in}}(h)| > \epsilon) \leq 2e^{-2\epsilon^2 N}$$

$E_{\text{in}}$  is random, but known;  $E_{\text{out}}$  fixed, but unknown.

$N \gg 1$

- If  $E_{\text{in}}(h) \approx 0 \implies E_{\text{out}}(h) \approx 0$  (with high probability), i.e.  $\mathbb{P}_{\mathcal{X}}[h(\mathbf{x}) \neq f(\mathbf{x})] = 0$ 
  - We have learned something about the entire  $f: f \approx h$  over  $\mathcal{X}$  (outside  $\mathcal{D}$ )
- If  $E_{\text{in}} \gg 0$ , we're out of luck.
  - But, we have still learned something about the entire  $f: f \approx h$  over  $\mathcal{X}$ ; it is not very useful though.

Questions:

1. Suppose that  $E_{\text{in}} = 1$ , have we learned something about the entire  $f$  that is useful?
2. What is the worst  $E_{\text{in}}$  for inferring about  $f$ ?

# Understanding PAC results

$$P(\mathcal{D}: |E_{out}(h) - E_{in}(h)| > \epsilon) \leq 2e^{-2\epsilon^2 N} \text{ for any } \epsilon > 0$$

- Let's consider  $\delta = 2e^{-2\epsilon^2 N}$  then

$$P(\mathcal{D}: |E_{out}(h) - E_{in}(h)| > \epsilon) \leq \delta \Leftrightarrow P(\mathcal{D}: |E_{out}(h) - E_{in}(h)| \leq \epsilon) > 1 - \delta$$

- Or equivalently:


$$E_{out}(h) \leq E_{in}(h) + \epsilon, \text{ with probability at least } 1 - \delta \text{ on } \mathcal{D}$$

- Let's write  $\epsilon$  as a function of  $N$  and  $\delta$ , then

$$E_{out}(h) \leq E_{in}(h) + \sqrt{\frac{1}{2N} \log \frac{2}{\delta}} \text{ with probability at least } 1 - \delta \text{ on } \mathcal{D}$$

- The higher  $N$  the narrower the interval (The sample size is important !!)
- The smaller  $\delta$  the larger the interval (The higher guarantee the lesser accuracy)

# Real Learning – Finite Learning Model



Pick the hypothesis with minimum  $E_{\text{in}}$ ; will  $E_{\text{out}}$  be small?

# The Hoeffding inequality for multiple hypothesis

NOW IS DIFFERENT: In Hoeffding's inequality  $h$  is fixed before knowing the data, BUT in REAL PROBLEMS the chosen hypothesis,  $g \in \mathcal{H}$  is identified using the data.

## SEARCH CAUSES SELECTION BIAS: THE COIN ANALOGY

**Question:** if toss one **fair coin** ten times, what is the probability that you will get ten heads ?

**Answer:**  $\approx 0.1$  (try it)

**Question:** if toss 1000 **fair coins** ten times, what is the probability that some coin will get ten heads ?

**Answer:**  $\approx 0.63$  (try it)

**Identifying coins with functions:** the higher the size of  $\mathcal{H}$  the higher the probability of having a hypothesis with  $E_{\text{in}} \approx 0$  error, BUT can we expect  $E_{\text{out}}$  to be small ?

# Guaranties for $\mathcal{H}$ -finite

- Adapting the Hoeffding's inequality to the **case of finite  $\mathcal{H}$** 
  1. Hoeffding:  **$g$  should be fixed before knowing the data sample.** (MANDATORY CONDITION)
  2. Now, the Learning Algorithm uses the training data to search for  $g$ .
  3. How to establish valid guarantees for any class element?
- A simple solution is to consider an event valid for all functions in  $\mathcal{H}$ .
  - Let  $g$  denote a generic hypothesis solution then,

$$\{\mathcal{D}: |E_{in}(g) - E_{out}(g)| > \epsilon\} = \bigcup_{h_i \in \mathcal{H}} (\mathcal{D}: |E_{in}(h_i) - E_{out}(h_i)| > \epsilon)$$

- Using  $P\left(\bigcup_{i=1:|\mathcal{H}|} B_i\right) \leq \sum_{i=1}^{|\mathcal{H}|} P(B_i)$

$$P(\mathcal{D}: |E_{in}(g) - E_{out}(g)| > \epsilon) < 2|\mathcal{H}|e^{-2\epsilon^2 N} \text{ for any } \epsilon > 0$$

**Uniform Convergence with N**

# Interpreting the bound

$$\mathbb{P}(\mathcal{D}: |E_{in}(\mathbf{g}) - E_{out}(\mathbf{g})| > \epsilon) \leq 2|\mathcal{H}|e^{-2\epsilon^2 N} \quad \text{for any } \epsilon > 0$$

$$\mathbb{P}(\mathcal{D}: |E_{in}(\mathbf{g}) - E_{out}(\mathbf{g})| \leq \epsilon) \geq 1 - 2|\mathcal{H}|e^{-2\epsilon^2 N} \quad \text{for any } \epsilon > 0$$

Generalization  
bar error

**Result:** With probability at least  $1 - \delta$ ,  $E_{out}(\mathbf{g}) \leq E_{in}(\mathbf{g}) + \sqrt{\frac{1}{2N} \ln \frac{2|\mathcal{H}|}{\delta}}$

To do that, denote  $\delta = 2|\mathcal{H}|e^{-2\epsilon^2 N}$  and writes  $\epsilon$  as a function of  $N$ ,  $\delta$  and  $|\mathcal{H}|$

$$E_{in} \text{ reaches outside to } E_{out} \text{ when } |\mathcal{H}| \text{ is small} \quad E_{out}(\mathbf{g}) \leq E_{in}(\mathbf{g}) + \mathcal{O}\left(\sqrt{\frac{\ln |\mathcal{H}|}{N}}\right)$$

if  $N \gg \ln |\mathcal{H}|$ , then  $E_{out}(\mathbf{g}) \approx E_{in}(\mathbf{g})$

- This bound does not depend on  $\mathcal{X}$ ,  $\mathbb{P}(\mathbf{x})$ ,  $f$  or how  $\mathbf{g}$  is found. (i.e. a worst case bound)
- Only requires  $\mathbb{P}(\mathbf{x})$  to generate the data points independently *including* the test point.

# A particular case: The Realizability Hypothesis

- **Definition ( $\mathcal{H}$  -realizable):** There exist  $h^* \in \mathcal{H}$  s.t.  $E_{out_{\mathcal{P},f}}(h^*) = 0$
- This means that the class of functions  $\mathcal{H}$  includes at least a function with error zero for any  $\mathcal{P}$  and  $f$ .
- Being a mathematical hypothesis can be applied in real tasks.
  - Example: Separated classes in classification
- Under this hypothesis can be shown that the ERM rule on finite classes  $\mathcal{H}$  always provides a function  $h_S$  s.t.  $E_{in}(h_S) = 0$ , obtaining the smallest error bound

$$E_{out}(h_S) \leq \frac{1}{N} \log \frac{|\mathcal{H}|}{\delta} \quad \text{with probability at least } 1 - \delta$$



# Formal Definition of PAC-Learning

- A hypothesis class  $\mathcal{H}$  is PAC learnable if there exist a function  $m_{\mathcal{H}}(\epsilon, \delta) \rightarrow \mathbb{N}$  and a learning algorithm  $\mathcal{A}$  with the following property:
  - For every,  $\epsilon, \delta \in (0, 1)$
  - For every distribution  $\mathcal{P}$  over  $\mathcal{X}$
  - For every labeling function  $f$if the realizable assumption holds with respect to  $\mathcal{H}, \mathcal{P}, f$  then when running the learning algorithm on  $N \geq m_{\mathcal{H}}(\epsilon, \delta)$  i.i.d samples generated by  $\mathcal{P}$  and labeled by  $f$ , the algorithm returns a hypothesis  $h$  such that , with probability at least  $1 - \delta$  (over the choice of the  $m$  training samples)

$$ERM_{\mathcal{P}, f}(h) \leq \epsilon$$

This means the learning algorithm always return a hypothesis  $h$  with error close to zero.

(*Understanding Machine Learning*, Shalev-Shwartz and Ben-David, 2014, free pdf)

# Sample Complexity in PAC-Learnability

- How many examples are required to guarantee a PAC solution in realizable classes?
- This depend on  $\epsilon$ ,  $\delta$ ,  $\mathcal{H}$  and the loss function range
- If  $\mathcal{H}$  is a PAC-realizable, there exist many functions  $m_{\mathcal{H}}(\epsilon, \delta)$  that satisfy the requirements given in the definition of Realizable PAC learnability
- The sample complexity for learning  $\mathcal{H}$  is defined as the “minimal integer”  $m_{\mathcal{H}}(\epsilon, \delta)$  that satisfies the requirements of realizable PAC learning with accuracy  $\epsilon$  and confidence  $\delta$
- Result for loss function with range in  $[0,1]$ : every realizable finite hypothesis class is PAC learnable with sample complexity

$$m_H(\epsilon, \delta) \leq \left\lceil \frac{1}{\epsilon} \log \frac{|\mathcal{H}|}{\delta} \right\rceil$$

# Formal Definition of Agnostic PAC-Learning

- A hypothesis class  $\mathcal{H}$  is agnostic PAC learnable if there exist a function  $m_{\mathcal{H}}(\epsilon, \delta) \rightarrow \mathbb{N}$  and a learning algorithm  $\mathcal{A}$  with the following property:
  - For every,  $\epsilon, \delta \in (0, 1)$
  - For every distribution  $\mathcal{P}$  over  $\mathcal{X}$

When running the learning algorithm on  $N \geq m_{\mathcal{H}}(\epsilon, \delta)$  i.i.d samples generated by  $\mathcal{P}$ , the algorithm returns a hypothesis  $h$  such that , with probability at least  $1 - \delta$  (over the choice of the  $m$  training samples)

$$E_{in \mathcal{P}}(h) \leq \min_{h' \in \mathcal{H}} E_{in \mathcal{P}}(h') + \epsilon$$

This means the learning algorithm always return a hypothesis  $h$  closer to the best possible inside of the class  $\mathcal{H}$

**The ERM rule is a succesful agnostic PAC learner for finite classes  $\mathcal{H}$**

# Agnostic-PAC Sample Complexity

- How many examples are required to guarantee a PAC solution?
- This depend on  $\epsilon$ ,  $\delta$ ,  $\mathcal{H}$  and the loss function range
- If  $\mathcal{H}$  is PAC learnable, there exist many functions  $m_{\mathcal{H}}(\epsilon, \delta)$  that satisfy the requirements given in the definition of PAC learnability
- The sample complexity of learning  $\mathcal{H}$  is defined as the “minimal integer”  $m_{\mathcal{H}}(\epsilon, \delta)$  that satisfies the requirements of PAC learning with accuracy  $\epsilon$  and confidence  $\delta$
- Result for loss function with range in  $[0,1]$ : every finite hypothesis class is PAC learnable with sample complexity

$$m_H(\epsilon, \delta) \leq \left\lceil \frac{2}{\epsilon^2} \log \frac{2|\mathcal{H}|}{\delta} \right\rceil$$

- Compare this new expression with that in the PAC-realizable case!

# Feasibility of Learning vs Complexity

- Learning is possible in a probabilistic setting (under conditions):
  - Samples from  $\mathcal{X}$  **must be i.i.d**
  - Same probability distribution in training and test
- To be succesful in learning means to find a function  $g$ , s.t.  $E_{out}(g) \approx 0$
- Nevertheless, we are only able to guarantee,

$$P(\mathcal{D}: |E_{in}(g) - E_{out}(g)| > \epsilon) < 2|\mathcal{H}|e^{-\epsilon^2 N} \text{ for any } \epsilon > 0$$

- Feasibility of Learning must answer two questions:
  1. Can we make sure that  $E_{out}(g)$  is close enough to  $E_{in}(g)$  ?
  2. Can we make  $E_{in}(g)$  small enough?
- What is the relationship between Feasibility of Learning and the complexity of  $\mathcal{H}$  and  $f$ ?

# Feasibility of learning : $E_{out} \approx 0$

Two conditions:


(1)  $E_{in} \approx E_{out}$   $\rightarrow$  Is verified thanks to the Hoeffding's inequality

(2)  $E_{in} \approx 0$   $\rightarrow$  Is achieved through the learning algorithm

Together, these ensure  $E_{out} \approx 0$

**BUT there is a tradeoff on  $\mathcal{H}$ :**

- **Small**  $|\mathcal{H}| \Rightarrow E_{in} \approx E_{out}$
- **Large**  $|\mathcal{H}| \Rightarrow E_{in} \approx 0$  is more likely



**What about the UNKNOWN complexity of  $f$ :**

- **Simple**  $f \Rightarrow$  can use small  $\mathcal{H}$  to get  $E_{in} \approx 0$  (need **smaller**  $N$ ).
- **Complex**  $f \Rightarrow$  need large  $\mathcal{H}$  to get  $E_{in} \approx 0$  (need **larger**  $N$ ).

# The Fundamental theorem of PAC learning

Let  $\mathcal{H}$  be a hypothesis class of functions from a domain  $\mathcal{X}$  to  $[0,1]$  and let the loss functions be the 0-1 loss. Then the following are equivalent:

- $\mathcal{H}$  has the uniform convergence properties
- Any ERM rule is a successful agnostic PAC learner for  $\mathcal{H}$
- $\mathcal{H}$  is agnostic PAC learnable
- $\mathcal{H}$  is PAC learnable
- Any ERM rule is a successful PAC learner for  $\mathcal{H}$
- $\mathcal{H}$  has a finite VC-dimension