


COOK ISLANDS






MOPT	TALITY [†]	NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MOK	Premature mortality		Total NCD deaths	2016			
	from NCDs	X	Risk of premature death between 30-70 years (%)	2016			
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	18	4	11
X	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	11	30	21
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	***		
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	30	21	26
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	24	19	21
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	27	26	27
i.	Obesity	✓	Obesity, adults aged 18+ (%)	2016	52	59	55
			Obesity, adolescents aged 10-19 (%)	2016	32	29	30
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
b	Household air pollution	-	Population with primary reliance on polluting fuels and technologies $(\!\%\!)$	2016	-	-	16
SELEC	CTED ADULT RISK FACTO	OR TRENDS					
	CURRENT TOBACO	CO SMOKING	OBESITY	F	RAISED BLOOD PRESSURE		
70			70	70			









Drug therapy to prevent heart attacks and strokes





Number of essential NCD technologies reported as



Essential NCD medicines and basic technologies to treat major NCDs