RWANDA

RISK OF PREMATURE DEATH DUE TO NCDS (%)*

Probability (%) of premature death

Past trends | Projected linear trends | Global targets
Males
Females

PROPORTIONAL MORTALITY*

14% Cardiovascular diseases
13% Other NCDs
13% Cancers
42% Communicable, maternal, perinatal and nutritional conditions
3% Chronic respiratory diseases
14% Injuries
2% Diabetes

NCDs are estimated to account for **44%** of all deaths.

9 100 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"

	NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*						
Premature mortality from NCDs	X	Total NCD deaths	2016	14 600	14 500	29 100
		Risk of premature death between 30-70 years (%)	2016	20	17	18
Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	7
RISK FACTORS						
Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	16	3	9
Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	10	17	13
Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	4	4	4
Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	18	4	11
Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	19	20	20
Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	3	3	3
Obesity	X	Obesity, adults aged 18+ (%)	2016	2	8	5
		Obesity, adolescents aged 10-19 (%)	2016	0	2	1
Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	4
Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	>95

SELECTED ADULT RISK FACTOR TRENDS

CURRENT TOBACCO SMOKING

% of the population

OBESITY

% of the population

RAISED BLOOD PRESSURE

% of the population

Past trends | Projected linear trends | Global targets
Males
Females

NATIONAL SYSTEMS RESPONSE

Drug therapy to prevent heart attacks and strokes	X	Proportion of population at high risk for CVD or with existing CVD (%)	-	...
		Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-	...
		Proportion of primary health care centres reported as offering CVD risk stratification	2017	None
		Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017	Yes
Essential NCD medicines and basic technologies to treat major NCDs	X	Number of essential NCD medicines reported as "generally available"	2017	10 out of 10
		Number of essential NCD technologies reported as "generally available"	2017	5 out of 6

... = no data available
* The mortality estimates for this country have a high degree of uncertainty because they are not based on any national NCD mortality data (see Explanatory Notes)