

BANGLADESH


2016 TOTAL POPULATION: 163 000 000 **2016 TOTAL DEATHS: 856 000**




66 400 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
	Premature mortality	X	Total NCD deaths	2016	306 700	265 900	572 600
	from NCDs	A	Risk of premature death between 30-70 years (%)	2016	23	20	22
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	6
RISK FACTORS							
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	0	0	0
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	15	37	26
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	9	9	9
\odot	Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	44	1	23
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	22	20	21
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	9	7	8
	Obesity	X	Obesity, adults aged 18+ (%)	2016	2	5	3
	Obesity	X	Obesity, adolescents aged 10-19 (%)	2016	2	2	2
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	6
4	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	82

SELECTED ADULT RISK FACTOR TRENDS








NATIONAL SYSTEMS RESPO







Proportion of population at high risk for CVD or with existing CVD (%) Proportion of high risk persons receiving any drug therapy






Reported having CVD guidelines that are utilized in at least 50% of health facilities









²⁰¹⁷ 6 out of 6

⁼ no data available