

2016 TOTAL POPULATION: 6 345 000 **2016 TOTAL DEATHS: 42 000**


23% 27% 50 Cardiovascular Other NCDs Probability (%) of premature death 40 diseases 16% 11% 30 Cancers Communicable, 20 NCDs are maternal, perinatal 3% and nutritional estimated to <u>.....</u> 10 account for 74% conditions Chronic of all deaths. respiratory 0 15% diseases 2000 2005 2010 2015 2020 2025 Injuries > 5% Past trends Projected linear trends Global targets Males0 Diabetes Females

2 700 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
\Box	Premature mortality	X	Total NCD deaths	2016	16 000	15 100	31 100
	from NCDs	А	Risk of premature death between 30-70 years (%)	2016	15	13	14
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	14
RISK FACTORS							
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	7	1	4
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016			
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	9	8	8
\odot	Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	19	3	10
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	18	16	17
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	8	10	9
Å	Obesity	X	Obesity, adults aged 18+ (%)	2016	17	27	23
			Obesity, adolescents aged 10-19 (%)	2016	9	11	10
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2
	Household air pollution	-	Population with primary reliance on polluting fuels and technologies $(\!\%\!)$	2016	-	-	14



SELECTED ADULT RISK FACTOR TRENDS







to treat major NCDs



Proportion of population at high risk for CVD or with existing CVD (%) Proportion of high risk persons receiving any drug therapy

and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Number of essential NCD medicines reported as "generally available"

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD technologies reported as

2017 Less than 25% 2017 Yes 2017 7 out of 10

5 out of 6

2017

⁼ no data available