

**2016 TOTAL POPULATION:** 540 000 **2016 TOTAL DEATHS: 2 900** 

## 36% 15% 50 Cardiovascular Other NCDs Probability (%) of premature death 40 diseases ▶ 20% 11% 30 Cancers Communicable, 20 NCDs are maternal, perinatal 4% and nutritional estimated to 10 account for 70% conditions Chronic of all deaths. respiratory 0 10% diseases 2000 2005 2010 2015 2020 2025 Injuries > 5% Past trends Projected linear trends Global targets Males Diabetes .....ō Females

3 500 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"							
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORT	TALITY*						
	Premature mortality from NCDs	X	Total NCD deaths	2016	930	1100	2 100
			Risk of premature death between 30-70 years (%)	2016	19	16	17
P	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	11
RISK	FACTORS						
	Harmful use of alcohol	<b>✓</b>	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	10	2	6
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	12	24	19
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	9	8	8
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	16	2	9
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	26	25	25
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	7	7	7
À	Obesity	<b>√</b>	Obesity, adults aged 18+ (%)	2016	6	15	11
			Obesity, adolescents aged 10-19 (%)	2016	2	3	3
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	3
	Household air pollution	-	Population with primary reliance on polluting fuels and technologies $(\!\%\!)$	2016	-	-	29









Males


Females
















## "generally available" Number of essential NCD technologies reported as





<sup>2017</sup> 6 out of 6

â