

MEXICO





PROPORTIONAL MORTALITY 24% 22% Cardiovascular Other NCDs diseases 12% 10% Cancers Communicable, maternal, perinatal and nutritional conditions Chronic respiratory diseases 10% 15% NCDs are estimated to account for 80% of all deaths.

			-								
97 400 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"											
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL				
MORTALITY											
Ä	Premature mortality from NCDs	x	Total NCD deaths	2016	272 700	249 000	521 800				
			Risk of premature death between 30-70 years (%)	2016	17	14	16				
K.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	5				
RISK FACTORS											
	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	11	2	7				
K	Physical inactivity	X	Physical inactivity, adults aged 18+ (%)	2016	25	32	28				
	Salt/Sodium intake	X	Mean population salt intake, adults aged 20+ (g/day)	2010	7	7	7				
8	Tobacco use	X	Current tobacco smoking, adults aged 15+ (%)	2016	22	7	14				
	Raised blood pressure	X	Raised blood pressure, adults aged 18+ (%)	2015	21	17	19				
À	Diabetes	X	Raised blood glucose, adults aged 18+ (%)	2014	10	11	10				
i i	Obesity	X	Obesity, adults aged 18+ (%)	2016	24	33	28				
			Obesity, adolescents aged 10-19 (%)	2016	15	12	14				
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2				
各	Household air pollution	-	Population with primary reliance on polluting fuels and technologies $(\!\%\!)$	2016	-	-	15				
SELECTED ADULT RISK FACTOR TRENDS											



NATIONAL SYSTEMS RESPONSE





		√	Proportion of population at high risk for CVD or with existing CVD (%)	-	
	Drug therapy to prevent heart attacks and strokes		Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-	
			Proportion of primary health care centres reported as offering CVD risk stratification	2017	Less than 25%
			Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017	Yes
	Essential NCD medicines and basic technologies	✓	Number of essential NCD medicines reported as "generally available"	2017	9 out of 10
	to treat major NCDs		Number of essential NCD technologies reported as "generally available"	2017	6 out of 6