




		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORT	TALITY [†]						
	Premature mortality	✓	Total NCD deaths	2016			
	from NCDs	V	Risk of premature death between 30-70 years (%)	2016			
<u>r</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK	FACTORS						
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	11	2	6
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	39	52	46
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010			
⊗	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	37	43	40
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	23	17	20
į.	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	30	28	29
673	Obesity	✓	Obesity, adults aged 18+ (%)	2016	59	63	61
₹			Obesity, adolescents aged 10-19 (%)	2016	32	32	32
9	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
in the	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	9

SELECTED ADULT RISK FACTOR TRENDS





Males Females



NATIONAL	







....o

....0



Number of essential NCD medicines reported as "generally available"
Number of essential NCD technologies reported as "generally available"

