
## **MARSHALL ISLANDS**

NO DATA AVAILABLE NO DATA AVAILABLE


		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY <sup>†</sup>							
	Premature mortality from NCDs	<b>√</b>	Total NCD deaths	2016			
			Risk of premature death between 30-70 years (%)	2016			
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK	ACTORS						
快	Harmful use of alcohol	X	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016			
X	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	41	54	47
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	7	6	6
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016			
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	23	18	20
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	20	21	21
À	Obesity	X	Obesity, adults aged 18+ (%)	2016	48	57	52
			Obesity, adolescents aged 10-19 (%)	2016	23	25	24
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)°	2016	-	-	-
h	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	35
ELEC	TED ADULT RISK FACT	OR TRENDS					

## NO DATA AVAILABLE

**CURRENT TOBACCO SMOKING** 




**OBESITY** 





## **RAISED BLOOD PRESSURE**









to treat major NCDs

